

Zeitschrift: Bulletin de la Société Neuchâteloise des Sciences Naturelles
Herausgeber: Société Neuchâteloise des Sciences Naturelles
Band: 116 (1993)

Artikel: The spider fauna of the Karatau mountain [i.e. mountain] range
Autor: Zyuzin, A.A. / Tarabaev, Chingis / Fyodorov, A.A.
DOI: <https://doi.org/10.5169/seals-89395>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The spider fauna of the Karatau mountain range

par

A.A. Zyuzin, Chingis Tarabaev & A.A. Fyodorov

The Karatau (or the Syr Darya Karatau) mountain range is the extreme north-western spur of Tien Shang and has its length above 400 km. Thanks to the unique geographical situation of Karatau, strongly wedging in the desert zone, and its considerable geological age the Karatau range has a number of endemic insects (PRAVDIN 1978): among mammals the endemic Karatau mountain sheep occur here. As to the endemic plants, the Karatau mountain range takes the first place among the other floristic areas of Kazakhstan (GVOZDEV, ANDREJTCHUK 1988).

The present paper includes the results of the determination of spiders collected by the authors in 1988-1989.

The spider fauna of the Karatau up to the last years was an absolute white spot: only recently MARUSIK and LOGUNOV (1990) registered here 6 species of the family Thomisidae from our collections; all these are presented in the Table 1.

During 1988-1989 we collected 110 species of spiders belonging to 83 genera from 24 families (Table 1). The representatives of the families Oecobiidae, Filistatidae, Oonopidae, as well as 10 genera and 17 species, are new for the Kazakhstan fauna: the last make up 15,5 per cent of the total number. Non-determined specimens designated as "sp" (44 species from 17 families, or 40,0 %) presumably are new species.

The most of the species collected (69, or 62,7 %) are rare (isolated) ones. At the same time, some species are usual and constant (31, or 28,2%: see Table 1). The dominating species are the four ones: *Mogrus antonius* (Salticidae), *Agelena orientalis* (Agelenidae), *Pardosa turkestanica* (Lycosidae), *Neoscona adianta* (Araneidae).

Table 1.- Species composition and relative abundance of spiders at Karatau mountain range

Families, species		Relative abundance
	*1. Oecobiidae	
K*	1 <i>Ambika nadiae</i> (Spas.)	+
	2. Eresidae	
	2 <i>Eresus niger</i> (Pet.)	+
K	3 <i>Stegodyphus lineatus</i> (Latr.)	++
	*3. Filistatidae	
K*	4 <i>Pritha crosbyi</i> (Spas.)	++
	*5 <i>Zaitunia</i> sp.	+
	4. Dictynidae	
	6 <i>Archaeodictyna</i> sp.	+
	7 <i>Brigittae latens</i> (Fabr.)+	
K	8 <i>B. innocens</i> (O.P.-C)	+
	*9 <i>Nigma</i> sp.	+
	10 <i>Lathys puta</i> (O.P.-C.)	++
	5. Titanoecidae	
	11 <i>Titanoeca veteranica</i> Herm.	+
	12 <i>Nursicia albosignata</i> Sim.	+
	6. Uloboridae	
	13 <i>Uloborus walckenaerus</i> Latr.	+
	7. Pholcidae	
K	14 <i>Pholcus ponticus</i> Thor.	++
	8. Palpimanidae	
	15 <i>Palpimanus</i> sp.	+
	*9. Oonopidae	
K*	16 <i>Dysderina loricata</i> (Sim.)	+
	10. Salticidae	
	17 <i>Synageles ramitus</i> Andr.	+
	18 <i>Pellenes seriatus</i> (Thor.)	+
	19 <i>Aelurillus ater</i> (Kroneb.)	++
	20 <i>Philaeus chrysps</i> (Poda)	++
	21 <i>Mogrus antoninus</i> Andr.	+++
	22 <i>Phlegra sogdiana</i> Charit.	+
	23 <i>Ph.sp.</i>	+
	24 <i>Chalcoscirtus</i> sp.	++
	25 <i>Bianor</i> sp.	+

Families, species	Relative abundance
26 <i>Euophrys nigrita</i> (Thor.)	+
27 <i>E.sp.</i>	++
28 <i>Heliophanus potanini</i> Schenk.	++
11. Gnaphosidae	
29 <i>Gnaphosa sp. 1</i>	+
30 <i>G.sp.2</i>	+
31 <i>G.sp.3</i>	+
K 32 <i>Leptodrassus memorialis</i> Spas.	+
33 <i>Drassodes sp.</i>	+
34 <i>Haplodrassus sp.</i>	+
35 <i>Synaphosus turanicus</i> Ovt.	++
36 <i>Talanites sp.</i>	+
K *37 <i>Phaeoedus braccatus</i> (L.Koch)	+
K 38 <i>Ph. rufescens</i> (Kroneb.)	+
39 <i>Zelotes jaxartensis</i> (Kroneb.)	+
40 <i>Z. praeficus</i> (L.Koch)	++
41 <i>Z.sp.1</i> (pr.puritanus)	+
42 <i>Z.sp.2</i> (pr.atrocoeruleus)	+
K 43 <i>Aphantaulax seminigra</i> Sim.	+
K 44 <i>Micaria albimana</i> O.P.-C	+
45 <i>M. formicaria</i> (Sund.)	++
K 46 <i>M. septempunctata</i> O.P.-C	+
12. Philodromidae	
47 <i>Philodromus sp.1</i>	+
48 <i>Ph.sp.2</i>	+
49 <i>Thanatus sp.1</i>	+
50 <i>Th.sp.2</i>	++
13. Thomisidae	
51 <i>Thomisus onustus</i> Walck.	+
52 <i>Runcinia tarabayevi</i> Marus., Log.	+
53 <i>Xysticus cristatus</i> (Cl.)	+
54 <i>X.sabulosus</i> (Hahn)	++
55 <i>S.turanicus</i> Charit.	+
56 <i>X.turlan</i> Marus., Log.	+
14. Clubionidae	
57 <i>Chiracanthium sp.1</i>	+
58 <i>Ch.sp.2</i>	++
K *59 <i>Phrurolithus pullatus</i> Kulcz.	++
15. Liocranidae	

Families, species	Relative abundance
*60 <i>Mesiotelus</i> sp.	++
*61 <i>Castianeira</i> sp.1	++
62 <i>C.</i> sp.2	+
16. Theridiidae	
63 <i>Latrodectus tredecimguttatus</i> (Ros.)	++
64 <i>Steatoda paykulliana</i> (Walck.)	++
65 <i>Lithyphantes albomaculatus</i> (De Geer)	++
66 <i>Teutana triangulosa</i> (Walck.)	++
67 <i>Theridion varians</i> Hahn	++
68 <i>Enoplognatha ovata</i> (Cl.)	+
69 <i>E.</i> sp.	++
70 <i>Euryopis</i> sp.	+
71 <i>Robertus</i> sp.	+
17. Oxyopidae	
72 <i>Oxyopes</i> sp.	+
18. Zodariidae	
73 <i>Zodarion</i> sp.1	++
74 <i>Z.</i> sp.2	+
75 <i>Z.</i> sp.3	+
19. Agelenidae	
K 76 <i>Agelena orientalis</i> C.L.Koch	+++
K 77 <i>A.marakandensis</i> (Charit.)	+
78 <i>Coelotes</i> sp.	+
20. Hahniidae	
79 <i>Hahnia</i> sp.1	+
80 <i>H.</i> sp.2	++
21. Lycosidae	
81 <i>Lycosa nordmanni</i> (Thor.)	++
82 <i>L.</i> sp.	++
83 <i>Allohogna singoriensis</i> (Laxam.)	+
84 " <i>Geolycosa</i> " sp.1	+
85 " <i>G.</i> " sp.2	+
86 <i>Alopecosa</i> sp.	+
87 <i>Arctosa leopardus</i> (Sund.)	++
88 <i>Pirata</i> sp.	++
89 <i>Evippa</i> sp.	++
90 <i>Pardosa turkestanica</i> (Roew.)	+++
22. Araneidae	
91 <i>Argiope lobata</i> (Pall.)	++

Families, species	Relative abundance
92 <i>Larinioides folium</i> (Schr.)	++
93 <i>Agalenatea redii</i> (Scop.)	+
94 <i>Aculepeira</i> sp.	+
95 <i>Neoscona adianta</i> (Walck.)	+++
96 <i>Mangora acalypha</i> (Walck.)	+
97 <i>Zygiella caspica</i> (Sim.)	++
23. Tetragnathidae	
98 <i>Tetragnatha extensa</i> (L.)	++
99 <i>Pachygnatha clercki</i> Sund.	+
24. Linyphiidae	
100 <i>Microlinyphia pusilla</i> (Sund.)	++
101 <i>Agyneta</i> (<i>Agyneta</i>) sp.	+
102 <i>A. (Meioneta) fuscipalpis</i> (C.L.Koch)	+
K 103 <i>Lepthyphantes kuhitangensis</i> Tanas.	+
K 104 <i>L. tchatkalensis</i> Tanas.	+
105 <i>Bolyphantes indexoides</i> Tanas.	+
106 <i>Erigone dentipalpis</i> (Wid., Reuss.)	+
107 <i>Oedothorax apicatus</i> (Blackw.)	+
108 <i>Ceratinopsis romana</i> (O.P.-C.)	+
109 <i>Mesasigone mira</i> Tanas.	+
*110 <i>Trachelocamptus</i> sp.	+

Total: 110 species, 24 families, 83 genera

- + - rare (isolated) species
- ++ - usual species
- +++ - dominating species
- * - families and genera new for Kazakhstan
- K - species new for Kazakhstan

The most numerous as to the number of species are the families Gnaphosidae (18 species, or 16,4%), Salticidae (12-10,9), Linyphiidae (11-10,0), Oycosidae (10-9,1), Theridiidae (9-8,2), Araneidae (7-6,4), Thomisidae (6 species, or 5,5 per cent), and as to the generic diversity the families Gnaphosidae and Salticidae (10 Genera), Linyphiidae (9), Lycosidae and Theridiidae (in 8), Araneidae (7 genera).

The main part of the species collected constitute geobionts (the spiders living in soil and on the ground surface): these are 84 species, or 76,4 per cent of the total number of species - the representatives of the families Titanoecidae, Pholcidae, Palpimanidae,

Oonopidae, Salticidae (except the genera *Synageles*, *Mogrus*, *Heliophanus*), Gnaphosidae (except *Aphantaulax*), Hahniidae, Lycosidae, Linyphiidae (except *Microlinyphia*) and the genera *Eresus* (Eresidae), *Zaitunia* (Filistatidae), *Lathys* (Dictynidae), *Xysticus* (Thomisidae), *Phrurolithus* (Clubionidae), *Zygiella* (Araneidae), *Pachygnatha* (Tetragnathidae). The most of geobionts are the species hiding from the sunrays under stones: these are the representatives of above mentioned families and genera, except the families Salticidae, Philodromidae and genera *Castianeira* (Liocranidae), *Agelena* (Agelenidae), *Pardosa* (Lycosidae), active under bright light. The separate group of geobionts constitute the species burrowing deep holes - trypanobionts (trupa-the hole): these are the species of the genera *Lycosa*, *Allohogna* and "Geolycosa" (Lycosidae). Some geobionts are strictly limited to their specific habitats (stenobionts). Thus, *Pardosa turkestanica* (Lycosidae) lives among stones and in a grass along the banks of streams. The lycosids *Arctosa leopardus* and *Pirata* sp. prefer marshy places. The species *Evippa* sp. (Lycosidae), living on screes, and *Xysticus turanicus* (Thomisidae), are typical petrobionts having uniform grey colour (Araneidae) builds its webs on the cracked rocks. *Castianeira* spp. (Liocranidae) live only in the *Scyrcus* along the streams.

Phytobionts (plant inhabitants) are basically represented by the hortobionts (grass inhabitants): these are the species of the genera *Archaeodictyna*, *Brigittea* (Dictynidae), *Synageles*, *Heliophanus* (Salticidae), *Aphantaulax* (Gnaphosidae), *Thomisus*, *Runcinia* (Thomisidae), *Chiracanthium* (Clubionidae), *Oxyopes* (Oxyopidae), *Larinioides*, *Agalenatea*, *Neoscona*, *Mangora* (Araneidae), *Tetragnatha* (Tetragnathidae), *Microlinyphia* (Linyphiidae). The inhabitants of bushes and low trees (thamnobionts) are represented by the genera *Uloborus* (Uloboridae), *Stegodyphus* (Eresidae), *Mogrus* (Salticidae), *Argiope*, *Aculepeira* (Araneidae).

The species *Pritha crosbyi* (Filistatidae), inhabiting the wall cracks, and *Ambika nadiae* (Oecobiidae), are synanthropic ones.

Acknowledgements: For the aid in the determination of our material and for the participation in spider collections we are grateful to the following persons: P.M. Dunin, S.L. Zonshtein, L.I. Komarova, D.V. Logunov, Y.M. Marusik, K.G. Mikhailov, V.I. Ovtsharenko, S.V. Ovtchinnikov, A.V. Tanasevitch.

REFERENCES

- GVOZDEV, E.V. & ANDREJTCHUK, A.L. (ed.-in-chief) - (1988). The new reserves of Kazakhstan. "Nauka" Publ., Alma-Ata, 108 pp. (in russian).
- MARUSIK, Y.M. & LOGUNOV, D.V. - (1990). The crab spiders of Middle Asia, USSR (Aranei, Thomisidae). 1. Descriptions and notes on distribution of some species. *Korean Arachnol.*, v. 6, N 1, p. 31-62 (in english).
- PRAVDIN, F.N., 1978. An ecological geography of Central Asian insects. *Orthopteroids*. Moscow, "Nauka" Publ., 271 pp. (in russian).

A.A. ZYUZIN, Chingis TARABAEV, A.A. FYODOROV
Institute of Zoology, Kazakhstan Academy of Sciences
CEI-480032 ALMA-ATA