

**Zeitschrift:** Bulletin de la Société Neuchâteloise des Sciences Naturelles  
**Herausgeber:** Société Neuchâteloise des Sciences Naturelles  
**Band:** 120 (1997)

**Artikel:** Notes on Chionea (Sphaeonophilus) species in Central Europe (Diptera ; Limoniidae)  
**Autor:** Reusch, Herbert  
**DOI:** <https://doi.org/10.5169/seals-89470>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 29.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## NOTES ON *CHIONEA (SPHAECONOPHILUS)* SPECIES IN CENTRAL EUROPE (DIPTERA; LIMONIIDAE)

HERBERT REUSCH

Wellendorf 70, 29562 Suhlendorf, Deutschland.

### Summary

New data about unstudied material of *Chionea (Sphaeconophilus) belgica* (Becker, 1912) and *C. (S.) lutescens lutescens* Lundström, 1907, collected in Germany and Switzerland are reported, and material from Denmark is revised. For the first time there is evidence for *C. belgica* from Denmark, Bavaria and Thuringia (Germany) and Switzerland. New discoveries are known from altitudes (above sealevel) between 10 m (Copenhagen) und 835 m (Milseburg/Rhön, Hessa). Until these were made, no specimens had been found above 500 m.

*C. lutescens* has to be removed from the Danish fauna, because this species had been mistaken for *C. belgica*, whereas *C. lutescens* has to be added to the fauna of the northern German lowlands (alt. below 100 m; formerly none had been found below 340 m) and to the mountains of the "Harz" and the "Thüringer Wald".

### INTRODUCTION

In Central Europe 10 species of the genus *Chionea* Dalman, 1816, are known (SAVCHENKO *et al.*, 1992). These can be distinguished using the keys written by KRZEMIŃSKI (1982) and GROOTAERT (1984). Until GROOTAERT (1984) redescribed *C. (Sphaeconophilus) belgica* (Becker, 1912), *C. (S.) belgica* and *C. (S.) lutescens lutescens* Lundström, 1907, had been confused. So older material has to be revised to obtain a survey of the actual spreading of both species (MENDL & REUSCH, 1989; REUSCH, 1988).

The following data are the result of the revision of old collections as well as of the study of newly dealt with material from Central Europe.

### RESULTS AND DISCUSSION

#### *Chionea (Sphaeconophilus) belgica* (Becker, 1912)

##### Denmark

For Denmark EDWARDS (1936) reported individuals of "*C. lutescens*", which had been found by G. P. KRYGER during the winter of 1934-35, in Gentofte near Copenhagen (alt. 10 m) and sent to the "British Museum (Natural History)" at London for identification (see also NIELSEN, 1941). This occurred again in Nov. 1935 when KRYGER found several pupae and adults just as the year before while digging his carrots as well as another collector Mr KLÖCKER.

Revising the material of the British Museum, REUSCH realized that these ani-

mals were actually *C. belgica*, formerly unknown in Denmark. This discovery was only reported to the authors of the "Catalogue of Palaearctic Diptera" (SAVCHENKO *et al.*, 1992).

The pupae of the *Chionea*-species were unknown when EDWARDS (1936) gave a rough description of what was really *C. belgica*. BYERS (1961, 1983) detailed these descriptions using the same animals which he had been borrowed from the British Museum as "*C. lutescens*".

On demand for *Chionea*-material at the "Zoologisk Museum" at Copenhagen REUSCH received individuals which were labeled "*C. lutescens*" from the NIELSEN-collection (1 ♂, 2 ♀♀, "18.11.34, det. PEDER NIELSEN"). These refered to the same material from KRYGER and KLÖCKER and belong to *C. belgica* as well.

## Germany

The existence of *C. belgica* has been proved for the first time in the hilly region of southern Lower Saxony and eastern Hessa by REUSCH, 1988 (see also HÖVEMAYER, 1996, WEBER & WEIDEMANN, 1993), and other evidences dating from different periods exist from both regions of the central low mountain range. This species appeared for the first time in ground-traps in North-Rhine-Westphalia during a research project of the "Bundesamt für Naturschutz" (RIECKEN & RIES, 1993a, 1993b). *C. belgica* is also new for Thuringia where REUSCH discovered it on checking several collections.

Until now *C. belgica* was unknown in Bavaria, *C. lutescens* however had been found in the "Allgäu" north of the Alps (MENDL & REUSCH, 1989). Recently BUCK (1997) found several specimens in ground-traps as well as in traps that were baited with carcasses (kidney and dead *Helix pomatia*). Even though the access openings of the traps were closed with gaze (3

x 3 mm), the relatively long-legged midges passed through without problems.

BAUMANN (1977), BYERS (1983) and NABAGŁO (1973) had observed necrophily with other *Chionea*-species, whereas none of this behaviour had been reported about *C. belgica*.

MATERIAL (COLLECTED WITH GROUND-TRAPS, IF NO FURTHER INFORMATION IS GIVEN):

**Bavaria:** Environs of Senden-Hittistetten, Landkreis Neu-Ulm; alt. 487 m; 14.10.1992 - 13.01.1993; 42 ♂♂ 33 ♀♀; leg. et coll. M. Buck; traps baited with carcasses and ground-traps.

**Hessia:** Schlitz; 220 - 250 m; 15.11.1969; 4 ♂♂ 5 ♀♀ (traps baited with carcasses); 28.10.1977; 2 ♂♂ 2 ♀♀; leg. et coll. P. Zwick.

**Hessia:** Milseburg, Rhön; 835 m; 1 ♂; 28.10.1979; leg. et coll. P. Zwick; traps baited with cheese.

**Hessia:** Obersberg, Bad Hersfeld; 250 m; Nov.-Dec. 1980; 38 ♂♂ 26 ♀♀; leg. V. Puthz, coll. P. Zwick.

**Lower Saxony:** Steinberg near Bad Salzdetfurth, Landkreis Hildesheim; 160 - 180 m; 25 ♂♂ 34 ♀♀; 7.11. - 22.12.1996; leg. L. Schmidt & P. Sprick, coll. H. Reusch.

**North-Rhine-Westphalia:** Drachenfels Ländchen, south of Bonn; 165 - 180 m; 3.-17.12.1992; leg. U. Riecken, coll. H. Reusch.

**Thuringia:** Ohrdruf, Thüringer Wald; 400 m; Oct. 1889; 1 ♂; leg. Dr. Thomas, coll. Zoologisches Institut, Martin-Luther-Universität Halle/Wittenberg.

**Thuringia:** Haßlebener Ried, Landkreis Sömmerda; 147 m; 20.12.1983 - 3.01.1984; 2 ♂♂ 1 ♀; leg. H. Grimm, coll. "Museum der Natur" Gotha.

**Thuringia:** Herrenschwende, Landkreis Sömmerda; 142 m; 8.11.1983 - 17.01.1984; 18 ♂♂ 15 ♀♀; leg. H. Grimm, coll. "Museum der Natur" Gotha.

**Thuringia:** Leutratal, 8 km south of Jena; 250 - 260 m; 17.01.1989; 5 ♂♂ 1 ♀; leg. et coll. R. Bährmann.

## Switzerland

From the Sensetal near Flamatt (alt. 490 - 530 m), south-west of Bern, BANGERTER (1939) reported "*Chionea lutescens* Lundström. 20.XII.37. Selten." The Bangerter-material was revised by MENDL (1979: 353) who confirmed the identification and labeled the accompanying micro-preparation "Nr. 186".

During an inspection of larvae and pupae of the borrowed BANGERTER-collection REUSCH discovered that the previously mentioned preparation belongs to *C. belgica* and not to "*lutescens*". The alcoholic preserved material contains another 2 ♂♂, labeled "*Chionea*. Flamatt XII.37". Therefore BANGERTER (1939) discovered the furthest south population of this species in Europe.

Other than that *C. belgica* is only known to exist in Belgium (GROOTAERT, 1984) and the Netherlands (HEIJERMANN, 1987; the material was revised and confirmed by REUSCH). It will be uncertain if the whole spreading-area has been registered (fig. 1), until *Chionea*-material from other countries has been revised (for distribution-data see SAVCHENKO *et al.*, 1992).

### *Chionea (Sphaeconophilus) lutescens* *lutescens* Lundström, 1907

## Denmark

There is no proof of the existence of *C. lutescens* because the material mentioned in previous publications (BYERS, 1961, 1983; EDWARDS, 1936; NIELSEN, 1941) belongs to *C. belgica* without exception.

## Germany

The evidence of *C. lutescens* could be proved in the northern German lowlands of Lower Saxony for the first time. She was caught with ground-traps near Lehrte, east of Hannover, as well as in the nature reserve "Lüneburger Heide". These places are both situated below 100 m, that is




Figure 1: Distribution of *Chionea belgica* according to revised material.

considerably lower than any previous known places where *C. lutescens* was found before (alt. 340 - 1700 m; WEBER & WEIDEMANN, 1993).

Other regions where material was found and where the species was unknown until then (Mendl & Reusch, 1989) are the "Brocken" in the "Nationalpark Hochharz" (Landesamt für Umweltschutz Sachsen-Anhalt, 1994) as well as the "Thüringer Wald".

## MATERIAL:

**Lower Saxony:** Environs of Lehrte, 10 km east of Hannover; alt. 60 m; 15.11.1991 - 15.01.1992; 4 ♂♂ 10 ♀♀; leg. W. Wölkerling, coll. A. Melber.

**Lower Saxony:** Nature Reserve "Lüneburger Heide" near Oberhaverbeck, north-west of Behringen, Landkreis Soltau-Fallingbostel; 90 m; 11.1994 - 01.1995; 2 ♂♂ 9 ♀♀; leg. L. Schmidt, coll. H. Reusch.

**Saxony-Anhalt:** Brocken, "Nationalpark Hochharz"; 1020 - 1080 m; 9.05.1996; 95 ♂♂ 88 ♀♀; leg. P. Sacher, coll. H. Reusch. 930 m; 9.05.1996; 2 ♂♂ 3 ♀♀; leg. P. Sacher, coll. R. Bährmann.

**Thuringia:** Frauengraben, Klein Schmalkalden, Thüringer Wald; 650 m; 12.12.1993; 2 ♂♂; leg. R. Borcherding, coll. R. Bellstedt.

Another report exists from the “Odenwald” in western Hessia from BÜRGIS (1992), who proved evidence of this species in knowledge of GROOTAERT’s work (1984). The proof of further records from Germany (MENDL & REUSCH, 1989) is uncertain for the time being until the material has been revised.

#### ACKNOWLEDGEMENTS

I owe the following persons and institutions a debt of gratitude for lending me their material to work, to keep, as well as their help with supplement data.

Material of public collection was provided by Ronald Bellstedt (Museum der Natur, Gotha), Dr. Charles Huber (Naturhistorisches Museum, Bern), Dr. Herman de Jong and Dr. W. Hogenes (Zoologisch

Museum, Universiteit van Amsterdam), Dr. Leif Lyneborg (Zoologisk Museum, Copenhagen), Dr. Karla Schneider and J. Händel (Zoologische Sammlung, Martin-Luther-Universität Halle/Wittenberg; arranged by Dr. Andreas Stark) and Dr. B.C. Townsend (British Museum - Natural History, London).

Material from their private collections were put to my disposal by Prof. Dr. Rudolf Bährmann (Universität Jena), Ronald Bellstedt (Gotha), Dr. Matthias Buck (Ulm), Dr. Albert Melber, Ludger Schmidt, Peter Sprick and Helfried Wölkerling (all of them Universität Hannover), Uwe Riecken (Bundesamt für Naturschutz, Bonn-Bad Godesberg), Dr. Peter Sacher (Nationalparkverwaltung, Blankenburg/Harz) as well as Prof. Dr. Peter Zwick (Max-Planck-Institut für Limnologie, Schlitz).

Last but not least I want to thank Yanna McLaughlin (Lüneburg) for translating the English manuscript.

#### REFERENCES

- BANGERTER, H. 1939. Die Eriopterini des Sensetales bei Flamatt-Neuenegg. *Mitt. Schw. Ent. Ges.* 17 (10): 482-487.
- BAUMANN, E. 1977. Untersuchungen über die Dipterenfauna subterrane Gangsystems und Nester von Wühlmäusen (*Microtus*, *Clethrionomys*) auf Wiesen der montanen Region im Naturpark Hoher Vogelsberg. *Zool. Jb., Abt. Syst. Ökol. Geogr. Tiere* 104: 368-414.
- BUCK, M. 1997. Untersuchungen zur ökologischen Einnischung saprophager Dipteren unter besonderer Berücksichtigung der Phoridae und Sphaeroceridae (Brachycera/Cyclorrhapha). *Dissertation, Universität Ulm (in press)*.
- BÜRGIS, H. 1992. Die Winterinsekten vom Heidenberg bei Lautertal-Raidelbach (nordwestlicher Odenwald). *Naturwiss. Ver. Darmstadt, Ber. (N.F.)* 15: 41-54.
- BYERS, G.W. 1961. Biology and classification of *Chionea* (Diptera: Tipulidae). XI. *Int. Kongr. Entomol. (Wien)* 1960 (1): 188-191.
- BYERS, G.W. 1983. The crane fly genus *Chionea* in North America. *Univ. Kansas Sci. Bull.* 52 (6): 59-195.
- EDWARDS, F.W. 1936. Pupae and living adults of *Chionea lutescens* Lundström from Denmark. *Proc. R. Ent. Soc. London* 10: 89-91.

- GROOTAERT, P. 1984. Redescription of *Niphadobata belgica* (Becker, 1912) comb. nov. (Diptera: Tipulidae), a snowfly from low altitudes in Belgium. *Annls Soc. r. zool. Belg.* 114 (2): 241-247.
- HEIJERMAN, T. 1987. De sneeuwmug *Niphadobata belgica* (Becker, 1912) in Nederland (Diptera: Tipulidae). *Ent. Ber., Amst.* 47 (8): 113-118.
- HÖVEMEYER, K. 1996. Die Dipterengemeinschaft eines Erlenuferwaldes in Südniedersachsen. *Braunschw. naturkdl. Schr.* 5 (1): 71-84.
- KRZEMIŃSKI, W. 1982. Contributions to the taxonomy of the European species of *Chionea* Dalman (Diptera: Limoniidae). *Ent. scand.* 13: 193-200.
- LANDESAMT FÜR UMWELTSCHUTZ SACHSEN-ANHALT, ABTEILUNG NATURSCHUTZ (ED.). 1994. Der Nationalpark Hochharz. *Naturschutz im Land Sachsen-Anhalt* 31 (Sonderh.): 1-72.
- MENDL, H. 1979. Revision der Limoniidae-Sammlung von Hans Bangerter im Naturhistorischen Museum zu Bern/Schweiz. *Beitr. Ent., Berlin* 29 (2): 343-372.
- MENDL, H. & REUSCH, H. 1989. Liste der aus der Bundesrepublik Deutschland bis jetzt bekannten Stelzmücken (Diptera Nematocera: Limoniidae). *Braunschw. naturkdl. Schr.* 3 (2): 429-453.
- NABAGŁO, Ł. 1973. Participation of invertebrates in decomposition of rodent carcasses in forest ecosystems. *Ekologia Polska* 21: 251-270.
- NIELSEN, P. 1941. Danmarks Stankelben. Tillæg og Rettelser til "Danmarks Fauna", Bd. 28. *Flora og Fauna* 1941: 81-91.
- REUSCH, H. 1988. *Niphadobata belgica* (Becker, 1912) neu für Deutschland (Diptera: Limoniidae). *Ent. Z.* 98 (20): 300-304.
- RIECKEN, U. & RIES, U. 1993a. Zur Bedeutung naturnaher Bachufer und Brachen in der Zivilisationslandschaft am Beispiel der Laufkäfer (Coleoptera: Carabidae). *Mitt. Dtsch. Ges. Allg. Angew. Ent.* 8 (4-6): 397-403.
- RIECKEN, U. & RIES, U. 1993b. Zur Bewertung und Bedeutung naturnaher Landschaftselemente in der Agrarlandschaft. Teil II: Laufkäfer (Coleoptera: Carabidae). *Verh. Ges. Ökol.* 22: 241-248.
- SAVCHENKO, E.N., OOSTERBROEK, P. & STARÝ, J. 1992. Family Limoniidae. - In: Soós, Á., PAPP, L. & OOSTERBROEK, P. (eds.): Catalogue of the Palaearctic Diptera 1: 183-374. Ed. Hungarian natural History Museum. Budapest.
- WEBER, G. & WEIDEMANN, G. 1993. *Niphadobata lutescens* (Lundström, 1907) and *N. belgica* (Becker, 1912) from the Solling mountains, Lower Saxony, FRG (Diptera, Limoniidae). *Dtsch. ent. Z. (N.F.)* 40: 259-263.