

Zeitschrift: Bulletin de la Société Vaudoise des Sciences Naturelles
Herausgeber: Société Vaudoise des Sciences Naturelles
Band: 88 (2002-2003)
Heft: 4

Artikel: Two negative ¹³Corg excursions near the Triassic-Jurassic boundary in the New York Canyon area (Gabbs Valles Range, Nevada)
Autor: Guex, Jean / Bartolini, Annachiara / Atudorei, Viorel
DOI: <https://doi.org/10.5169/seals-281438>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Two negative $\delta^{13}\text{C}_{\text{org}}$ excursions near the Triassic-Jurassic boundary in the New York Canyon area (Gabbs Valley Range, Nevada)

by

Jean GUEX¹, Annachiara BARTOLINI², Viorel ATUDOREI³
and David TAYLOR⁴

Abstract.—GUEX J., BARTOLINI A., ATUDOREI V. and TAYLOR D., 2003. Two negative $\delta^{13}\text{C}_{\text{org}}$ excursions near the Triassic-Jurassic boundary in the New York Canyon area (Gabbs Valley Range, Nevada). *Bull. Soc. vaud. Sc. nat.* 88.4: 445-448. In this paper we present a new Late Rhaetian (Crickmayi Zone) to Early Jurassic (Polymorphum-Mulleri Zones) $\delta^{13}\text{C}_{\text{org}}$ curve established in New York Canyon (Gabbs Valley, Nevada) and discuss its significance for worldwide correlations. Our curve shows two significant negative shifts of the organic carbon which are correlated to the very precise biochronological data that we have established and published during the last few years in several reports (TAYLOR *et al.* 1999, TAYLOR and GUEX 2000, GUEX 1995, GUEX *et al.* 1998).

Keywords: Triassic Jurassic boundary, Carbon Isotopes, New York Canyon Nevada.

Résumé.—GUEX J., BARTOLINI A., ATUDOREI V. et TAYLOR D., 2003. Deux excursions négatives du $\delta^{13}\text{C}_{\text{org}}$ au voisinage de la limite Trias-Jurassic dans la région du New York Canyon (Gabbs Valley Range, Nevada). *Bull. Soc. vaud. Sc. nat.* 88.4: 445-448.

Nous présentons ici une nouvelle courbe du $\delta^{13}\text{C}_{\text{org}}$ allant du Rhétien supérieur (Zone à Crickmayi) au Jurassique inférieur (Zones à Polymorphum-Mulleri) établie dans la région du New York Canyon (Gabbs Valley Range, Nevada). Sa signification pour les corrélations à grande échelle est discutée. Cette courbe montre deux anomalies négatives significatives du carbone organique. Celles-ci sont précisément calibrées aux données biochronologiques que nous avons établies et publiées au cours des dernières années (TAYLOR *et al.* 1999, TAYLOR and GUEX 2000, GUEX 1995, GUEX *et al.* 1998).

Mots clés: Limite Trias Jurassique, Isotopes du Carbone, New York Canyon Nevada.

¹Dept. of Geology, Univ. of Lausanne, BFSH2, CH-1015 Lausanne, Switzerland

²Laboratoire de Micropaléontologie, UPMC, Paris, France

³Dept. of Geochemistry, University of New Mexico, USA

⁴NorthWest Museum of Natural History, Portland, Oregon, USA

INTRODUCTION

The goal of this paper is to present new data on the $\delta^{13}\text{C}_{\text{org}}$ variations in the New York Canyon area (Gabbs Valley, Nevada) and discuss its significance for worldwide correlations (Fig.1). Our curve shows the variation of the $\delta^{13}\text{C}_{\text{org}}$ from the Late Rhaetian (Crickmayi Zone) up to the Early Jurassic (Polymorphum-Mulleri Zones). It shows two significant negative shifts of the organic carbon which are correlated with the very precise biochronological data that we have established and published during the last few years in several reports (TAYLOR *et al.* 1999, TAYLOR and GUEX 2000, GUEX 1995, GUEX *et al.* 1998, 2002).

DISCUSSION

The beginning of the first excursion is located approximately at the top of the last occurrence of relatively abundant Rhaetian ammonites like *Arcestes nevadanus*, *A. gigantogaleata*, *Placites*, *Choristoceras crickmayi*, *C. marshi*, and *Rhacophyllites* spp. This negative excursion terminates 4 meters below the first occurrence of undisputable Jurassic ammonites belonging to the genus *Psiloceras*, i.e. *P.spelae* and *P. tilmanni* (bed N-8).

The second negative excursion starts near the first occurrence (bed M-2) of *Odoghertyceras* and *Psiloceras marcouxii*. These forms are accompanied by rare and very tiny survivors of the genus *Choristoceras*, i.e. *C.minutum*.

It is interesting to note that the first jurassic bivalves, represented by *Kalentera* and *Palmoxytoma*, first occur just above the last typical Triassic ammonites.

In this paper, we leave open the problem of the formal definition of the base of the Jurassic: it can be drawn either at the FO of the Jurassic bivalves or at the FO of the oldest Jurassic ammonites (i.e. *P.tilmanni*: see GUEX *et al.* 1998).

Three interesting $\delta^{13}\text{C}_{\text{org}}$ local curves have been published in recent years: HESSELBO (2002) in southern England, PALFY (2001) in Hungary and WARD *et al.* (2001) in western Canada.

Correlating these curves with our new isotopic data is quite obvious except for the third one.

The most complete $\delta^{13}\text{C}_{\text{org}}$ sequence around the TJB is that established by HESSELBO *et al.* (2002). The curve shows clearly two negative excursions: the first one is located within the range of the last conodonts occurring in the Cotham Member of the Lilstock Formation: it is here assigned to the Rhetian. The second excursion starts a few meters below the first local occurrence of *Psiloceras planorbis* and is interpreted as correlative of our second excursion




Figure 1.—Lithology, $\delta^{13}C_{org}$ variation and biochronology. A: Extinction of Triassic ammonites. B: First local occurrence of Jurassic Bivalves. C: First occurrence of *Psiloceras tilmanni*. Data from TAYLOR *et al.* 1999, TAYLOR and GUEX 2000 GUEX 1995, GUEX *et al.* 1998.

which is of lowermost Jurassic age. We note briefly that the Rhenium and Osmium anomalies described by COHEN and COE (2002) in the same section are clearly Jurassic and not Rhaetian.

The organic carbon curve published by PALFY *et al.* (2001) in Hungary also shows two negative excursions. The first one is located within the range of the last typical Rhaetian conodonts and is correlative with the first negative shift in England. A second negative shift is bracketed within a 20 meters stratigraphic interval located between a topmost Rhaetian (or lowermost Hettangian) *Rhacophyllites* and a *Kammerkarites* (our revised taxonomy). This second negative excursion correlates well with the second shift observed in New York Canyon, both in amplitude and chronologic position.

A third curve with only one negative shift was published by WARD *et al.* (2001) established at Kennecott Point (BC) in Canada. The biochronologic data accompanying that curve are not clear for two reasons. This negative shift overlaps the first Jurassic radiolarians and is supposedly located about 10 meters above the last occurrence of truly Triassic *Choristoceras*. This would

imply that the recorded excursion could be of Jurassic age. However we question the accuracy of these ammonite biostratigraphic data and suppose now that this shift corresponds to the first negative excursion in New York Canyon.

CONCLUSION

In summary our biostratigraphic observations and new isotope data show definitely two distinct negative carbon excursions close the TJB. The first one is clearly associated with the late-Triassic extinction and the second one, located in the Lower Hettangian, is concomitant with the major diversification of Jurassic ammonites, plants and radiolarians.

REFERENCES

- COHEN A.S. and COE A.L., 2002. New geochemical evidence for the onset of volcanism in the Central Atlantic magmatic province and environmental change at the Triassic-Jurassic boundary. *Geology (Boulder)* 30: 267-270.
- GUEx J., 1980. Remarques préliminaires sur la distribution stratigraphique des ammonites hettangiennes du New York Canyon (Gabbs Valley Range, Nevada). *Bull. Soc. vaud. Sc. nat.* 75.2: 127-140.
- GUEx J., 1995. Ammonites hettangiennes de la Gabbs Valley Range (Nevada). *Mémoires de Géologie, Lausanne* 27: 1-130.
- GUEx J., TAYLOR D., RAKUS M. and BUCHER H., 1998. Deux nouveaux genres et quatre nouvelles espèces d'ammonites (Cephalopoda) du Lias inférieur. *Bull. Soc. vaud. Sc. nat.* 86.1: 73-85.
- HESSELBO S.P., ROBINSON S.A., SURLYK F. and PIASECKI S., 2002. Terrestrial and marine extinction at the Triassic-Jurassic boundary synchronized with major carbon-cycle perturbation: a link to initiation of massive volcanism ? *Geology (Boulder)* 30: 251-254.
- PALFY J., DEMENY A., HAAS J., HETENYI M., ORCHARD M.J. and VETO I., 2001. Carbon isotope anomaly and other geochemical changes at the Triassic-Jurassic boundary from a marine section in Hungary. *Geology (Boulder)* 29: 1047-1050.
- TAYLOR D.G., SMITH P.L., LAWS R.A. and GUEx J., 1983. The stratigraphy and biofacies trends of the lower Mesozoic Gabbs and Sunrise formations, west-central Nevada. *Canad. J. Earth Sci.* 20: 1598-1608.
- TAYLOR D., BOELLING K. and GUEx J., 1999. The Triassic/Jurassic system boundary in the Gabbs Formation, Nevada, Symposium on Jurassic Stratigraphy, Vancouver, Vol. 6, Trans. Tech. Pub.:225-237.
- TAYLOR D. and GUEx J., 2002. The Triassic/Jurassic System boundary in the John Day Inlier, East-Central Oregon. *Oregon Geology* 64: 3-28.
- WARD P.D., HAGGART J.W., CARTER E.S., WILBER D., TIPPER H.W. and EVANS T., 2001. Sudden productivity collapse associated with the Triassic-Jurassic boundary mass extinction. *Science* 292: 1148-1151.

Manuscrit reçu le 29 septembre 2003

Remarque:

Cette note a fait l'objet d'un Numéro Spécial du Bulletin de Géologie, Lausanne (N° 360) paru en novembre 2003.