

Zeitschrift: Candollea : journal international de botanique systématique =
international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 17 (1959-1961)

Artikel: Three new species of Astragalus L. from West Pakistan

Autor: Ali, S. I.

DOI: <https://doi.org/10.5169/seals-880407>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Three new species of *Astragalus* L. from West Pakistan

by

S. I. ALI

(Departement of Botany, The University, Karachi, Pakistan)

Recently the present author published a paper (ALI, S. I. Notes on the genus *Astragalus* L. from West Pakistan and N.-W. Himalayas. *Kew Bull.* 1958: 303-318. 1958) wherein new and interesting taxa belonging to the genus *Astragalus* from West Pakistan and the N.-W. Himalayas have been discussed. Subsequently some more material was received for identification from the Conservatoire et Jardin botaniques, Geneva and from Gordon College, Rawalpindi by the courtesy of Dr. R. R. STEWART. A critical study of the material and the specimens available at the herbarium of Gordon College, Rawalpindi, revealed the existence of three new species. The material on which these new species are based was collected by F. SCHMID for the Swiss Entemological Expedition in 1953-1954 and by Mr. J. D. A. STAINTON, who led the British Museum Expedition to Chitral in 1958.

The present opportunity has been taken to describe these new species. The classification followed is the same as that discussed earlier by the author (ALI 1958). Further material of this genus is being studied and is expected to yield more interesting data.

Astragalus staintonianus Ali, spec. nov.

Herba perennis. Radix lignosa elongata. Caule super terram emergente valde abbreviato, internodiis quasi deficientibus; parte subterranea elongata. *Folia* 5-8,5 cm. longa, imparipinnata, stipulis c. 11 mm. longis, c. 3-4 mm. latis, glabris; rhachide c. 2-4 cm. longa. *Foliola* 25-43, petiolulo quam 1 mm. longiore, lamina c. 3-6 mm. longa, c. 2-4 mm. lata, elliptica, ovata, supra glabra (margine excepto), infra pilosa, margine integra, apice obtusa. *Inflorescentia* 4-flora, pedunculo 4-5 mm. longo. Flores pedicellis c. 3 mm. longis, bracteis c. 1 cm. longis, lineatis. *Calyx* extus pilosus, parte connata c. 9-10 mm. longa, duobus dentibus superioribus c. 6 mm., 3 dentibus inferioribus c. 5 mm. longis acutis. *Corolla* (in sicco) flava apicibus violascentibus, vexillo c. 19-22 mm. longo, c. 8-10 mm. lato, apice retuso; alis c. 19-21 mm. longis, c. 4 mm. latis, clave c. 11-12 mm. longa; carina c. 16 mm.



FIG. 13. — *Astragalus staintonianus* Ali, holotype (Stainton 2874, RAW).




FIG. 14. — *Astragalus imitensis* Ali, holotype (Schmid 2072, G).

longa, c. 4 mm. lata, clave c. 10-11 mm. longa. *Stamina* diadelpha, 9+1, filamentis c. 16-18 mm. longis. *Ovarium* stipitatum, c. 7-8 mm. longum, c. 2 mm. latum, dense albo-pilosum, stylus glaber c. 2 mm. ab apice angulo recto superne curvatus; stigmate capitato. Legumen sessile 17-18 mm. longum, c. 5-6 mm. latum, pilis albis hirsutum, partim biloculare, seminibus c. 8. Semina reniformia, c. 3-4 mm. longa, c. 2 mm. lata, colore brunneo.

HOLOTYPE: CHITRAL, Khot Au, alt. 4700 m., July 12, 1958, *Stainton* 2874 (RAW) (Fig. 13).

COTYPES: KASHMIR, Anesarbar, pierriers, 3900-4900 m., 15-16 août 1954, *Schmid* 2265 (G, RAW); deux versants de l'Ishkuman Aghost, 4210-4540 m, 15-16 août 1954, *Schmid* 2462 (G); Chitral, haute vallée de Yarkhun, hauts pâturages, 4000 m., 22-27 août 1954, *Schmid* 2327 (G).

This species belongs to the section *Myobroma* of the subgenus *Caprinus* and is related to *A. volkii* Rech. f. from which it differs in the characters of leaf, keel, vexillum and ovary. The size of the leaves in *A. volkii* Rech. f. is up to 12 cm., whereas in *A. staintonianus* Ali the size is of the order of 2-4 cm. The vexillum in *A. volkii* Rech. f. is more than three times longer than broad, whereas in *A. staintonianus* Ali it is less than three times as long as broad (19-22 mm. long, c. 8-10 mm. broad). The ratio of the length of the claw and lamina of the keel is also different in the two cases (*A. volkii*: 17-18/15; *A. staintonianus*: 11/6). Further, the ovary is only sparsely hairy in *A. volkii* Rech. f. but very densely hairy in *A. staintonianus* Ali (Fig. 16).

Astragalus imitensis Ali, spec. nov.

Herba perennis, c. 30-40 cm. alta. *Radix* lignosa. *Caulis* erectus pilosus. *Folia* c. 3-5,5 cm. longa, imparipinnata, stipulis lateralibus liberis, subulatis, c. 2 mm. longis. *Petiolum* c. 7-15 cm. longus. *Foliola* 9-11, petiolulo quam 1 mm. longiore, lamina c. 5-10 mm. longa, 2-4 mm. lata, obovata vel elliptica, margine integra, apice acuto, mucronato, utrinque aequaliter pilosa. *Inflorescentia* racemosa, c. 10-31 cm. (in fructu) longa, pedunculo c. 5-11 cm. longo. *Bracteae* c. 1-2 mm. longae. *Pedicelli* c. 1 mm. longi. *Calyx* aequaliter pilosus, parte connata c. 5-5,5 mm. longa, dentibus c. 1,5-2 mm. longis. *Corolla* (in sicco) flava, vexillo c. 12-13 mm. longo, c. 6 mm. lato, apice retuso; alis c. 11 mm. longis, c. 1,5-2 mm. latis, clave c. 5-6 mm. longa, auriculis c. 5 mm. longis. *Carina* c. 8 mm. longa, c. 2 mm. lata, clave c. 5 mm. longa. *Stamina* diadelpha, 9+1, filamentis c. 7-9 mm. longis. *Ovarium* c. 4 mm. longum, c. 1 mm. latum, aequaliter albo-pilosum, stylus glaber, c. 5 mm. longus; stigma capitatum. Legumen sessile 17 mm. longum, c. 2 mm. latum, pilis albis aequaliter pilosum, biloculare, seminibus numerosis (c. 8). Semina glabra, matura non visa.

HOLOTYPE: KASHMIR, Imit, 2900 m., 2-3 août 1954, *Schmid* 2072 (G) (Fig. 14).

This species belongs to the subgenus *Cercidothrix* and comes nearest to *A. subuliformis* DC. and *A. suffulcatus* DC. but differs from these in having a calyx smaller than 1 cm. (c. 7 mm.) (Fig. 17).

***Astragalus khalifatensis* Ali, spec. nov.**

Herba perennis, radice elongata. Caulis super terram reductus, internodiis stipulis persistentibus. *Folia* imparipinnata, stipulis lateralibus, c. 1-2 mm. longis, deltoidis, supra pilosis, infra glabris, rhachide 6-13 mm. longa, petiolo c. 3-8 mm. longo. Foliola 9, petiolulo quam 1 mm. longiore, lamina c. 2,5-5 mm. longa, c. 1-1,5 mm. lata, margine integra, apice obtuso. *Racemi* floriferi capitati ± densiusculi, pedunculo c. 3-4,5 cm. longo, albo-nigro piloso. Flores pedicellis c. 1,5 mm. longis. *Calyx* 17-18 mm. longus, tubulosus, tubo appresse, densiuscule albo-nigro-piloso, dentibus c. 3-3,5 mm. longis. *Corolla* (in sicco) lutea; vexillo c. 20 mm. longo, c. 6 mm. lato; alis c. 19 mm. longis, c. 1,5 mm.



FIG. 15. — *Astragalus khalifatensis* Ali, holotype (Schmid 152, G).




FIG. 16. — *Astragalus staintonianus* Ali. A: calyx and ovary. B: vexillum. C: wing. D: keel (*Stainton* 2874, RAW).




FIG. 17. — *Astragalus imitensis* Ali. A: calyx and ovary. B, C: vexillum. D: wing. E: keel. F: stamens (Schmid 2072, G).




FIG. 18. — *Astragalus khalifensis* Ali. A: calyx and ovary. B: vexillum. C: wing.
D: keel (Schmid 152, G).

latis, clave c. 12 mm. longa, auriculis c. 5 mm. longis; carina 16-17 mm. longa, c. 2,5 mm. lata, clave c. 11-12 mm. longa. *Stamina* diadelpha, 9+1, filamentis c. 17-18 mm. longis. *Ovarium* stipite c. 2 mm. longo sustentum c. 6 mm. longum, c. 1 mm. latum, glabrum, stylus c. 9-10 mm. longus; stigma capitatum. *Fructus* non vidi.

HOLOTYPE: BELOUTCHISTAN, sommet du Khalifat près de Ziarat, 3300 m. 7 mai 1953, Schmid 152 (G) (Fig. 15).

This species belongs to the subgenus *Cercidothrix* and resembles *A. patulipilosus* Sirj. & Rech. f. but differs in the glabrous nature of the ovary and in the panduriform vexillum (Fig. 18). In *A. patulipilosus* the ovary is pilose and the vexillum is elliptical. It resembles *A. virgaeformis* Sirj. & Rech. f. in having vertical rows of black hairs on the calyx, but its most marked difference lies in its having a compact inflorescence.

Acknowledgements:

My thanks are due to Dr. R. R. STEWART for allowing me the use of the herbarium and library of Gordon College, Rawalpindi and for sending me the specimens on loan. I also beg to express my indebtedness to the authorities of the Conservatoire et Jardin botaniques, Geneva.

