

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève
Band: 25 (1970)
Heft: 1

Artikel: New chromosome counts in Euphrasia
Autor: Yeo, P. F.
DOI: <https://doi.org/10.5169/seals-880317>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

New chromosome counts in *Euphrasia*

P. F. YEO

RÉSUMÉ

Le nombre chromosomique diploïde $n = 11$ est rapporté pour *Euphrasia alpina*, *E. picta* et *E. hirtella* ($n = \text{env. } 11$) et le nombre tétraploïde $n = 22$ pour *E. pectinata*. Un comptage précédent pour "*E. hirtella*" de Grande-Bretagne se réfère en réalité à *E. rostkoviana*. La discussion comporte des remarques sur la parenté d'autres espèces d'euphrasies grandiflores.

SUMMARY

The diploid chromosome number $n = 11$ is reported in *Euphrasia alpina*, *E. picta*, and *E. hirtella* ($n = \text{c. } 11$), and the tetraploid number $n = 22$ in *E. pectinata*. A previous count for "*E. hirtella*" from Britain is now referred to *E. rostkoviana*. The discussion includes remarks on the relationships of other large-flowered *Euphrasia* species.

ZUSAMMENFASSUNG

Die diploide Chromosomenzahl $n = 11$ wurde für *Euphrasia alpina*, *E. picta* und *E. hirtella* ($n = \text{ca. } 11$), und die tetraploide Zahl $n = 22$ für *E. pectinata* ermittelt. Eine frühere Zählung für "*E. hirtella*" der britischen Inseln wird jetzt auf *E. rostkoviana* bezogen. Die Diskussion bringt Bemerkungen über die Verwandtschaftsbeziehungen weiterer grossblütiger *Euphrasia*-Arten.

Results

In a previous paper (Yeo 1954) I have published chromosome counts for 15 British *Euphrasia* taxa. Using the method there described I have now obtained further counts at the stage of pollen mother cell meiosis from material originating in the Alps and cultivated in Cambridge in 1969. In the list that follows, numbers with the prefix E are the serial numbers of the samples; CGE indicates specimens preserved in the University Herbarium, Cambridge. The "fixed specimens" are those from which shoot apices were cut and fixed for cytological examination.

1. *E. alpina* Lam.

E 1072, Alpes-Maritimes, France; progeny of plants grown in 1968 from seed supplied by the Muséum national d'histoire naturelle, Paris (CGE 1968, 1969, including fixed specimens; herb. P. F. Yeo 1968, 1969): $n = 11$ (metaphase I, anaphase I, metaphase II).

E 1091, roadside above Brunnen, Simplon Pass, Valais, Switzerland, 1380 m alt., 5.8.1968 (CGE; herb. P. F. Yeo), cult. 1969 (CGE; herb. P. F. Yeo, including fixed specimens): $n = 11$ (metaphase I).

2. *E. picta* Wimmer

E 1119A, Enzinger Boden, Stubbachtal, Land Salzburg, Austria, 1480 m alt., 15.8.1968 (CGE; herb. P. F. Yeo), cult. 1969 (herb. P. F. Yeo): $n = 11$ (metaphase I, anaphase, I).

3. *E. pectinata* Ten.

E 1084, near Epinel, val de Cogne, Aosta, Italy, 1400 m alt., 3.8.1968 (CGE; herb. P. F. Yeo), cult. 1969 (CGE, including fixed specimens; herb. P. F. Yeo): $n = 22$ (metaphase I, anaphase I). In some cells at metaphase I in which a clear count was not possible one or two univalents were seen; in another such cell there were apparently four univalents.

4. *E. hirtella* Jordan ex Reuter

E 1101, Rothwald, Simplon Pass, Valais, Switzerland, 1745 m alt., 6.8.1968 (CGE, herb. P. F. Yeo), cult. 1969 (CGE): $n = c. 11$ (metaphase I, only 3 cells, none of which gave a perfectly clear count).

The general appearance of the chromosome complements in these samples was the same as in those previously illustrated (Yeo 1954, 1956).

Discussion

In my previous paper on the chromosomes of *Euphrasia* (Yeo 1954), I reported a count of $n = 11$ for *E. hirtella* var. *polyadena* (Gren. & Roux) Pugsley, from Lawers, Mid Perth, Scotland. I have since come to the conclusion that *E. hirtella* does not occur in the British Isles, and that the Lawers material is best referred to *E. rostkoviana* Hayne. The tentative count given here for *E. hirtella* is therefore the

first that is truly applicable to that species. Another taxonomic decision of mine, not yet published, is that *E. tatarica* Fischer ex Sprengel is not separable from *E. pectinata* Ten.; therefore, although the foliage of E 1084 is strongly hairy, so that these plants would in the past have been referred to *E. tatarica*, they are here indicated as *E. pectinata*.

As previously pointed out (Yeo 1954), whereas Wettstein divided the species with long glandular hairs between series *Grandiflorae* and series *Parviflorae*, Pugsley placed them all together in series *Hirtellae*, a decision supported by the apparent restriction of the diploid chromosome number ($2n = 22$) to this group. However, this treatment involved disregarding the marked similarity between the large-flowered *E. rostkoviana* Hayne (the commonest species of series *Hirtellae*) and the eglandular *E. kernerii* Wettst., a similarity strongly emphasised by Wettstein (1896). Pugsley (1936), following Rothmaler (1935), placed *E. kernerii*, and the earlier-flowering species *E. picta* Wimmer and *E. versicolor* Kerner, in series *Alpinae*. Recently Schaeftlein (1967) has thoroughly studied the occurrence of the large-flowered glandular species *E. rostkoviana* and *E. montana* Jord. and the large-flowered eglandular species *E. kernerii*, *E. versicolor* and *E. picta*, and has found that over a large part of the eastern Alps, Bavaria, Moravia and neighbouring areas, glandular and eglandular forms, otherwise indistinguishable, grow together. Furthermore, he found that from such populations the progeny of glandular plants included some eglandular plants, and vice versa. He concluded that such mixed populations constitute single breeding groups ("gamodemes") with gene-exchange between the two forms, and that the glandless taxa should provisionally be incorporated in an aggregate species under the name *E. rostkoviana*. In view of this one would expect the three eglandular taxa of this group to be diploid, and this is partly confirmed by the count here reported from Enzinger Boden. These plants came from a mixed population of glandular and eglandular plants; they are rather more branched and begin to flower at a slightly higher node than is the case with the most typical forms of *E. picta*, but agree with a description of this species which I drew up for "Flora Europaea" before I was aware of the intergradation with *E. rostkoviana*. The chromosomes were counted separately in two individual plants.

This group of plants evidently presents an awkward taxonomic problem, and I am not attempting to provide a solution here. However, the facts indicate that at least some of the species of series *Alpinae* ought to be placed together with some or all of those in series *Hirtellae*.

The relationship of *E. alpina* (the type species of series *Alpinae*) to the *E. kernerii* group is not yet clear. The chromosome number is the same, but the morphology is so different that *E. alpina* may deserve separate classification (together with its ally, *E. cisalpina* Pugsley). Indeed, it is possible that *E. alpina* and a member of the *E. rostkoviana* aggregate are joint parents of some or all of the European tetraploid groups other than subsection *Angustifoliae*, the *E. salisburgensis* group (see Yeo 1954: 106-107 and Yeo 1968: 736-737). The discovery of the diploid number in *E. alpina* is relevant to the origin of *E. christii* Favrat. This species was believed by Wettstein (1896) and Pugsley (1930) to be a hybrid between *E. alpina* and *E. minima* Jacq. ex Lam. & DC., which is a tetraploid species (Witsch 1932). *E. christii* may, therefore, be analogous to *E. vigursii* Davey, a diploid species believed to have originated by the crossing of another diploid species with a tetraploid (Yeo 1956).

Nomenclatural Note.

For convenience, the nomenclature used here for the subdivisions of the genus *Euphrasia* is that of Pugsley (1936), although this is not in accord with the International Code of Botanical Nomenclature (ed. 1966); it is hoped that a publication containing correct infrageneric nomenclature will appear soon (Sell and Yeo ined.).

BIBLIOGRAPHY

- Pugsley, H. W. (1930) A revision of the British Euphrasiae. *J. Linn. Soc. Bot.* 48: 467-542.
- (1936) Enumeration of the species of *Euphrasia* L., sect. *Semicalcaratae* Benth. *J. Bot.* 74: 273-288.
- Rothmaler, W. (1935) *Generum plantarum ibericum revisio critica*, III. *Euphrasia* L. *Cavanillesia* 7: 5-28.
- Schaeftlein, H. (1967) Beiträge zur Kenntnis einiger mitteleuropäischer Euphrasien, I. Der taxonomische Wert der drüsigen Behaarung. *Phyton (Horn)* 12: 48-90.
- Sell, P. D. & P. F. Yeo (ined.) *A revision of the North American species of Euphrasia* L.
- Wettstein, R. von (1896) *Monographie der Gattung Euphrasia*. Engelmann, Leipzig.
- Witsch, H. von (1932) Chromosomenstudien an mitteleuropäischen Rhinantheen. *Österr. Bot. Z.* 81: 108-141.
- Yeo, P. F. (1954) The cytology of the British species of *Euphrasia*. *Watsonia* 3: 101-108.
- (1956) Hybridisation between diploid and tetraploid species of *Euphrasia*. *Watsonia* 3: 253-269.
- (1968) The evolutionary significance of the speciation of *Euphrasia* in Europe. *Evolution* 22: 736-747.