

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	32 (1977)
<b>Heft:</b>	1
<b>Artikel:</b>	Studies on the flora of Jordan : 4. On the desert flora north-east of Aqaba
<b>Autor:</b>	Boulos, Loutfy / Lahham, Jamil
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-880220">https://doi.org/10.5169/seals-880220</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Studies on the flora of Jordan

### 4. On the desert flora north-east of Aqaba

LOUTFY BOULOS & JAMIL LAHHAM

#### Résumé

Boulos, L. & J. Lahham (1977). Etude de la flore de Jordanie 4. Flore du désert au nord-est d'Aqaba. *Candollea* 32: 81-98. En anglais.

Les auteurs énumèrent 250 espèces de plantes vasculaires récoltées en 1974 et en 1975 entre Wadi Yutum et Wadi Rum, en y joignant à l'occasion des remarques d'écologie, de phénologie et de répartition.

#### Abstract

Boulos, L. & J. Lahham (1977). Studies on the flora of Jordan 4. On the desert flora north-east of Aqaba. *Candollea* 32: 81-98. French abstract.

The authors enumerate 250 species of vascular plants collected, mainly in 1974 and 1975, in the area between Wadi Yutum and Wadi Rum. Occasional notes on ecology, phenology and distribution are added.

The area covered in the present study is defined from the north by the paved junction to Wadi Rum Rest House from the Amman—Aqaba highway at about 50 km N.E. of Aqaba, and from the east by the track to Wadi Rum Police Station, which joins the highway at Khirbet Kithara, some 15 km N.E.E. of Aqaba. At Khirbet Kithara, Wadi Yutum downstream extends to the west for about 6 km, while upstream ends at about 15 km N.E. of Khirbet Kithara.

Our area is located within the basement complex that is a part of the Arabian Shield. The rocks are mainly granodioritic and granitic, with numerous dikes of diabase and aplite granite. In the northern part, Cambrian sandstone overlies the Precambrian basement complex. The soil varies from sandy to sandy-clayey, with gravels in some areas.

The mean annual rainfall at Wadi Rum Police Station is 75.2 mm. The highest observed rainfall in 24 hours is 34.5 mm (March 1972). The mean maximum air temperature is 34.6°C in July. The mean minimum air temperature is 4.6°C in January. The altitude is 952 m.

The following is a list of the localities from which collections were made (see map, fig. 1).




Fig. 1. – Map of the area N.E. of Aqaba, Jordan.

- 6085-6117: 6 km N. of Aqaba, 13.3.1974, *B., A-E. & J.*<sup>1</sup>
- 163-166: idem, *J.*
- 6118-6202: 10-12 km N. of Aqaba, 13.3.1974, *B., A-E. & J.*
- 152-162: idem, *J.*
- 6203-6217: Wadi Rum, near the Rest House, 14.3.1974, *B., A-E. & J.*
- 178-181: idem, *J.*
- 6218: 5 km W. of Wadi Rum Rest House, 14.3.1974, *B., A-E. & J.*
- 6650-6695: 9 km N. of Aqaba, 5.4.1974, *B., J. & L.*
- 316-325: idem, *J.*
- 167-177: 9 km N. of Aqaba, 13.3.1974, *J.*
- 7445-7505: Wadi Yutum upstream, 15 km N. of Aqaba, 21.3.1975, *B., J., L. & A-H.*
- 251-252: idem, *L.*
- 50-55: idem, *T.*

<sup>1</sup>Collectors' names are abbreviated as follows: *A.* = D. Addis; *A-E.* = D. Al-Eisawi; *A-H.* = M. A. Abu-Hmaidan; *B.* = L. Boulos; *J.* = W. Jallad; *L.* = J. Lahham; *Q.* = M. Qumsiyeh; *T.* = M. Z. Tarabeih.

- 7506-7534: Wadi Yutum, on the slopes along the road, 9 km N. of Aqaba, 21.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 253: idem, *L.*
- 56-57: idem, *T.*
- 7535-7561: Wadi Yutum, on the roadside, 12 km N. of Aqaba, 21.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 7562-7576: 30 km S.W. of Rum Police Station, along the track to Aqaba, 23.3.-1975, *B.*, *J.*, *L.* & *A-H.*
- 7577-7596: 15-16 km S.E. of Rum Police Station, along the track to Aqaba, 23.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 7597-7669: 6-7 km S. of Wadi Rum Rest House, 23.3.1975, *B.*, *J.*, *L.* & *A-H.*

Some of the species recorded from this locality are halo- and hydrophytes, e.g.: *Mentha longifolia* subsp. *typhoides*, *Zannichellia palustris*, *Juncus bufonius* and *Samolus valerandi*. These plants grow in and around a small water pool. The site could easily be seen from a distance, due to the presence of a large *Acacia gerrardii* var. *negevensis* tree on the western side of the wadi at a few hundred metres up the hilly slopes.

Another microclimate is created due to the presence of a protected area within eroded rocks near the pool, which forms an open cave with a large shelf as a shelter. This supports the growth of some shade-loving plants, e.g.: *Adiantum capillus-veneris*, some mosses, and shade forms of several desert species. The drooping branches of *Capparis spinosa* are also typical of such habitats; the latter species forms a conspicuous part of the shrub vegetation.

- 7670-7709: 4 km S. of Wadi Rum Rest House, along the road to Rum Police Station, 23.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 7710-7750: 3-4 km W. of Wadi Rum Rest House (c. 18 km from the main road to Aqaba), 23.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 7751-7791: vicinity of Wadi Rum Rest House, 23.3.1975, *B.*, *J.*, *L.* & *A-H.*
- 7812-7821: 7 km S. of Wadi Rum junction, 4.4.1975, *B.* & *J.*
- 7822-7826: 2 km S. of Wadi Rum Rest House, 4.4.1975, *B.* & *J.*
- 8545: 10 km N. of Aqaba, 1.5.1976, *B.*, *A.* & *Q.*

The number of taxa recorded from the area is 256, which includes 250 species and 6 varieties belonging to 45 families. The largest families are: *Compositae*: 41 species; *Cruciferae*: 24 species and 1 variety; *Gramineae*: 23 species, and *Leguminosae*: 20 species and 2 varieties. It is remarkable that out of about 80 species of *Umbelliferae* known from Jordan, only 1 species is recorded from our area. The main centre of distribution of this family in Jordan seems to be in the N.W.

Specimens of the above-mentioned collections were deposited in the Herbarium, Faculty of Science, University of Jordan, Amman, Jordan. Duplicates were distributed to the following herbaria: AAU, ATH, ATHU, BM, C, CAI, E, G, H, K, LD, RNG, S and ZT.

The following is an alphabetically arranged list of the families, with the genera and species related to them. The numbers refer to the localities from which the specimens were collected. The abbreviations used in the text are:

- fl. = flowering specimen
- fr. = fruiting specimen
- ster. = sterile specimen, without flowers or fruits.

*Acanthaceae*

**Blepharis ciliaris** (L.) B. L. Burtt  
7532 (seedlings, ster.).

*Aizoaceae*

**Aizoon canariense** L.  
6106 (fl., fr.) – 6681 (fl., fr.).

*Amaranthaceae*

**Aerva persica** (Burm. fil.) Merr.  
6141 (fl.) – 6664 (fl.) – 7557 (fl.).  
Fairly common shrublet, often with thicker growth in protected parts.

*Asclepiadaceae*

**Caralluma** cf. *aaronis* (Hart) N. E. Br.  
7750 (young fr.). Locally fairly common in sheltered fissures facing southeast.

**Pergularia tomentosa** L.  
6183 (fl.) – 6655 (fl., fr.) – 7556 (fl., fr.).

**Periploca angustifolia** Labill.  
7653 (fl., fr.).

*Boraginaceae*

**Anchusa aegyptiaca** (L.) DC.  
7501 (fl., fr.) – 7661 (fl., fr.).

*Asperugo procumbens* L.

7598 (fl., fr.).

*Echium longifolium* Delile  
6095 (fl.).

**Gastroctyle hispida** (Forsskål) Bunge  
6158 (fl.) – 6211 (fl.) – 7657 (fl., fr.) – 7676 (fl., fr.) – 7774 (fl., fr.).  
Common annual in sandy and alluvial soils.

**Heliotropium arbainense** Fresen.  
6654 (fl.) – 7503 (fl.).

**Heliotropium ramosissimum** (Lehm.) DC.  
6123 (fl.).

**Lappula spinocarpos** (Forsskål) Ascher-  
son  
6121 (fl.) – 6207 (seedlings, ster.) –  
7473 (fl., fr.) – 7618 (fl., fr.) –  
7693 (ster.) – 7769 (fl., fr.).

**Onosma orientalis** L.  
7653 (fl.).

**Paracaryum intermedium** (Fresen.) Lipsky  
6126 (fl., fr.) – 6670 (fl., fr.) –  
7481 (fl.) – 7542 (fl., fr.) – 7605 (fl., fr.) –  
7691 (fl., fr.) – 7726 (fl., fr.) –  
7785 (fl., fr.) – 7816 (fl.). Delicate small annual, common.

**Paracaryum rugulosum** (DC.) Boiss.

7681 (fl., fr.). Perennial, rare.

**Trichodesma africanum** (L.) R. Br. var.  
*heterotrichum* Bornm. & Kn.

6162 (fl.) – 7553 (fl.) – 7680 (fl.).

*Capparaceae***Capparis spinosa** L.

7636 (ster.). The species concept and nomenclature of this and other related *Capparis* species are discussed by several authors, e.g.: Zohary (1960, 1966), Bočancev (1964b), St John (1965), Täckholm & Boulos (1974).

*Caryophyllaceae***Gymnocarpos decandrum** Forsskål52 (fl.) – 55 (fl.) – 7704 (fl.) –  
7747 (fl.). Common shrub.**Gypsophila arabica** Barkoudah

56 (ster.).

**Gypsophila viscosa** Murr.

7606 (fl.).

**Herniaria hirsuta** L.

6153 (fl.) – 6673 (fl., fr.).

**Minuartia picta** (Sibth. & Sm.) Bornm.6203 (fl.) – 7671 (fl., fr.) – 7771  
(fl., fr.).**Paronychia arabica** (L.) DC.

7609 (fl.).

**Paronychia sinaica** Fresen.

6193 (fl., fr.) – 7712 (fl., fr.). The above two species of *Paronychia* are generally restricted to the southern

desert in Jordan. They are replaced by the common *Paronychia argentea* Lam. in the north.

**Polycarpaea repens** (Forsskål) Ascher-  
son & Schweinf.

6187 (fl.).

**Robbairea delileana** Milne-Redhead

50 (fl.) – 54 (fl.) – 57 (fl.) – 6085  
(fl., fr.) – 6173 (fl., fr.) – 6678 (fl.) –  
7570 (fl., fr.) – 7587 (fl., fr.) –  
7689 (fl.) – 7710 (fl.) – 7753 (fl., fr.).  
One of the commonest desert species  
in southern Jordan.

**Silene colorata** Poiret subsp. *oliveriana*  
(Otth) Rohrb.6148 (fl.) – 7579 (fl.) – 7623 (fl.) –  
7789 (fl.).**Silene conoidea** L.

7619 (fl.).

**Silene villosa** Forsskål

7581 (fl.).

In Jordan *Silene* L. is represented by 25 species, mostly distributed in the temperate, rather than the warmer parts of the country. Zohary (1973), deals with *Silene* species of Palestine and the Middle East, with special emphasis on their geographical distribution, endemism, ecology, etc.

**Spergularia diandra** (Guss.) Heldr. &  
Sart.

7603 (fl., fr.).

**Telephium sphaerospermum** Boiss.51 (fl., fr.) – 7686 (fl.) – 7714  
(fl., fr.).

*Chenopodiaceae****Atriplex dimorphostegia* Kar. & Kir.**

6210 (ster.) – 7649 (ster.) – 7755  
(ster.) – 7808 (fl., fr.).

***Bassia latifolia* (Fresen.) Ascherson &  
Schweinf.**

7751 (fl.) – 7810 (fl.).

***Bassia muricata* (L.) Murr.**

6091 (fl.) – 6152 (fl., fr.) – 6206  
(seedlings) – 7644 (fl.) – 7761 (fl.) –  
7811 (fl., fr.).

***Chenopodium murale* L.**

7475 (fl.).

***Halogeton alopecuroides* (Delile) Moq.**

7737 (ster.).

***Haloxylon persicum* Bunge**

6218 (fl.) – 7791 (fl.). Shrub or  
small tree up to 4 m high. Unfortunately the larger specimens are becoming rare as the Bedouins cut them down for fuel.

*Cistaceae****Helianthemum kahiricum* Delile**

7510 (fl.) – 7734 (fl.).

***Helianthemum ledifolium* (L.) Miller**

7600 (fl., fr.) – 7759 (fl., fr.).

***Helianthemum lippii* (L.) Dum.-Courset**

6200 (fl.) – 7703 (fl.) – 7736 (fl.,  
fr.).

***Helianthemum ventosum* Boiss.**

7496 (fl., fr.).

*Cleomaceae****Cleome amblyocarpa* Barratte & Murb.**

6159 (fl., fr.) – 6665 (fl., fr.) –  
7486 (fl., fr.) – 7522 (fl., fr.) – 7573  
(fl.).

***Cleome arabica* L.**

6137 (fl., fr.) – 7484 (fl., fr.) –  
7517 (fl., fr.) – 7571 (fl., fr.). The  
nomenclature of the above two species  
is discussed by Bočancev (1964a,  
1968) and Kers (1966).

***Cleome droserifolia* (Forsskål) Delile**

6197 (fr.) – 7519 (fr.).

*Compositae*<sup>1</sup>***Aaronsohnia faktorovskyi* Warburg &  
Eig**

6119 (fl., fr.) – 7728 (fl., fr.).

***Anthemis deserti* Boiss.**

6695 (fl.)

***Anthemis melampodina* Delile**

6154 (fl.).

***Anthemis* sp.**

7778 (fl.).

***Artemisia inculta* Delile**

= *A. herba-alba* Asso

6174 (fl.) – 6677 (ster.).

***Artemisia judaica* L.**

7593 (fl., fr.) – 7821 (fl., fr.). Three  
species are known from Jordan: *Arte-  
misia judaica* is restricted to the south-  
western desert; *A. inculta*, a very com-

---

<sup>1</sup> See Täckholm (1974), and Davis (1975).

mon desert shrublet, is dominant in many regions forming the *Artemisia-tum herbae-albae* association (Zohary 1962, 1973); *A. monosperma* Delile seems to be rare, and was only collected once by Boulos, Al-Eisawi & Jallad from Ras en Naqb area in 1974.

**Asteriscus graveolens** Less.

7475 (fl.) – 7527 (fl.).

**Asteriscus pygmaeus** (DC.) Cosson & Durieu

6125 (fl.) – 6684 (fl.).

**Astropterus leyseroides** (Desf.) Rothm.

6149 (fl., fr.) – 7544 (fl., fr.) – 7630 (fl., fr.) – 7732 (fl., fr.). A rather common annual in shallow soils. For synonyms, see Täckholm (1974).

**Atractylis carduus** (Forsskål) C. Chr.

7575 (fl., fr.) – 7781 (fl.).

**Calendula arvensis** L.

7620 (fl., fr.) – 7767 (fl., fr.). According to Heyn & al. (1974), our specimens may be referred to the *Calendula arvensis* complex. For synonyms, chromosome records, and relationships see Heyn & al., l.c.

**Carthamus cf. glaucus** M.B.

7707 (ster.).

**Centaurea cf. araneosa** Boiss.

7787 (fl.).

**Centaurea calcitrapella** Bornm. & Dinsm.

7633 (fl., fr.) – 7813 (fl., fr.).

**Centaurea eryngioides** Lam.

7740 (fl.).

**Centaurea laxa** Boiss. & Hausskn.

7678 (fl.).

**Centaurea pallescens** Delile

6104 (fl., fr.) – 6133 (fl.) – 6688 (fl., fr.) – 7543 (fl.) – 7631 (fl.).

**Filago desertorum** Pomel

6176 (fl.) – 6675 (fl.) – 6694 (fl., fr.) – 7447 (fl., fr.).

**Filago** sp.

7537 (fl., fr.). A tiny ephemeral, 1-5 cm. Specimens collected c. 5 km S.E. of Aqaba, *Boulos 5396*, may refer to this species; see Boulos & Lahham (1977).

**Ifloga spicata** (Forsskål) Schultz Bip.

6092 (fl., fr.) – 6157 (fl., fr.) – 6663 (fl., fr.) – 7452 (fl., fr.) – 7604 (fl., fr.) – 7768 (fl., fr.). Common small annual, in shallow sandy soils.

**Iphiona mucronata** (Forsskål) Ascher-  
son & Schweinf.

7746 (ster.). Another species: *Iphiona scabra* DC. is recorded from a desert wadi 8 km S. of Aqaba (Boulos & Lahham 1977). Both species are restricted to the S.W. corner of Jordan, which is the hottest desert in the country. Zohary (1962), cites *Iphiona* Cass. among other genera such as *Cleome* L., *Caralluma* R. Br., *Trichodesma* R. Br., etc., in the "Flora palaestina" as being of Sudanian or tropical origin.

**Jasonia montana** (Vahl) Botsch.

7555 (ster.).

**Koelpinia linearis** Pallas

7562 (fl., fr.) – 7607 (fl., fr.) – 7776 (fl., fr.).

- Lasiopogon muscoides** (Desf.) DC.  
*6146* (fl., fr.) – *6204* (fl.) – *7679* (fl., fr.).
- Lasiospermum brachyglossum** DC. (det. C. Jeffrey)  
*7455B* (fl., fr.). Native of S. Africa, new to Jordan.
- Launaea capitata** (Sprengel) Dandy  
*6672* (fl., fr.).
- Launaea mucronata** (Forsskål) Muschler  
*6094* (fl.) – *6175* (ster.) – *7588* (fl.).
- Launaea nudicaulis** (L.) Hooker fil.  
*6163* (fl., fr.) – *7520* (fl., fr.) – *7551* (fl., fr.) – *7730* (fl., fr.).
- Launaea spinosa** (Forsskål) Schultz Bip.  
*6171* (ster.) – *7498* (fr.). The genus *Launaea* Cass. is represented in Jordan by the above four species.
- Phagnalon rupestre** (L.) DC.  
*7650* (fl., fr.) – *7720* (fl.).
- Picris asplenoides** L.  
= *P. radicata* Forsskål; *P. lyrata* Delile; *P. pilosa* Delile  
*6098* (fl., fr.) – *6135* (fl., fr.) – *6682* (fl., fr.).
- Picris babylonica** Hand.-Mazz.  
*7779* (fl., fr.) – *7817* (fl., fr.).
- Picris longirostris** Schultz Bip.  
= *P. damascena* Boiss. & Gaill.; *P. blancheana* Boiss.; *P. desertorum* Nábelék  
*7464* (fl., fr.) – *7694* (fl., fr.).
- Pulicaria undulata** (L.) Kostel.  
*6101* (fl., fr.) – *6132* (fl., fr.) – *6674* (fl., fr.) – *7497* (fl., fr.) – *7508* (fl., fr.).
- Reichardia tingitana** (L.) Roth  
*6120* (fl., fr.) – *6686* (fl., fr.) – *7462* (fl., fr.) – *7744* (fl., fr.) – *7782* (fl., fr.).
- Scorzonera mollis** M.B.  
*7683* (ster.).
- Scorzonera pseudolanata** Grossh.  
*7698* (fl., fr.).
- Senecio coronopifolius** Desf.  
*6156* (fl., fr.) – *7450* (fl., fr.) – *7576A* (fl., fr.) – *7675* (fl., fr.).
- Senecio flavus** (Decne) Schultz Bip.  
*6090* (fl., fr.) – *6151* (fl., fr.) – *6683* (fl., fr.) – *7460* (fl., fr.) – *7541* (fl., fr.) – *7645* (fl., fr.).
- Tripleurospermum auriculatum** (Boiss.) Rech. fil.  
= *Chamaemelum auriculatum* Boiss.  
*7455A* (fl., fr.).
- Urospermum picroides** (L.) F. W. Schmidt  
*7647* (fl., fr.).
- Cruciferae*<sup>1</sup>
- Biscutella didyma** L.  
*7601* (fl., fr.) – *7690* (fl., fr.).

---

<sup>1</sup>The following specimens belonging to this family were determined by W. Jallad, and most of the collections were made by him.

**Brassica tournefortii** Gouan

175A (fl., fr.).

**Clypeola jonthlaspi** L.

7696 (fl., fr.).

**Diplotaxis acris** (Forsskål) Boiss.

156B (fl., fr.).

**Diplotaxis harra** (Forsskål) Boiss.

156A (fl., fr.) – 161 (fl., fr.) – 175B (fl.) – 325 (fl., fr.) – 7489 (fl., fr.) – 7506 (fl., fr.). Specimens of this species kept together with *Cleome amblyocarpa* Barratte & Murb. in the same polyethylene bag during collection changed colour from yellow to purple; the new colour is still retained by the herbarium specimens. Boulos & Jallad (1975) report five species of *Diplotaxis* DC. in Jordan.

**Eremobium aegyptiacum** (Sprengel)  
Ascherson & Schweinf. ex Boiss.  
var. *lineare* (Delile) Zoh.

7566 (fl., fr.) – 7584 (fl., fr.).

**Eremobium diffusum** (Decne) Botsch.

179B (fl.) – 320 (fl., fr.).

**Erucaria boveana** Cosson

169 (fl., fr.) – 322 (fl., fr.).

**Farsetia aegyptia** Turra

152 (fl., fr.) – 7554 (fl., fr.) – 7663 (fl.).

**Isatis lusitanica** L.

170 (fl.) – 319 (fl., fr.).

**Leptaleum filifolium** (Willd.) DC.

7818 (fl., fr.).

**Malcolmia africana** (L.) R. Br.

181 (fl.).

**Maresia pygmaea** (Delile) O. E. Schulz

180 (fl., fr.).

**Matthiola longipetala** (Vent.) DC. var.  
*bicornis* (Sibth. & Sm.) Zoh.

160 (fl., fr.) – 165 (fl., fr.).

**Matthiola longipetala** (Vent.) DC. var.  
*longipetala*

172 (fl., fr.) – 323 (fl., fr.).

**Morettia canescens** Boiss.

154 (fl., fr.) – 166 (fl., fr.) – 167 (fl., fr.) – 171 (fl., fr.) – 317 (fl., fr.) – 7493 (fl., fr.) – 7733 (fl.).

**Morettia parviflora** Boiss.

7495 (fl., fr.) – 7513 (fl., fr.) – 7552 (fl., fr.) – 7569 (fl.).

**Notoceras bicone** (Sol.) Caruel

158 (fl., fr.) – 173 (fl., fr.) – 178 (fl.) – 318 (fl., fr.) – 7814 (fl.).

**Schimpera arabica** Hochst. & Steudel  
ex Boiss.

7580 (fl.) – 7639 (fl., fr.).

**Sinapis arvensis** L. var. *arvensis*

467A (fr.).

**Sisymbrium bilobum** (C. Koch) Grossh.

7455 (fl., fr.).

**Sisymbrium erysimoides** Desf.

155 (fl., fr.) – 159 (fl., fr.) – 163 (fl., fr.) – 174 (fl., fr.) – 321 (fl., fr.).

**Sisymbrium irio** L.

7499 (fl., fr.) – 7610 (fl., fr.).

**Stigmatella longistyla** Eig

179A (fl.) – 7625 (fl., fr.).

**Zilla spinosa** (Turra) Prantl

153 (ster.) – 168 (fl., fr.) – 316  
(fl., fr.) – 7467 (fl., fr.).

*Cucurbitaceae***Citrullus colocynthis** (L.) Schrader

6169 (ster.).

**Cucumis prophetarum** L.

6109 (fl.) – 6165 (fl., fr.) – 6679  
(fl., fr.) – 7525 (fl., fr.).

*Cyperaceae***Scirpus holoschoenus** L.

7662 (fl.). In the vicinity of a water  
pool.

*Euphorbiaceae***Andrachne aspera** Sprengel

251 (fl., fr.) – 6190 (fl., fr.).

**Andrachne telephioides** L.

6195 (fl., fr.).

**Chrozophora oblongifolia** (Delile) A.  
Juss. ex Sprengel

253 (fl., fr.) – 6199 (fl., fr.).

**Euphorbia chamaepeplus** Boiss. & Gaill.

252 (fl., fr.) – 6196 (fl., fr.) –  
6669 (fl., fr.) – 7715 (fl., fr.).

*Geraniaceae***Erodium deserti** (Eig) Eig

7448 (fl., fr.) – 7617 (fl., fr.).

**Erodium glaucophyllum** (L.) Aiton

6112 (fl., fr.).

**Erodium hirtum** Willd.

7483 (fl.) – 7702 (ster.).

**Erodium laciniatum** (Cav.) Willd. var.  
*laciniatum*

6130 (fl., fr.) – 6213 (fl., fr.) –  
7577 (fl., fr.).

**Erodium laciniatum** (Cav.) Willd. var.  
*pulverulentum* (Cav.) Boiss.

7461 (fl., fr.) – 7611 (fl., fr.) –  
7719 (fl., fr.) – 7757 (fl., fr.) – 7812  
(fl., fr.).

**Erodium oxyrrhynchum** M.B.

7692 (fl.).

**Monsonia nivea** (Decne) Decne

6103 (fl., fr.) – 7766 (fl., fr.).

*Gramineae***Aegilops caudata** L.

7825 (fl., fr.).

**Aegilops kotschy** Boiss.

7822 (fl.).

**Aristida adscensionis** L.

7480 (fl.).

**Bromus tectorum** L.

7624 (fl.) – 7758 (fl.).

**Bromus** sp.

7672 (fl.).

**Cenchrus ciliaris** L.

6198 (fl.) – 7454 (fl.).

**Cutandia dichotoma** (Forsskål) Trabut

7590 (fl.).

- Cutandia memphitica** (Sprengel)  
Bentham  
7613 (fl.) – 7786 (fl.) – 7826 (fl.).
- Cynodon dactylon** (L.) Pers.  
7638 (fl.).
- Enneapogon desvauxii** Beauv.  
6116 (fl.) – 7533 (fl.).
- Enneapogon** sp.  
7528 (fl.).
- Hordeum glaucum** Steudel  
7568 (fl.) – 7627 (fl.).
- Lophochloa pumila** (Desf.) Bor  
7446 (fl.).
- Poa bulbosa** L.  
7688 (fl.) – 7742 (fl.). A polymorphic species that needs revision in Jordan. Specimens from the above and other localities could be confused with *Poa sinaica* Steudel. Bor (1968) shows that in Iraq both species occur in almost separated geographical areas.
- Schismus barbatus** (L.) Thell.  
6086 (fl.) – 6662 (fl., fr.) – 7515 (fl.) – 7545 (fl.) – 7565 (fl.) – 7597 (fl.) – 7773 (fl.). Täckholm & al. (1941), Täckholm (1974), consider *Schismus arabicus* Nees as a synonym of *S. barbatus*. Other authors consider it as a subspecies: *S. barbatus* (L.) Thell. subsp. *arabicus* (Nees) Maire & Weiller. Haines (1962, cited by Bor 1968), Bor (1964, 1968) and others consider both taxa as two distinct species. However, Bor (1968: 377) writes: "It is proposed in this work to retain them as distinct, mainly on the authority of Haines (in Agnew, Bull. Coll. Sci. Univ. Baghdad 6, suppl.: 87.

1962) who has had very considerable experience of this genus in the field. It is extremely likely that hybridisation occurs, since both these species grow together, and this may be the real reason for much of the difficulty". It is obvious that it is extremely difficult for a non-specialist to separate both 'taxa', especially when one deals with intermediate forms. It is proposed here to list *Schismus arabicus* Nees in the synonymy under *S. barbatus* (L.) Thell. unless future studies reveal distinct differences between them.

**Sclerochloa woronowii** (Hack.) Tzvelev  
6209 (fl.).

**Stipa capensis** Thunb.  
6088 (fl.) – 7474 (fl.) – 7784 (fl.).

**Stipagrostis obtusa** (Delile) Nees  
6194 (fl.) – 7820 (fl.).

**Stipagrostis plumosa** (L.) Munro ex T. Anders.  
6110 (fl.) – 6177 (fl.) – 7591 (fl.).

**Stipagrostis** sp.  
7530 (fl.).

**Tetrapogon villosus** Desf.  
7526 (fl.) – 7705 (fl.) – 7713 (fl.).

**Trachynia distachya** (L.) Link  
6189 (fl.).

**Tricholaena teneriffae** (L. fil.) Link  
6185 (fl.) – 7548 (fl.).

#### *Hypocoaceae*

**Hypcum imberbe** Sibth & Sm.  
7780 (fl., fr.).

**Hypocoum pendulum L.**

6161 (fl., fr.) – 7809 (fr.).

*Iridaceae***Iris sibiricum L.**7594 (fl., fr.) – 7701 (fl.) – 7741  
(fl., fr.).*Juncaceae***Juncus bufonius L.**

7641 (fl.).

**Juncus rigidus Desf.**= *J. arabicus* (Ascherson & Buch)  
Adams7660 (fl., fr.). Both *Juncus* species  
were growing on the borders of a small  
water pool.*Labiatae***Ajuga chia (Poiret) Schreber**7608 (fl., fr.). Ball (in Tutin & al.  
1972), treats this perennial species as a  
subspecies of the annual *Ajuga cha-*  
*maepitys* (L.) Schreber.**Ballota undulata (Sieb. ex Fresen.)  
Bentham**

7735 (ster.).

**Lavandula stricta Delile**6139 (fl.) – 7500 (fl., fr.) – 7523  
(fl.) – 7738 (fl.).**Mentha longifolia (L.) L. subsp. ty-**  
**phoides (Briq.)**

7654 (fl.).

**Micromeria myrtifolia Boiss. & Hohen.**

7477 (fl.).

**Micromeria sp.**

7700 (ster.) – 7748 (ster.).

**Salvia aegyptiaca L.**

7456 (fl., fr.) – 7634 (fl., fr.).

**Salvia deserti Decne**

6201 (fl.) – 7470 (fl.) – 7511 (fl.).

**Salvia lanigera Poiret**

7458 (fl., fr.).

**Stachys aegyptiaca Pers.**

7664 (fl.).

**Teucrium polium L.**

6142 (ster.).

*Leguminosae*<sup>1</sup>**Acacia gerrardii** Bentham subsp. nege-  
vensis Zoh.

7667 (fr.).

**Argyrolobium crotalarioides** Jaub. &  
Spach

7559 (fl.).

**Astragalus acinaciferus** Boiss.

7592 (fl., fr.).

**Astragalus bombycinus** Boiss.7491 (fl., fr.) – 7512 (fl., fr.) –  
7682 (fl., fr.).**Astragalus eremophilus** Boiss.

6089 (fl., fr.).

---

<sup>1</sup> See Davis (1970), Zohary (1972),  
Townsend & Guest (1974) and Täckholm  
(1974).

- Astragalus sparsus** Delile ex Decne  
6155 (fl.) – 7550 (fl.). Calyx red-dish-brown, corolla greenish-yellow.
- Astragalus spinosus** (Forsskål) Muschler  
7482 (fl., fr.) – 7595 (fl.).
- Astragalus tribuloides** Delile var. elarishensis Eig  
6129 (ster.).
- Astragalus tribuloides** Delile var. minutus (Boiss.) Boiss.  
7725 (fl.).
- Astragalus tribuloides** Delile var. tribuloides  
6131 (fl., fr.) – 6685 (fl., fr.) – 7536 (fl., fr.) – 7564 (fl.) – 7589 (fl., fr.) – 7752 (fl., fr.).
- Cassia italica** (Miller) Lam. ex Steudel  
6652 (fl., fr.).
- Crotalaria aegyptiaca** Bentham  
6144 (fl., fr.) – 7560 (fl., fr.).
- Lotononis platycarpa** (Viv.) Pichi-Sermanni  
6099 (fl., fr.) – 7502 (fl., fr.) – 7529 (fl., fr.) – 7538 (fl., fr.) – 7558 (fl., fr.).
- Medicago laciniata** (L.) Miller var. laciniata  
6191 (fl.) – 7578 (fl., fr.) – 7615 (fl., fr.) – 7756 (fl., fr.).
- Ononis natrix** L.  
7616 (fl.) – 7788 (fl.).
- Ononis serrata** Forsskål  
7567 (fl.) – 7582 (fl., fr.) – 7724 (fl.) – 7754 (fl.).

- Ononis sicula** Guss.  
6687 (fl., fr.).
- Retama raetam** (Forsskål) Webb & Berth.  
7709 (fl.) – 7790 (fl.).
- Tephrosia apollinea** (Delile) Link  
6143 (fr.).
- Trigonella astroites** Fischer & Meyer  
7599 (fl., fr.) – 7824 (fl., fr.).
- Trigonella cf. moabitica** Zoh.  
6691 (fl., fr.).
- Trigonella stellata** Forsskål  
6102 (fl., fr.) – 6170 (fl., fr.) – 6215 (old fr.) – 7449 (fl., fr.) – 7546 (fl., fr.) – 7629 (fl., fr.) – 7718 (fl., fr.).
- Liliaceae*
- Asphodelus tenuifolius** Cav.  
6105 (fl., fr.) – 6659 (fl., fr.) – 7487 (fl.) – 7572 (fl., fr.) – 7727 (fl., fr.) – 7783 (fl., fr.).
- Bellevalia cf. zoharyi** Feinbrun  
7628 (fr.).
- Bellevalia** sp.  
7670 (fr.).
- Colchicum ritchii** R. Br.  
6216 (fl.) – 7656 (fr.) – 7685 (fr.) – 7739 (fr.).
- Gagea reticulata** (Pallas) A. & H. Schulthes  
6214 (fl.).

***Urginea maritima* (L.) Baker**

7504 (ster.).

*Loranthaceae****Loranthus acaciae* Zucc.**

7505 (ster.) – 8545 (fl.). Flowers bright crimson red; parasitic on *Acacia raddiana*, was not observed on *A. tortilis*.

*Malvaceae****Abutilon fruticosum* Guill. & Perr.**

6178 (ster.) – 7520 (fl.).

***Hibiscus micranthus* L. fil.**

6180 (ster.).

***Malva parviflora* L.**

7468 (fl., fr.) – 7622 (fl., fr.) –  
7765 (fl.).

***Malvella sherardiana* (L.) Jaub. & Spach**

6179 (ster.).

*Moraceae****Ficus pseudosycomorus* Decne**

6167 (ster.) – 7666 (fr.).

*Neuradaceae****Neurada procumbens* L.**

7763 (fl., fr.).

*Nyctaginaceae****Boerhavia diffusa* L.**

6653 (ster.).

***Boerhavia repens* L.**

6118 (fl.) – 7472 (ster.) – 7534 (fl.).

*Orobanchaceae****Cistanche tubulosa* (Schenk) Wight**

7596 (fl.).

*Papaveraceae****Papaver glaucum* Boiss. & Hausskn. ex Boiss.**

6671 (fr.) – 7476 (fr.) – 7602 (fl., fr.) – 7674 (fr.). For distribution in Jordan see Boulos & al., 1975.

***Roemeria hybrida* (L.) DC.**

7677 (fl., fr.) – 7716 (fl., fr.) – 7819 (fl., fr.).

*Plantaginaceae****Plantago afra* L.**

6184 (fl., fr.) – 7445 (fl., fr.) – 7642 (fl., fr.) – 7722 (fl., fr.).

***Plantago ciliata* Desf.**

6147 (fl.) – 6690 (fl., fr.).

***Plantago coronopus* L.**

7815 (fl., fr.).

***Plantago cylindrica* Forsskål**

6145 (fl.).

***Plantago ovata* Forsskål**

6093 (fl., fr.) – 6693 (fl., fr.) – 7463 (fl.) – 7673 (fl., fr.) – 7717 (fl., fr.) – 7764 (fl., fr.).

*Polygonaceae****Emex spinosa* (L.) Campd.**

6124 (fl.) – 6212 (ster.) – 7466 (fl., fr.).

**Rumex cyprius** Murb. emend. Sam.

6108 (fl.) – 6127 (fl.) – 6651 (fl., fr.) – 7695 (fl., fr.).

*Primulaceae***Samolus valerandi** L.

7643 (ster.).

*Ranunculaceae***Ranunculus asiaticus** L.

7687 (fl.). Flowers scarlet red; in Jordan this seems to be the only colour for this species. The present locality is apparently the most southern for the species in Jordan. This may be attributed to the high altitude, c. 950 m. The species is mainly located in the northwest; but it is also found as far S. as the Amman area.

*Resedaceae***Caylusea hexagyna** (Forsskål) Green

6122 (fl., fr.) – 6676 (fl.) – 7488 (fl., fr.).

**Ochradenus baccatus** Delile

6689 (fl.) – 7531 (fl.) – 7632 (fl., fr.) – 7749 (fl.).

**Oligomeris linifolia** (Vahl) Macbride

5506 (fl., fr.).

**Reseda arabica** Boiss.

6097 (fl.).

**Reseda decursiva** Forsskål

6164 (fl.).

**Reseda muricata** C. Presl

6166 (fl.) – 6658 (fl.) – 7485 (fl.) – 7524 (fl.) – 7547 (fl.) – 7658 (fl.).

*Rubiaceae***Callipeltis cucullaris** (L.) Rothm. var. *aptera* (Boiss. & Buhse) Rech. fil. & Ehrend.

6128 (fl.) – 7721 (fl., fr.).

**Crucianella herbacea** Forsskål

6115 (ster.) – 6138 (ster.) – 7612 (fl.) – 7775 (ster.).

**Galium canum** Req.

7640 (ster.).

**Galium setaceum** Lam.

6188 (fl.) – 6667 (fl.) – 7453 (fl.) – 7535 (fl.).

*Scrophulariaceae***Kickxia acerbiana** (Boiss.) Täckh. & Boulos

7540 (fl.) – 7516 (fl., fr.).

**Kickxia cf. commutata** (Bernh. ex Reichenb.) Fritsch

6202 (fl.).

**Kickxia floribunda** (Boiss.) Täckh. & Boulos

6650 (fl.) – 7494 (fl., fr.) – 7684 (fl.) – 7711 (fl.).

**Kickxia spartioides** (Brouss.) Janchen

6134 (fl., fr.) – 7490 (fl., fr.).

**Linaria haelava** (Forsskål) Delile

6150 (fl., fr.) – 6205 (fl.) – 6668 (fl., fr.) – 7576B (fl., fr.) – 7585 (fl.).

**Linaria simplex** (Willd.) DC.

7823 (fr.).

**Misopates orontium** (L.) Rafin.  
= *Antirrhinum orontium* L.

6666 (fl., fr.).

**Scrophularia deserti** Delile

6140 (fl.) – 7549 (fl.) – 7574 (fl., fr.).

**Verbascum** cf. *schimperanum* Boiss.

6656 (fl.) – 7561 (fl.).

**Veronica anagallis-aquatica** L.

7614 (fl., fr.). Growing on the edge of a small water pool.

*Solanaceae*

**Hyoscyamus aureus** L.

7646 (fl.).

**Hyoscyamus pusillus** L.

6111 (fl., fr.) – 6217 (fl.) – 6692 (fl.) – 7451 (fl., fr.) – 7586 (fl., fr.) – 7659 (fl., fr.) – 7760 (fl., fr.).

**Lycium shawii** Roemer & Schultes

7469 (fr.) – 7706 (ster.).

**Solanum nigrum** L.

7478 (fl., fr.) – 7563 (fl., fr.) – 7651 (fl., fr.).

**Withania somnifera** L.

7479 (ster.).

*Umbelliferae*

**Anisosciadium isosciadium** Bornm.

7637 (fl.).

*Urticaceae*

**Forsskaolea tenacissima** L.

6113 (fl.) – 6186 (fl.) – 6661 (fl.) – 7518 (fl.).

**Parietaria alsinifolia** Delile

6117 (fl.) – 6136 (fl., fr.) – 6660 (fl., fr.) – 7539 (fl., fr.) – 7723 (fl., fr.).

*Zannichelliaceae*

**Zannichellia palustris** L.

7669 (fr.).

*Zygophyllaceae*

**Fagonia bruguieri** DC. var. *bruguieri*

7509 (fl., fr.).

**Fagonia bruguieri** DC. var. *laxa* Boiss.

6172 (fl., fr.) – 7699 (fl.) – 7745 (fl.). Flowers pink, fading whitish.

**Fagonia glutinosa** Delile var. *grandiflora* Boiss.

7583 (fl., fr.) – 7626 (ster.) – 7697 (fl., fr.) – 7770 (ster.).

**Fagonia mollis** Delile var. *densiglandulosa* Hadidi

7655 (fl.) – 7743 (fl.).

**Fagonia mollis** Delile var. *mollis*

6160 (fl., fr.) – 6657 (fl., fr.) – 7492 (fl., fr.) – 7507 (fl., fr.).

**Fagonia olivieri** DC.

7509 (fl., fr.).

**Peganum harmala** L.

7648 (ster.).

**Tribulus bimucronatus** Viv.

7471 (fl., fr.) – 7514 (fl., fr.).

**Tribulus terrestris** L.

6100 (fl., fr.).

<i>Adiantaceae</i>	<i>Ephedraceae</i>
<b>Adiantum capillus-veneris L.</b>	<b>Ephedra alte C. A. Meyer</b>
7665 (fertile).	7708 (male: fl.; female: fl.).
<i>Sinopteridaceae</i>	<i>Ephedra peduncularis</i> Boiss.
<b>Cheilanthes catanensis (Cosent.) H. P.</b>	6168 (female: fl.).
Fuchs	
7729 (fertile).	
<b>Cheilanthes fragrans (L.) Swartz</b>	
7731 (fertile).	

### Acknowledgements

The authors are indebted to the Keeper and Staff of the Herbarium, Royal Botanic Gardens, Kew, and to the Keeper and Staff of the Botany Department, British Museum (Natural History) for the facilities and help offered during their summer visits in 1974 and 1975. Special thanks are due to the following colleagues who kindly determined many specimens: Mr. C. C. Townsend (*Leguminosae* and *Chenopodiaceae*), Mr. C. Jeffrey (*Compositae*), Mr. S. A. Renvoize (*Gramineae*), Dr. R. M. Harley (*Labiatae*), Mr. R. D. Meikle (different plants), Mr. F. N. Hepper and Dr. I. K. Ferguson (*Scrophulariaceae*) and Professor M. N. El-Hadidi (*Zygophyllaceae*). Thanks are also due to the Director and Staff of the Section of Botany, Natural History Museum, Stockholm, for sending loans to Kew Herbarium in 1974 and 1975. Dr. Jalal Farhat, University of Jordan, kindly provided the geological information. Mr. Walid Jallad helped during the field trips. The map was drawn by Mr. M. Adam of the Geography Department, University of Jordan. Meteorological data were kindly provided by the Meteorological Department, Amman. The research grant offered by the Jordan Research Council is much appreciated.

### REFERENCES

- Ball, P. W. (1972). *Ajuga L.* In: T. G. Tutin, V. H. Heywood, N. A. Burges, D. M. Moore, D. H. Valentine, S. M. Walters & D. A. Webb (eds.), *Flora europaea* 3: 128-129. Cambridge.
- Bor, N. L. (1964). *Gramineae*. In: K. H. Rechinger (ed.), *Flora of lowland Iraq*: 33-119. Weinheim.
- (1968). *Gramineae*. In: C. C. Townsend, E. Guest & A. Al-Rawi (eds.), *Flora of Iraq* 9. Baghdad.
- Bočancev, V. P. (1964a). Čto takoe Cleome arabica Jusl.? *Nov. Sist. Viss. Rast.* 1964: 129-131.
- (1964b). Dopolnenija k flore OAR (Egipet). *Nov. Sist. Viss. Rast.* 1964: 349-378.
- (1968). Zametki o nekotoryh rastenijah Alžirskoj flory. *Nov. Sist. Viss. Rast.* 1968: 233-239.

- Boulos, L. & W. Jallad (1975). Studies on the flora of Jordan. 1. *Diplotaxis villosa* sp. nov. (Cruciferae). *Bot. Not.* 128: 365-367.
- W. Jallad & J. Lahham (1975). Studies on the flora of Jordan. 2. Seven species new to the flora of Jordan. *Bot. Not.* 128: 368-370.
- & J. Lahham (1977). Studies on the flora of Jordan. 3. On the flora of the vicinity of the Aqaba Gulf. *Candollea* 32: 73-80.
- Davis, P. H. (1970, 1975). *Flora of Turkey and the East Aegean Islands* 3, 5. Edinburgh.
- Heyn, C. C., O. Dagan & B. Nachman (1974). The annual *Calendula* species: Taxonomy and relationships. *Israel J. Bot.* 23: 169-201.
- Kers, L. E. (1966). On the identities of *Cleome angustifolia* Forssk. and *Cleome arabica* L. *Acta Horti Berg.* 20: 335-342.
- St John, H. (1965). Revision of *Capparis spinosa* and its African, Asiatic and Pacific relatives. *Micronesia* 2: 25-44.
- Täckholm, V. (1974). *Students' flora of Egypt*, ed. 2. Beirut.
- & L. Boulos (1974). Supplementary notes to Students' flora of Egypt, ed. 2. *Publ. Cairo Univ. Herb.* 5 [“1972”].
- G. Täckholm & M. Drar (1941). Flora of Egypt, 1. *Bull. Fac. Sci. Fouad I Univ.* 17.
- Townsend, C. C. & E. Guest (1974). Leguminosae. In: C. C. Townsend & E. Guest (eds.), *Flora of Iraq* 3. Baghdad.
- Zohary, M. (1960). The species of *Capparis* in the Mediterranean and the Near Eastern countries. *Bull. Res. Council Israel, Sect. D, Bot.* 8/2: 49-64.
- (1962). *Plant life of Palestine*. New York.
- (1966, 1972). *Flora palaestina* 1, 2. Jerusalem.
- (1973). *Geobotanical foundations of the Middle East* 1, 2. Amsterdam.

---

Addresses of the authors: L. B.: Department of Environmental Sciences and Plant Production, International Livestock Centre for Africa, P.O. Box 5689, Addis Ababa, Ethiopia.

J. L.: Department of Biological Sciences, Faculty of Science, University of Jordan, Amman, Jordan.