

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	33 (1978)
Heft:	1
Artikel:	Three new species of Anaphalis (Compositae) from "Flora iranica" area
Autor:	Georgiadou, Elisabeth E.
DOI:	https://doi.org/10.5169/seals-880205

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Three new species of *Anaphalis* (Compositae) from "Flora iranica" area

ELISABETH E. GEORGIADOU

Résumé

GEORGIADOU, E. (1978). Trois nouvelles espèces d'*Anaphalis* (Compositae) dans l'aire de la "Flora iranica". *Candollea* 33: 135-138. En anglais, résumé français.

Trois nouvelles espèces de l'aire de la "Flora iranica" sont décrites dans le genre *Anaphalis*, famille des *Compositae*, tribu des *Inuleae*, sous-tribu des *Gnaphaliinae*: *A. candollei* Georgiadou de l'Afghanistan, *A. staintonii* Georgiadou du Pakistan et du Cachemire ainsi que *A. boissieri* Georgiadou de l'Afghanistan et du Pakistan. Des remarques sont faites quant à leur dimorphisme sexuel et leur tendance à la diœcie.

Abstract

GEORGIADOU, E. (1978). Three new species of *Anaphalis* (Compositae) from the "Flora iranica" area. *Candollea* 33: 135-138. In English, French abstract.

Three new species of *Anaphalis* (Compositae), tribe *Inuleae*, subtribe *Gnaphaliinae*, are described from "Flora iranica" area. *A. candollei* Georgiadou from Afghanistan, *A. staintonii* Georgiadou from Pakistan and Kashmir, and *A. boissieri* Georgiadou from Afghanistan and Pakistan. Notes on subdioecism and sex-linked dimorphism are given.

Introduction

In the course of preparation of the account of *Anaphalis* for "Flora iranica", three new species came to light and are described here.

Anaphalis has its centre of distribution in the mountains of Central Asia. The genus reaches its western limit in the mountains of Northwest Pakistan, in the adjacent section of East Afghanistan and in the Pamir-Alai mountain system of soviet Central Asia.

According to DRURY (1970), the genus is subdioecious, although GRIERSON (1972) pointed out that this is not absolute, as some species, such as the Ceylon species and some of the Sino-Himalayan are not. In the three new taxa, the predominantly hermaphrodite capitula contain exclusively the hermaphrodite (functionally male) type of florets, whereas in the predominantly female capitula of *A. candollei* and *A. boissieri*, there are some hermaphrodite florets in the centre of the capitulum. Female capitula were not seen in *A. staintonii*. This was also noted by GRIERSON (1972) in *A. aureo-punctata* Lingelsh. & Borza, *A. bicolor* Franchet, *A. griffithii* Hooker fil., *A. hancockii* Maxim., *A. larium* Hand.-Mazz. and *A. likianensis* Chang.

Sex-linked dimorphism in *Anaphalis* was long ago observed by BEAUVERD (1910, 1913) in the florets, achenes and pappus. *A. candollei* and *A. boissieri* have filiform female florets whereas the hermaphrodite ones are tubular. The achenes of the female florets, where seen, are well developed and appear fertile, whereas in the functionally male capitula the hermaphrodite florets have reduced, rudimentary and sterile achenes. Finally the pappus of the female florets is finer with acute bristles, whereas that of the hermaphrodite florets is wider and clavicate, except in *A. boissieri* which has acute narrow bristles.

In habit many Asiatic species of *Anaphalis* resemble *Helichrysum* which, however, has either homogamous capitula with fertile hermaphrodite florets or heterogamous with peripheral female florets and central hermaphrodite florets, the latter predominating in the capitulum. Sex-linked dimorphism is unknown in the representatives of *Helichrysum* capitula at least from W. Asia and the Mediterranean region.

Anaphalis is taxonomically an extremely complicated genus. A thorough revision is urgently needed. Such an enterprise would be hampered by the conflicting interpretation of such widespread species as *A. virgata* Thomson.

In view of the variable sex distribution, a clearer indication of affinities can probably be obtained by the more consistent evaluation of relative stable vegetative characters, such as growth, leaf arrangement and direction, presence or absence of sterile axillary shoots.

1. *Anaphalis candollei* Georgiadou, spec. nova.

Holotypus: Breckle 3118, W.

Suffrutex multicaulis. Caules 20-30 cm alti laxe floccoso-tomentosi tota longitudine foliati, basi ramuligera. Folia supra virescentia, laxe floccoso-tomentosa, subtus dense albido-tomentosa, glandulis paucis indumento occultis, folia intermedia maxime circiter 20 x 6.5 mm, plana, margine leviter revoluta, basi late sessilia, apice breviter attenuata et apiculata. Folia superiora paulo decrescentia, synflorescentiam fulcrantia, quam inferiora proportione angustiora. Synflorescentia laxe corymbosa circiter 19.2-47.7 mm diametro. Capitula 7-30, subdioecia disciformia vel discoidea. Pedicelli 4-6.2 mm albo-tomentosi, parce glandulosi. Involucrum circiter 5-6.5 mm, flores paulo superans. Phylla 40-45, 6-7-seriata, alba, opaca, ± acuta, basi brunnescentia tomentosa. Flores in capitulis hermaphroditis (functione masculi) 29-31, in capitulo subfoemineo circiter 43 foeminei, 4 hermaphroditici. Achenia in capitulo submasculo 0.5 mm longa, oblonga sterilia. In capitula subfoemineo achenia florium foemineorum 1.3 mm longa, glabra. Pappus albus, uniseriatus, radii acheniorum foemineorum capillares ad apices non incrassati. Radii acheniorum hermaphroditorum clavicolati ad apicem incrassati. — Differt ab *A. boissieri* Georgiadou achenio foemineo glabro, involuci phyllis 40-45, 6-7 seriatis.

Afghanistan: E: Kunar: Bashgal, Garm Chashma, 3250 m, *Rasoul* in Breckle 3518! In valle Suyengal, affluentis fluvii Bashgal, 3600 m, 4.9.1969, Breckle 3118! (holotypus). Barge-Matal, in valle orientali, 2450-2650 m, Anders 5101! Darrah-i Pengar ad conjunctionem vallis Pushal, 3350-3400 m, Podlech 16388!

2. *Anaphalis staintonii* Georgiadou, spec. nova.

Typus: Stainton 3032 (holotypus BM!; isotypus G!).

Suffrutex multicaulis. Caules 35-40 cm alti, erecti vel arcuati, tota longitudine foliati, appresse canescenti-tomentosi, parce glandulosi. Folia a basi usque ad apicem omnia aequalia circiter 22 x 1.7 mm, anguste linearia, acuta, arcte convoluta, utrimque subaequaliter tomentosa, inferiora foliorum fasciculos nonnullos minutos proferentia, superiora inflorescentiam attingentia. Synflorscentia longe ramosa, laxe corymbosa, 22-57 mm diametro, 15-57-cephala. Pedicelli 1.5-7.5 mm longi, albo-tomentosi et glandulosi. Involucrum 5 x 7.5 mm. Phylla 32-41, 5-6-seriata, acuta, irregulariter laxe disposita, inferne coriacea, superne scariosa, amoene rosea, interiora quam exteriora duplo longiora. Flores hermaphroditi in capitulis hermaphroditis (functione masculi) 26-36. Achenia in capitulo submasculo circiter 0.5 mm longa, oblonga, omnia sterilia. Pappus albus uniseriatus, scabridulus, radii clavulati ad apicem incrassati. Capitula foeminea non visa. — Habitu *A. virgatae* Thomson ex C. B. Clarke similis! Differt ab ea inflorescentia ampla, involucris roseis, pappo claviculato.

Pakistan: Chitral: Baroghil Pass, 1100 ft., 28.7.1858, *Stainton* 3032! Lager Rumosto, in Bachsand, 2900 m, *Kerstan* 1570! *Kashmir:* Kar-Gah trans Gilgit, 2600-2900 m, Schmid 1997!

3. *Anaphalis boissieri* Georgiadou, spec. nova.

Typus: Gilli 3984, W!

Suffrutex multicaulis, omnino indumento subfloccoso-tomentoso canescens vel albescens, indumento in partibus inferioribus magis floccoso, in superioribus magis appresso. Caules floriferi 20-35 cm alti, erecti vel arcuato-ascendentes, tota longitudine foliati, prope basin ramulos steriles proferentes. Folia omnia indumento supra laxiore atro- vel sordide canescenti-viridia, subtus indumento denso tomentoso appresso albida, inferiora usque ad 20-25 x 4-7 mm, lanceolata, obtusiuscula vel acutiuscula, plana, quam folia caulina superiora semper evidenter latiora et proportione breviora, folia caulina media et superiora basi latiuscula vel sat breviter attenuata sessilia, lanceolata usque anguste linear-lanceolata, acuta, margine ± revoluta, angulo ± recto patentia. Synflorescentia 28-35 mm diametro, ± 15-cephala, composite corymboso-paniculata, plerumque densiuscula. Rami inferiores plerumque 10-20 mm longi. Pedicelli albo-tomentosi, involucro breviores vel paulo longiores. Involucrum depresso-semiglobosum, floriferum, ± 5 mm, fructiferum usque ad 8 mm diametro. Phylla 28-35, 5-6-seriata, alba, opaca, basi saepe rubescens vel brunnescentia, longitudine inter se non valde diversa. Flores hermaphroditi in capitulis hermaphroditis 24, in capitulis subfoemineis circiter 20, hermaphroditi 3. Achenia in capitulo submasculo 0.5 mm longa sterilia, in capitulis subfoemineis, 1.5 mm longa, papillosa. Pappus albus, uniseriatus, radii scabridi, acuti. — Proxima *A. darvasicae* Borissova, a qua differt involucri phyllis 5-6-nec 3-4-seriatis, papii radiis acheniorum hermaphroditorum acutis nec clavato-dilatatis.

Afghanistan: E: In fissuris rupium gneiss, faucium Nedschrab inter Gulbahar et Sarobi, 3100-3240 m, 13.9.1951, *Gilli* 3984!, 3985! Logar: In valle fluvii Cacobay in saxosis humidis, 3100 m, *Lalande R* 1072-E 12! *Pakistan*: Kurram: Parachinar, *Afendi* 40!

Acknowledgements

The author is most grateful to Prof. Dr. K. H. Rechinger for his collaboration during the preparation of the account of *Anaphalis* and *Helichrysum* for "Flora iranica". Thanks are also due to the Conservateurs and technical staff of the Conservatoire et Jardin botaniques de Genève, for the facilities and help offered during her visits there.

REFERENCES

- BEAUVERD, G. (1910). Recherches sur la tribu des Gnaphaliées. *Bull. Soc. Bot. Genève*, sér. 2, 2: 207-252.
— (1913). La constante générique des Anaphalis. *Bull. Soc. Bot. Genève*, sér. 2, 5: 146-147.
DRURY, D. G. (1970). A fresh approach to the classification of the genus *Gnaphalium* with particular reference to the species present in New Zealand. *New Zealand J. Bot.* 8: 222-248.
GRIERSON, A. J. C. (1972). A new species of *Anaphalis* (Compositae) from Mexico. *Notes Roy. Bot. Gard. Edinburgh* 31: 389-392.