

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	38 (1983)
<b>Heft:</b>	1
<b>Artikel:</b>	Studies on the flora of Jordan : 10. Nine new species to the Flora of Jordan
<b>Autor:</b>	Al-Eisawi, Dawud
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-879869">https://doi.org/10.5169/seals-879869</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 01.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Studies on the flora of Jordan. 10. Nine new species to the Flora of Jordan

DAWUD AL-EISAWI

## RÉSUMÉ

AL-EISAWI, D. (1983). Etude sur la flore de Jordanie. 10. Neuf nouvelles espèces pour la flore de Jordanie. *Candollea* 38: 359-364. En anglais, résumé français.

Neuf espèces sont repérées pour la première fois en Jordanie, ce sont: *Anisosciadium lanatum* Boiss. et *Opopanax hispidum* (Friv.) Griseb. (Umbelliferae), *Ambrosia maritima* L. et *Zoegea purpurea* Fres. (Compositae), *Euclidium syriacum* (L.) R. Br. (Cruciferae), *Lappula barbata* (Bieb.) Gürke (Boraginaceae), *Calligonum tetrapterum* Taub. & Spach (Polygonaceae), *Teucrium rigidum* Benth. (Labiatae) et *Cytinus hypocistis* (L.) L. (Cytinaceae ou Rafflesiaceae).

## ABSTRACT

AL-EISAWI, D. (1983). Studies on the flora of Jordan. 10. Nine new species to the Flora of Jordan. *Candollea* 38: 359-364. In English, French abstract.

Nine species are recorded for the first time in Jordan, these are: *Anisosciadium lanatum* Boiss. and *Opopanax hispidum* (Friv.) Griseb. (Umbelliferae), *Ambrosia maritima* L. and *Zoegea purpurea* Fres. (Compositae), *Euclidium syriacum* (L.) R. Br. (Cruciferae), *Lappula barbata* (Bieb.) Gürke (Boraginaceae), *Calligonum tetrapterum* Taub. & Spach (Polygonaceae), *Teucrium rigidum* Benth. (Labiatae), and *Cytinus hypocistis* (L.) L. (Cytinaceae or Rafflesiaceae).

The nine species reported in this paper belong to nine different genera of which five are recorded as new to Jordan. The genera are: *Opopanax*, *Ambrosia*, *Zoegea*, *Euclidium* and *Cytinus*. All the specimens of the nine species are deposited at the University of Jordan Herbarium, Amman.

**Anisosciadium lanatum** Boiss.

The record of this species *A. lanatum* in addition to the already known species *A. isociadium* will represent two out of three species belonging to the genus *Anisosciadium* (the third species is *A. orientalis*). HEDGE & LAMOND (1973) in their review of the tribe *Echinophoreae* do not show in the map of distribution of the genus *Anisosciadium* that *A. lanatum* occurs in Jordan. ZOHARY (1972), POST & DINSMORE (1932) as well do not enumerate this species from Jordan. Although this species was collected from eastern desert along the borders with Saudi Arabia. MIGAHID (1978) doesn't mention either that this species occurs in Saudi Arabia.

*Specimen collected from Jordan*

Eastern desert: Al-Inab, 75 km S.E. of Al-Jafr, 20 June 1980, *Rasmi Jarrar D 8615*.

**Opopanax hispidum** (Friv.) Griseb.

CHAMBERLAIN (1972) gives the geographical distribution as Italy, Balkans, Syria, Cyprus, N. Iraq, Soviet Armenia, W. Iran without mentioning Jordan or Palestine. POST & DINSMORE (1932) enumerate this species as known from Palestine and not Jordan, but ZOHARY (1972) does not describe this genus from Palestine or Jordan.

*Specimen collected from Jordan*

Ajloum: Ischtafaina; in a *Quercus* and *Arbutus* natural forest, 8 June 1979, *D. Al-Eisawi & S. Al-Eisawi 7599*. The plant was seen in the same locality several times in the following years.

**Ambrosia maritima** L.

POST & DINSMORE (1932) as well as FEINBRUN-DOTHAN (1978) give the geographical distribution of this species as it occurs in Palestine along the coastal area mainly and they don't include Jordan as a place of occurrence for this mediterranean element. Recently this species was collected from the eastern side of the southern end of the Dead Sea in Jordan near a cultivated field close to the sea shore. This occurrence makes this species as a new element to the tropical vegetation in the area.

*Specimen collected from Jordan*

Ghor Al-Hadeetha; the eastern side of the southern end of the Dead Sea, near Ghor al-Mazra'a, 14 March 1979, *D. Al-Eisawi & R. Jarrar 4756*.




Fig. 1. — The distribution of the new taxa within Jordan.

**Zoegea purpurea** Fres.

This species is very rare in the area. POST & DINSMORE (1932) mention this species from two localities: Sinai and the Dead Sea. FEINBRUN-DOTHAN (1978) confirms the occurrence of this species in Palestine but from only C. & S. Negev. TÄCKHOLM (1974) gave the distribution as very rare and known only from Sinai. MIGAHID (1978) shows that the distribution in Saudi Arabia is only restricted to Najd. Finally BOULOS & AL-EISAWI (1977) do not record this species from Ras En-Naqab. This species was collected two years later from Ras En-Naqab area and identified at Kew by C. Jeffrey

*Specimen collected from Jordan*

Ras En-Naqab: 30 km W. of Dlatha, at the end of the old road which leads to Gharandal in Wadi Araba, about 10 km E. of Gharandal, 2 May 1979, *D. Al-Eisawi* 7475.

**Euclidium syriacum** (L.) R. Br.

This species according to POST & DINSMORE (1932) occurs in the three localities Aleppo, Hums and Hauran in Syria only and is not recorded from Jordan or Palestine. ZOHARY (1966) does not enumerate this species from Palestine and Jordan either. HEDGE (1965) gives the distribution of this species as S. Russia, S.W. Asia and Central Asia. The occurrence of this species in three well separated localities in Jordan may suggest that it was distributed from Hauran (S. Syria) through the eastern side of Jordan from the north down to the south in Petra.

*Specimens collected from Jordan*

Zarka: Zarka river; Bani Hassan, near Sukhnah, 18 March 1978, *D. Al-Eisawi & K. Abd-Alla* 3075.

Amman: Al-Jubaiha; University housing area, near the university fence, 11 April 1978, *D. Al-Eisawi* 3546.

Petra: Al-Hiesha Al-Byda, 5 km N. Petra near a water spring, 23 April 1981, *O. Abul-Rub & D. Al-Eisawi* 151.

**Lappula barbata** (M. Bieb.) Gürke

The specimens were identified by R. Meikle at Kew and collected from S. Jordan near Tafila under an old protected stand of wild *Cupressus sempervirens*. Distribution is given by POST & DINSMORE (1932) as known from two localities only, Damascus in Syria and Báalbak in Lebanon. Therefore it was not known from Palestine or Jordan. Similarly FEINBRUN-DOTHAN

(1978) does not enumerate this species from Palestine or Jordan. EDMONDSON (1978) gives the distribution of this species as Balkans, S. Russia, Caucasia, N., N.W., W. & S. Iran, Khorassan, Afghanistan.

*Specimen collected from Jordan*

Tafila, 25 km S. Tafila along the road to Shubak, near Ain Lahtha (Khonag Al-Arz), 11 April 1979, *D. Al-Eisawi & R. Jarrar 6057*. The species was seen in the same locality in the following years.

**Calligonum tetrapterum** Taub. & Spach

This species was collected from the eastern desert of Jordan near the borders with Saudi Arabia and Iraq. POST & DINSMORE (1932), ZOHARY (1966) and MIGAHID (1978) do not record this species from Palestine, Jordan or Saudi Arabia. BOISSIER (1879) gave the distribution as known from Baghad and Aleppo. RECHINGER (1964) gives the distribution from few localities in Iraq especially Al-Fallujah desert.

*Specimen collected from Jordan*

Eastern desert: Al-Wisad, 5 km E. of Al-Wisad, near Qa Al-Gathoum, 4 April 1979, *D. Al-Eisawi 5608*.

**Teucrium rigidum** Benth.

This is a rare species in the area. The distribution as given by BOISSIER (1879) is Assyria and Basian. POST & DINSMORE (1932) as well FEINBRUNDOTHAN (1978), do not enumerate this species. The specimen was collected from the eastern desert near the borders with Saudi Arabia, even though MIGAHID (1978) does not enumerate this species either.

*Specimen collected from Jordan*

Eastern desert: along the way to Alwisad; 20 km W. Alwisad, 3 April 1979, *D. Al-Eisawi 5594*.

**Cytinus hypocistis** (L.) L.

POST & DINSMORE (1932) give the geographical distribution of this species as Lebanon and Palestine in the Carmel mountain. FEINBRUNDOTHAN (1978) as well says that it is a rare species and known only from Carmel. The specimen collected from Jordan was found as a parasite on *Cistus* roots.

*Specimen collected from Jordan*

Jarash: Dibbeen National Park, 24 April 1980, *K. Shukri, K. Dieb & I. Ali* s.n.

## ACKNOWLEDGEMENT

I would like to thank the University of Jordan for their financial support. I'm greatly indebted to the Director, Curator and Staff of the Herbarium and Library, Royal Botanic Gardens, Kew, for the facilities and help. I would like to thank C. C. Townsend, R. Meikle and C. Jeffrey for their special help. Finally I would like to thank the Institute of botany, Munich, especially Prof. Merxmüller and Prof. Podlech for their help during the preparation of this manuscript.

## REFERENCES

- BOISSIER, E. (1879). *Flora Orientalis IV*. Geneva.
- BOULOS, L. & D. M. AL-EISAWI (1977). Studies on the flora of Jordan, 6. On the flora of Ras en Naqb. *Candollea* 32(1): 111-120.
- CHAMBERLAIN, D. F. (1973). Opopanax in: DAVIS, P. H., *Flora of Turkey* 4: 471-473. Edinburgh.
- EDMONSON, J. R. (1978). Lappula in: DAVIS, P. H., *Flora of Turkey* 6: 255-261. Edinburgh.
- FEINBRUN-DOTHAN, N. (1978). *Flora Palaestina*, 3. Jerusalem.
- HEDGE, I. C. (1965). Euclidium in: DAVIS, P. H., *Flora of Turkey* 6: 255-261. Edinburgh.
- & J. M. LAMOND (1973). A review of the tribe Echinophoreae (Umbelliferae). *Notes Roy. Bot. Gard. Edinb.* 32: 167-188.
- MIGAHID, A. M. (1978). *Flora of Saudi Arabia*, 1 & 2. Riyad University.
- POST, G. E. & J. E. DINSMORE (1932, 1933). *Flora of Syria, Palestine and Sinai*, 1 & 2. Beirut.
- RECHINGER, K. H. (1964) *Flora of Lowland Iraq*. Weinheim.
- TÄCKHOLM, V. (1974). *Student Flora of Egypt*, ed. 2. Beirut.
- TOWNSEND, C. C. (1974). *Flora of Iraq*, ed. 3. Baghdad.
- ZOHARY, M. (1966, 1972). *Flora Palestina*, 1 & 2. Jerusalem.