

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	40 (1985)
Heft:	1
Artikel:	Minuartia favargeri : a new species from Peloponnesus (Greece)
Autor:	Iatrou, Gregory / Georgiadis, Theodore
DOI:	https://doi.org/10.5169/seals-879774

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Minuartia favargeri: a new species from Peloponnesus (Greece)

GREGORY IATROÚ
&
THEODORE GEORGIADIS

RÉSUMÉ

IATROÚ, G. & T. GEORGIADIS (1985). *Minuartia favargeri*: une espèce nouvelle du Péloponnèse (Grèce). *Candollea* 40: 129-138. En anglais, résumé français.

Minuartia favargeri a été trouvée sur des rochers calcaires dans la partie orientale du Mt. Parnon, en Arcadie (Péloponnèse). Cette espèce, qui appartient à *Minuartia* sect. *Acutiflorae* ser. *Pichleriae* est en rapport avec l'endémique du sud du Péloponnèse *M. pichleri* et l'endémique du sud de l'Anatolie *M. rimarum*. Les nombres chromosomiques ($2n = 26$) et les caryotypes des deux endémiques grecques sont nouveaux. Les auteurs présentent aussi le nombre total des taxa endémiques grecs du genre *Minuartia* ainsi que leur distribution.

ABSTRACT

IATROÚ, G. & T. GEORGIADIS (1985). *Minuartia favargeri*: a new species from Peloponnesus (Greece). *Candollea* 40: 129-138. In English, French abstract.

Minuartia favargeri is described from calcareous rocks of the E. side of Mt. Parnon, in the province of Arkadhia, Peloponnesus. This species belonging to *Minuartia* sect. *Acutiflorae* ser. *Pichleriae* is related to the Peloponnesian endemic *M. pichleri* and to the S. Anatolian endemic *M. rimarum*. The chromosome number ($2n = 26$) and the karyotypes of the two Greek endemic species are new. The total number of the Greek endemic taxa of *Minuartia* as well as their distribution are presented.

Minuartia favargeri Iatroú & Georgiadis, sp. nova (Fig. 1).

Typus: Greece: Peloponnesus, prov. Arkadhia, in declibus orientalibus montis Parnon; prope Monasterium Elona, in saxosis calcareis alt. ca. 550 m, 7.VII.1983, Gr. Iatroú & E. Kokkinos 1915 (holotypus UPA).

Planta perennis, dense pulvinato-caespitosa. Caules erecti vel adscendententes 3-8 cm alti, cum foliis dense glandulosso-pilosis; folia basalia 3-nervia 8-10(-12) mm longa, 1 mm lata. Flores (2)-3-6(-8) in cymas terminalis dispositi. Pedicelli dense glandulosso-pilos, calycibus campanulatis 3-vel 5-plo longiores. Sepala ovato-lanceolata, dense glandulosso-pilosa, anguste scarioso marginata 2.5-3 mm longa 1 mm lata. Petala cuneato-oblongo-ovovata 4.5-5 mm longa calyce sesqui vel duplo longiora. Capsula calyce sesquilonior. Glandes staminales 0.1-0.2 mm longae. Semina dorso tenuiter tuberculata.

Numerus chromosomatum: 2n = 26.

Ex affinitate *M. pichleri* (Boiss.) Maire & Petitmengin et *M. rimarum* (Boiss. & Bal.) Mattf.

Densely caespitose, cushion-forming, perennial. The whole plant densely glandular pubescent. Stems 3-8 cm high, erect. Basal leaves 8-10(-12) mm long, 1-1.2 mm broad, 3-veined, somewhat falcate. Cymes (2)-3-6(-8) flowered; pedicels 3-5 times as long as sepals. Calyx campanulate with sepals ovate-lanceolate 2.5-3 × 1 mm, with narrow membranous margins. Petals 4.5-5 mm long, at least 1½ times as long as sepals. Capsule very obviously exserted calyx, about 1½ times as long as calyx. Staminal glands 0.15-0.2 mm long, forming a short finger-like process with an apical nectar pit. Seeds black, slightly tuberculate on the back. Flowering period from late May to early July.




Fig. 1. — *Minuartia favargeri* (holotypus).

Minuartia favargeri is endemic on Mt. Parnon of E. Peloponnesus. It is a typical chasmophyte growing in rock crevices of hard limestone, at an altitude of 500-600 m above sea level. The station in which it prospers consists of a rocky limestone ecosystem, just before the Monastery of Elona, in the gorge which starts from the village of Leonidion and ends before the village of Kosmas. In this ecosystem predominate the following plants: *Asperula boryana* (Walpers) Ehrend., *Asperula taygetea* Boiss. & Heldr., *Asperula elonea* Iatrou & Georgiadis, *Athamantha macedonica* (L.) Sprengel subsp. *arachnoidea* (Boiss. & Heldr.) Tutin, *Stachys chrysanthra* Boiss. & Heldr., *Trachelium asperuloides* Boiss. & Orph., *Petrorhagia grandiflora* Iatrou, *Centaurea raphanina* Sibth. & Sm. subsp. *mixta* (DC.) Runemark, *Onosma graeca* Boiss., *Ceterach officinarum* DC., *Micromeria juliana* (L.) Bentham ex Reichenb., *Briza minor* L., *Parietaria* sp., *Galium* sp., *Sedum* sp., *Inula* sp., *Odontites* sp. The area around this station belongs to the sub-humid Mediterranean bioclimatic zone, with a mean annual rainfall of 600-800 mm and a mean annual sunlight of 2800 hrs.

M. favargeri and *M. pichleri* are closely related to each other and have the same chromosome number, $2n = 26$. They are schizoendemics (FAVARGER, 1964). *M. rimarum* is closely related and geographically vicarious with both. The two former species are restricted to Peloponnesus and the latter is endemic in C. Anatolia, with its main centre of distribution in the Taurus mountain range.

Taxon	Locality	Collector	Chromosome number
<i>M. favargeri</i>	Greece, Peloponnesus: Mt. Parnon, Monasterium Elona, ca. 550 m	Iatrou & Kokkinos 1915	$2n = 26$
<i>M. pichleri</i>	Greece, Peloponnesus: Mt. Taygetos, Langada gorge, ca. 500 m	Iatrou & Georgiadis 2142	$2n = 26$
<i>M. rimarum</i> var. <i>rimarum</i>	Turkey, Izmir: Ödemiş, Bozdağ Boz Dağ	Contandriopoulos 68-196	$n = 13$ (ÇELEBIOĞLU & FAVARGER, 1982)
<i>M. rimarum</i> var. <i>rimarum</i>	Turkey, Kayseri: Erciyas Dağı, alt. 2200 m	Welter & Hébert 34 A. W.	$n = 13$ (ÇELEBIOĞLU & FAVARGER, 1982)
<i>M. rimarum</i> var. <i>multiflora</i>	Turkey, Konya: Ermerek, Yellibel, alt. 2000 m	Contandriopoulos	$n = 13$ (ÇELEBIOĞLU & FAVARGER, 1982)

Table 1. — Collecting sites of populations used for the cytological study. The Turkish material is according to ÇELEBIOĞLU & FAVARGER, 1982.




Fig. 2. — Mitotic chromosome plates: **a**, *Minuartia favargeri*; **b**, *Minuartia pichleri*.




Fig. 3. — Differences between basal leaves, hair-covering, petals, sepals and staminal glands.

A. *Minuartia favargeri*: **a**, basal leaf; **b**, hair-covering; **c**, petals; **d**, sepal; **e**, staminal gland.

B. *Minuartia pichleri*: **a**, basal leaf; **b**, hair-covering; **c**, petals; **d**, sepal; **e**, staminal glands.




Fig. 4. — Differences between basal leaves, hair-covering, petals, sepals and staminal glands.

B. *Minuartia rimarum* var. *rimarum*: a, basal leaf; b, hair-covering; c, petals; d, sepal; e, staminal gland.

D. *Minuartia rimarum* var. *multiflora*: a, basal leaf; b, hair-covering; c, petals; d, sepal; e, staminal gland.




Fig. 5. — SEM photographs of the seeds (a, b) *Minuartia pichleri*: a, whole seed ($\times 100$); b, outer cells of the median zone ($\times 240$).

(c, d) *Minuartia favargeri*: c, whole seed ($\times 100$); d, outer cells of the median zone ($\times 240$).




Fig. 6. — Map of geographical distribution of the taxa endemic in Greece of the genus *Minuartia*. 1. *Minuartia wettsteinii* Mattf. 2. *M. verna* (L.) Hiern subsp. *idaea* (Halacsy) Hayek. 3. *M. favargeri* Iatrou & Georgiadis. 4. *M. pichleri* (Boiss.) Maire & Petitmengin. 5. *M. eurytanica* (Boiss.) & Heldr.) Hand.-Mazz. 6. *M. pseudosaxifraga* (Mattf.) Greuter & Burdet. 7. *M. confusa* (Heldr. & Sart.) Maire & Petitmengin.

<i>Minuartia favargeri</i>	<i>Minuartia pichleri</i>	<i>Minuartia rimarum</i>
Perennial plants, cushion-forming, densely caespitose glandular pubescent.	Perennial plants, cushion-forming, densely caespitose glandular pubescent.	Perennial, mat- to cushion-forming glandular or densely glandular pubescent.
Stems 3-8 cm high, erect.	Stems 3-10 cm high, erect.	Stems up to 8 cm high erect.
Leaves 3-veined, basal 8-10(-12) × 1 mm.	Leaves 3-veined, basal 5.5-6 × 1 mm.	Leaves 3-veined, basal 4-8 × 1 mm.
Cymes (2-)3-6(-8) flowered, pedicels 3-5 times as long as sepals.	Cymes (1)2-4(6) flowered, pedicels 3-5 times as long as sepals.	Cymes 1-5 flowered, pedicels 3-5 times as long as sepals.
Calyx campanulate with sepals ovate-lanceolate 2.5-3 × 1 mm, with narrow membranous margin, densely glandular pubescent.	Calyx funnelled with sepals oblong-lanceolate 4.5-5 × 1 mm, with broad membranous margin, glandular pubescent.	Calyx ± campanulate with sepals ovate-lanceolate, acuminate, 4.5 × 1 mm, with ± broad membranous margin, glandular or densely glandular pubescent.
Petals 4.5-5 mm long, at least 1 1/2 times as long as sepals.	Petals 7-7.5 mm long, up to 1 1/2 times as long as sepals.	Petals 5.5-8.5 mm long, 1 1/2 times as long as sepals.
Capsule very obviously exserted calyx (about 1 1/2 times as long as calyx).	Capsule slightly exceeds calyx.	Capsule shorter than calyx.
Staminal glands 0.15-0.2 mm, forming a very short finger-like process with an apical nectar pit.	Staminal glands 0.5-0.75 mm, forming a long finger-like process with an apical nectar pit.	Staminal glands c.o. 1 mm consisting of a sessil nectar pit.
Seeds, black, slightly tuberculate on the back.	Seeds dark brown to black acutely tuberculate on the back.	Seeds, black, slightly tuberculate on the back.
Chromosome number $2n = 26$.	Chromosome number $2n = 26$.	Chromosome number $2n = 26$.
Distribution: endemic in Greece, Peloponnesus (Parnon, ca. 550 m.	Distribution: endemic in Greece, Peloponnesus (Kyllene, ca. 1300 m, Taygetos, ca. 500 m).	Distribution: endemic in Turkey (Cilicia, Erzincan, Antalya, Maras Sivas, 1750-2900 m).

Table 2.—Differences between *Minuartia favargeri* and related species. Measurements are based on herbarium material checked by the authors.

Genus <i>Minuartia</i>		<i>x or x'</i>
subgenus I	<i>Rhodalsine</i> (J. Gay) Graebner	9
subgenus II	<i>Spergella</i> (Fenzl) McNeill	11
subgenus III	<i>Minuartia</i> (L.) Hiern	13 (12)
	Sect. <i>Spectabiles</i>	15
	Sect. <i>Plurinerviae</i>	16
	Sect. <i>Lanceolatae</i> (Fenzl) Graebner	16
	ser. <i>Graminifoliae</i>	18
	ser. <i>Diantifoliae</i>	18
	ser. <i>Lanceolatae</i>	18
	ser. <i>Grigneenses</i>	18
	Sect. <i>Aretioideae</i> Mattf.	18
	Sect. <i>Sclerophylla</i>	15
	Sect. <i>Acutiflorae</i> (Fenzl) Hayek	13
	Sect. <i>Tryphane</i> (Fenzl) Hayek	12 (13)
	Sect. <i>Alsinanthe</i> (Fenzl) Graebner	15
	Sect. <i>Minuartia</i>	15 (14 & 13)
	ser. <i>Montanae</i>	15
	ser. <i>Minuartia</i>	15
	ser. <i>Leucocephalae</i>	15
	ser. <i>Setaceae</i>	15 (14)
	ser. <i>Xeralsine</i>	15 (14)
	ser. <i>Campestres</i>	15
	Sect. <i>Sabulina</i> (Reichb.) Graebner	11, 12, 23, 35

Tab. 3. — The basic chromosome numbers (*x* or *x'*) of sections and series of genus *Minuartia* according to ÇELEBIOĞLU & FAVARGER, 1982.

Cytology

The cytological study is based on material collected in nature and cultivated in pots outdoors. For the study of the mitotic chromosomes, actively growing root tips were pretreated in a saturated solution of 1-Bromonaphthalene, 4°C, 6 hrs.; fixed in carnoy's solution, 4°C, 24 hrs.; hydrolized in 1N HCl, 60°C, 12 min.; stained with Feulgen reagent, room temperature, 3 hrs.; treated in 4% pectinase, 20 min. Squashes were prepared and made permanent according to ÖSTERGREN & HENEEN (1962). The drawings were made with the aid of a camera lucida.

M. favargeri and *M. pichleri* have the same chromosome number $2n = 26$. This number coincides with the ones, studied before, of all the other species belonging to the same section (FAVARGER, 1962a, 1967, ÇELEBIOĞLU & FAVARGER, 1982). The chromosomes are small 1-2 μ m, but with well visible constrictions. Due to their small size all the chromosomes appear very similar and seem to belong to the m, sm categories (LEVAN & al., 1964), and are difficult to identify. This makes the observation of any differences in the karyotype almost impossible (Fig. 2a, b). For the same reasons morphological details could not be studied, except for two pairs of satellites chromosomes present in both karyotypes.

This new species belongs to *Minuartia* (L.) Hiern sect. *Acutiflorae* (Fenzl) Hayek ser. *Pichleriae* Mattf. (McNEILL, 1963) and is undoubtedly related to *M. pichleri* endemic also in Peloponnesus (HALIDAY, 1964), as well as to *M. rimarum* endemic in S. Anatolia (McNEILL, 1967), but it differs from both in a number of characters (Tabl. 1). The most important differences between *M. favargeri* and the other two species are the size of the leaves and petals, the shape and the size of sepals, the exsertion of the capsule from the calyx, the size of the staminal glands and the morphology of the seeds. The leaves of *M. favargeri* are 8-10(-12) mm, almost double the size of the leaves of *M. pichleri* (5.5-6 mm) and much longer than the ones of *M. rimarum* (4-8 mm). The petals of *M. favargeri* are 4.5-5 mm much smaller than the ones of the other two, being for *M. pichleri* 7-7.5 mm and for *M. rimarum* 5.5-8.5 mm. The sepals of *M. favargeri* are 2.5-3 mm, almost half the size of the ones of *M. pichleri* (4.5-5 mm) and *M. rimarum* (4-5 mm). The capsule in *M. favargeri* exceeds very obviously the calyx (1 $\frac{1}{2}$ times as long as calyx), in *M. pichleri* is equal to the calyx or slightly exceeds it; in *M. rimarum* the capsule is shorter than the calyx. The staminal glands in *M. favargeri* have an intermediate size (0.15-0.2 mm) between the very long ones of *M. pichleri* (0.5-0.7 mm) and the almost sessile ones of *M. rimarum* (ca. 0.1 mm).

The chief characters for the comparative study of the seeds are: the ornamentation of the outer cells of the median zone, the shape and the ornamentation of the papillae and the degree of differentiation of the cells of the mediane zone (ÇELEBIOĞLU & al., 1983). All the above characters

have been studied by SEM and proved very different between *M. favargeri* and *M. pichleri*, with as most striking difference, the ornamentation of the outer cells of the median zone (base) (Fig. 3b, c). Unfortunately we did not study seeds of *M. rimarum*.

In genus *Minuartia* each section, and sometimes the series as they have been divided by MATTFELD (1922) & McNEILL (1962, 1963), is characterized by a unique basic chromosome number, usually different between them (Tabl. 3).

The cytological independance of each section and series confirms the antiquity of this genus. The different sections are all very isolated from each other and own their origin to allopolyploidy or dispoloidy effects, what has been called secondary swarming ("foisonnement secondaire", ÇELEBIOĞLU & FAVARGER, 1982). According to the same authors (1982) the section *Acutiflorae* ($n = 13$) and probably also the sect. *Spectabiles*, which are considered to be the less evolved groups of the genus (McNEILL, 1962), seem to have been derived, by amphidiploidy, from the primitive numbers $x = 6$ and $x = 7$ ($6 + 7 \rightarrow n = 13$). MATTFELD (1922) believes that sect. *Acutiflorae* is derived from a primitive type of *Univerviae*. In fact one of its primitive basic numbers ($x = 7$) exists today in the species of *Univerviae* (ÇELEBIOĞLU & FAVARGER, 1982).

ACKNOWLEDGEMENTS

We are deeply indebted to Prof. C. Favarger, not only because he has put at our disposal his herbarium material of *M. rimarum* but also for his kind help and constructive criticism throughout the course of this study.

REFERENCES

- ÇELEBIOĞLU, T. & C. FAVARGER (1982). Contribution à la cytotaxonomie du genre *Minuartia* L. (Caryophyllacées) en Turquie et dans quelques régions voisines. *Biol. Ecol. Med.* 9(2-3): 139-160.
- ÇELEBIOĞLU, T., K. FAVARGER & K.-L. HUYNH (1983). Contribution à la micromorphologie de la testa des graines du genre *Minuartia* (Caryophyllaceae). I. Sect. *Minuartia*. *Bull. Mus. Natl. Hist. Nat.*, 4^{me} sér., 5, section B, *Adansonia* 4: 415-435.
- FAVARGER, C. (1962a). Contribution à l'étude cytologique des genres *Minuartia* et *Arenaria*. *Bull. Soc. Neuch. Sci. Nat.*, Ser. 3, 85: 53-81.
- FAVARGER, C. (1964). Cytotaxonomie et endémisme. *Compt. Rend. Soc. Biogéogr.* 40: 23-44.
- FAVARGER, C. (1967). Nombres chromosomiques de quelques taxa principalement balkaniques du genre *Minuartia* (L.) Hiern. *Bot. Jahrb.* 86: 280-292.
- HALLIDAY, G. (1964). *Minuartia* L. In: TUTIN, T. G. & al. (ed.), *Flora europaea* 1: 125-132. Cambridge.
- LEVAN, A., K. FEDGA & A. SANDBERG (1965). Nomenclature for centrometric position on chromosomes. *Hereditas* 52: 201-220.
- MATTFELD, J. (1922). Geographisch, genetische Untersuchungen über die Gattung *Minuartia* (L.) Hiern. *Rep. Spec. Nov. Regni Veg.* 100: 146-169.
- MATTFELD, J. (1929). *Minuartia* (L.) Hiern (Karte 51-61). *Pflanzenareale*. 2 Reihe, Heft 6. Verlag von Gustav Fischer (Ed.), Jena.
- MCNEILL, J. (1962). Taxonomic studies in the Alsinoideae: I. Genetic and infragenetic groups. *Notes Roy. Bot. Gard. Edinburgh* 24(2): 79-155.
- MCNEILL, J. (1967). *Minuartia* L. In: P. H. DAVIS (ed.), *Flora of Turkey* 2: 38-67. Edinburgh.
- ÖSTERGREN, G. & W. K. HENEEN (1962). A squash technique for chromosome morphological studies. *Hereditas* 48: 332-341.