

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	41 (1986)
<b>Heft:</b>	2
<b>Artikel:</b>	A new species of Ecdysanthera Hook. & Arn. (Apocynaceae) from Assam
<b>Autor:</b>	Srivastava, S.K. / Mehrotra, B.N.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-880005">https://doi.org/10.5169/seals-880005</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# A new species of *Ecdysanthera* Hook. & Arn. (Apocynaceae) from Assam

S. K. SRIVASTAVA  
&  
B. N. MEHROTRA

## RÉSUMÉ

SRIVASTAVA, S. K. & B. N. MEHROTRA (1986). *Ecdysanthera* Hook. & Arn. (Apocynaceae), espèce nouvelle d'Assam. *Candollea* 41: 381-383. En anglais, résumé français.

Une nouvelle espèce d'Assam (Indes), est décrite et illustrée: *Ecdysanthera lakhimpurensis* Srivastava & Mehrotra spec. nov. (Apocynaceae).

## ABSTRACT

SRIVASTAVA, S. K. & B. N. MEHROTRA (1986). A new species of *Ecdysanthera* Hook. & Arn. (Apocynaceae) from Assam. *Candollea* 41: 381-383. In English, French abstract.

A new species *Ecdysanthera lakhimpurensis* Srivastava & Mehrotra spec. nov. (Apocynaceae) from Assam, India, is described with illustrations.

During the botanical exploration of the forest areas of North Lakhimpur district in Assam, one of the authors (SKS) collected a few specimens later identified as *Ecdysanthera* Hook. & Arn. This genus is represented by 8-10 species distributed in Malayan region and China; only one species viz. *E. micrantha* A. DC. is known so far from Assam, Khasia and Cachar hills in India. Thorough study of the literature and herbaria of Botanical Survey of India, Shillong (Assam), National Botanical Research Institute, Lucknow (LWG) and also of CDRI, reveals that this taxon represents a hitherto unrecognized species.

### ***Ecdysanthera lakhimpurensis* Srivastava & Mehrotra spec. nov. (Fig. 1).**

**Typus:** India: Assam, North Lakhimpur, Dulung R. F., ± 350 m, 24.4.1986, S. K. Srivastava 16105 (holo: CDRI; iso: CDRI).

*Ecdysanthera lakhimpurensis* Srivastava & Mehrotra *E. micranthae* affinis. Sed differt foliis parvioribus, ovatis vel ovato-ellipticis, cum glandibus in axillis nervorum lateralium, floribus corollisque permajoribus, pistillisque bis longioribus.

*Ecdysanthera lakhimpurensis* Srivastava & Mehrotra is allied to *E. micrantha* A. DC. but differs in having smaller leaves, ovate or ovate-elliptic, glands on the axil of the lateral nerves, flowers and corolla much larger, stamens sagittate, flattened, much longer, pistil twice long.

Large shrub, branches terete, young one flattened, channelled, glabrous. Leaves simple, opposite, estipulate, (sheathing cover on each node) 4.5-6.5 × 1.6-2.3 cm, ovate or ovate-elliptic, acute or acuminate, base cuneate, entire, glabrous, membranous or somewhat chartaceous, midrib distinct on both the surfaces, lateral nerves 3-5 pairs, nearly inconspicuous; petioles 1-1.2 cm long, glabrous.

Cymes axillary or terminal, minutely pubescent, trichotomous, 4-7 × 3-4 cm, bracts 1 mm long, ovate, acute, glabrous. Flowers 5 × 2 mm, pedunculate, hermaphrodite, peduncle 1.5-2.8 cm long, pedicels 2 mm long, minutely pubescent, bracteoles tringular, 1-1.5 mm long. Calyx




Fig. 1. — *Ecdysanthera lakhimpurensis* Srivastava & Mehrotra.  
1, habit; 2, flower; 3, bract; 4, corolla (cut open); 5, stamen; 6, pistil.

5-lobed, sepals 1.5 mm long, ovate, acute, glabrous, jointed each other at half of the length. Corolla membranous 5-lobed, jointed each other at  $1/3$  of the length from the apex somewhat rounded. Stamens 5, epipetalous, nearly 2 mm long, attached nearly at the base of the petal lobe, filament short, anther sagittate, spur long. Pistil 2-2.5 mm long, ovary superior, rounded, nearly 1.5 mm in diam., minutely pubescent at the top, style nearly 1 mm long, stigma simple conical. Fruit not seen.

*Flowering.* — March-April.

*Ecology.* — Plant is found growing on the slopes along the dry rocky rivers in moist climatic condition.

#### ACKNOWLEDGEMENTS

Authors are thankful to the Director, Central Drug Research Institute, Lucknow for providing facilities. Thanks are also due to Dr. N. C. Majumdar, Regional Botanist, Botanical Survey of India, Howrah, for latin translation.

Central Drug Research Institute Communication No. 3931.

