

Zeitschrift: Candollea : journal international de botanique systématique =
international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 42 (1987)

Heft: 2

Artikel: A new species of Heliotropium L. (Boraginaceae) from Pakistan

Autor: Nasir, Yasin J.

DOI: <https://doi.org/10.5169/seals-879963>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A new species of *Heliotropium* L. (Boraginaceae) from Pakistan

YASIN J. NASIR

RÉSUMÉ

NASIR, Y. J. (1987). Une espèce nouvelle du genre *Heliotropium* L. (Boraginaceae) du Pakistan. *Candollea* 42: 553-554. En anglais, résumé français.

Description et discussion des affinités d'une espèce nouvelle du Pakistan: *Heliotropium alii* Y. J. Nasir.

ABSTRACT

NASIR, Y. J. (1987). A new species of *Heliotropium* L. (Boraginaceae) from Pakistan. *Candollea* 42: 553-554. In English, French abstract.

Description and discussion of the affinities of a new species from Pakistan: *Heliotropium alii* Y. J. Nasir.

While revising the heliotrophs for the flora of Pakistan, one gathering, namely *S. I. Ali*, *S. A. Faruqi & Sultanul Abedin* II01 from S. Baluchistan turned out to be distinct from other related species. It is described here as *H. alii* Y. J. Nasir. Its affinities are briefly discussed.

***Heliotropium alii* Y. J. Nasir, spec. nov. (Fig. 1A-C).**

Herba annua usque 20 cm alta. Radix palaris gracilis, usque 14 cm longa. Caules e basi plures, ascendentis, pubescentes, pilis brevioribus adpressis longioribus (usque 0.8 mm longis) subpatentibus commixtis. Folia basalia atique caulina petiolata, suprema autem subsessilia. Petiole usque 2 cm longi. Lamina 13-40 × 5-11 mm, elliptico-lanceolata vel oblanceolata, cuneata; pili illis in caule similes, sed pilis validioribus e basi tuberculata exortis. Inflorescentia laxa. Flores uniseriales, ebracteati. Calyx 2 mm longus, lobis obtusis. Corolla 3 mm longa, intus glabra, alba, lobis 1-1.3 mm longis lanceolatus basi latioribus inflexis, sinibus squamigeris, squamis bilobatis.

Holotype: Baluchistan: S. Makran, 8 miles from Sunstar, Turbat road, *S. I. Ali*, *S. A. Faruqi & Sultanul Abedin* II01 (KUH).

As yet known only from the type locality. The type sheet is a gathering of 5 specimens and several twigs. The new taxon is named after Dr. S. I. Ali, Professor of Botany, Karachi University, Karachi.

In the nature of the internally glabrous corolla with lanceolate inflexed lobes, *H. alii* bears resemblance to *H. dasycarpum* Bunge, but differs in the annual habit, longer petioles and obtuse calyx lobes (not acute). The inflorescence is like *H. remotiflorum* Rech. f. & Riedl (sect. *Adelotropium* Rech. f. & Riedl), which however has a corolla pubescent internally. In habit and leaves *H. alii* resembles the annual *H. europeum* L. var. *lasiocarpum* (Fisch. & Mey.) Kazmi and its ally *H. dolosum* De Not., but differs from both in the lanceolate inflexed corolla lobes (not obtuse) and obtuse calyx lobes. There is also some resemblance of this taxon with two extra-Pakistan species, namely *H. mamamense* Bunge and *H. mesinanum* Bunge (sect. *Pseudobucanion* Riedl); from the




Fig. 1. — *Heliotropium alii* Y. J. Nasir, spec. nov.
A, flower, $\times 20$; B, carpel showing hairy stigma and short style, $\times 40$; C, corolla opened, $\times 20$ (drawn by Shahid Hamid).

former, *H. alii* differs in being pubescent (not densely so), obtuse calyx lobes, smaller corolla and obtuse anthers (not apiculate); from the latter, the species differs in the more or less ascending habit, obtuse calyx lobes and corolla limb which is only slightly shorter than tube.

The flowering period is February.

ACKNOWLEDGEMENTS

I wish to acknowledge my thanks to the Curator, Karachi University Herbarium, Karachi, for the general loan of *Heliotropium* material. My thanks are also due to Mr. A. Radcliffe-Smith, Royal Botanic Gardens, Kew, for the latin diagnoses to the new species. The figures were rendered by our herbarium artist, Mr. Shahid Hamid, for which I am grateful.