

Zeitschrift: Candollea : journal international de botanique systématique =
international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 44 (1989)

Heft: 1

Artikel: Review of the genus Litosanthes Bl. (Rubiaceae)

Autor: Deb, D.B. / Gangopadhyay, Mohan

DOI: <https://doi.org/10.5169/seals-879610>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Review of the genus *Litosanthes* Bl. (Rubiaceae)

D. B. DEB
&
MOHAN GANGOPADHYAY

RÉSUMÉ

DEB, D. B. & M. GANGOPADHYAY (1989). Révision du genre *Litosanthes* Bl. (Rubiacee). *Candollea* 44: 209-223. En anglais, résumés français et anglais.

Révision du genre *Litosanthes* Bl. (Rubiaceae). Le statut taxonomique du genre est réévalué. Quinze espèces sont transférées de *Lasianthus* Sect. *Pedunculatae* Hook. f. dans *Litosanthes* Bl. Des combinaisons nouvelles sont faites: 1. *Litosanthes biermannii* (King ex Hook. f.) Deb & Gang.; 2. *L. capitulatus* (Wt.) Deb & Gang.; 3. *L. cinereus* (Gamble) Deb & Gang.; 4. *L. dichotomus* (Wt.) Deb & Gang.; 5. *L. longicauda* (Hook. f.) Deb & Gang.; 6. *L. micranthus* (Hook. f.) Deb & Gang.; 7. *L. venulosus* (Wt. & Arn.) Deb & Gang.; 8. *L. filiformis* (King & Gamble) Deb & Gang.; 9. *L. filiformis* var. *bracteata* (King & Gamble) Deb & Gang.; 10. *L. pendulus* (Ridley) Deb & Gang.; 11. *L. pseudo-lucidus* (King) Deb & Gang.; 12. *L. robinsonii* (Ridley) Deb & Gang.; 13. *L. scalariformis* (King & Gamble) Deb & Gang.; 14. *L. subspicatus* (King & Gamble) Deb & Gang.; 15. *L. varians* (Hook. f.) Deb & Gang.; 16. *L. villosus* (Ridley) Deb & Gang. Mots clé: taxonomie de la plante, Rubiacées, *Litosanthes*, description de la clé, nouvelles combinaisons.

ABSTRACT

DEB, D. B. & M. GANGOPADHYAY (1989). Review of the genus *Litosanthes* Bl. (Rubiaceae). *Candollea* 44: 209-223. In English, French and English abstracts.

The paper presents a review of the genus *Litosanthes* Bl. (Rubiaceae). The taxonomic status of the genus is reviewed and reinstated. 15 species are transferred from *Lasianthus* Sect. *Pedunculatae* Hook. f. to *Litosanthes* Bl. New combinations made are: 1. *Litosanthes biermannii* (King ex Hook. f.) Deb & Gang.; 2. *L. capitulatus* (Wt.) Deb & Gang.; 3. *L. cinereus* (Gamble) Deb & Gang.; 4. *L. dichotomus* (Wt.) Deb & Gang.; 5. *L. longicauda* (Hook. f.) Deb & Gang.; 6. *L. micranthus* (Hook. f.) Deb & Gang.; 7. *L. venulosus* (Wt. & Arn.) Deb & Gang.; 8. *L. filiformis* (King & Gamble) Deb & Gang.; 9. *L. filiformis* var. *bracteata* (King & Gamble) Deb & Gang.; 10. *L. pendulus* (Ridley) Deb & Gang.; 11. *L. pseudo-lucidus* (King) Deb & Gang.; 12. *L. robinsonii* (Ridley) Deb & Gang.; 13. *L. scalariformis* (King & Gamble) Deb & Gang.; 14. *L. subspicatus* (King & Gamble) Deb & Gang.; 15. *L. varians* (Hook. f.) Deb & Gang.; 16. *L. villosus* (Ridley) Deb & Gang. Key words: plant taxonomy, Rubiaceae, *Litosanthes*, key descriptions, new combinations.

The Indian species are described with key, synonymy, citations, types, distribution, flowering and fruiting time, etc. Malayan and Sri Lankan and others species are not sufficiently available for thorough study of those species.

The genus *Litosanthes* Bl. Bijdr. 994. 1826 was postulated as a monotypic genus from Java. Subsequently 4 species were described from Formosa and New Guinea. DE CANDOLLE (1830), MIQUEL (1855), HOOKER (1873), SCHUMANN (1891) and others recognized the genus. HOOKER (1880), however, reduced it under *Lasianthus* Sect. *Pedunculatae* Hook. f. RIDLEY (1923) followed it. BAKHUIZEN (1965) reestablished the generic status mainly for imbricate aestivation of corolla lobes, divided stipules and pedunculate inflorescence, as mentioned in his key to these two genera, even though in the description of the genus he erroneously mentioned the aestivation as valvate like all earlier workers, even by BLUME himself (1826) in course of describing the monotypic genus.

In course of study of the Indian *Lasianthus* the authors of the present paper observed the presence of these three characters in the species of *Lasianthus* Sect. *Pedunculatae*, distributed in India, Sri Lanka, Java and Malaya as available in herb. CAL. It is now generally accepted that there is only one kind of aestivation of corolla in a genus in Rubiaceae. On the other hand, the combination of these three characteristics is sufficient to exclude such species from the genus *Lasianthus* and place them under the most allied genus *Litosanthes* as rightly distinguished and reinstated by Bakhuizen (l.c.).

The present paper describes the genus *Litosanthes* Bl. in India with key to and descriptions of the species with nomenclatural changes, synonymy, types, citations, distributions, flowering and fruiting time, etc. Nomenclatural changes in the names of Sri Lankan and Malayan species are made but they are not described as some of their types and other specimens were not available for thorough study.

Taxonomic position of the genus

DE CANDOLLE (1830) treated the genus in subtribe *Guettardeae* of the tribe *Guettardaceae* Kunth. Hook. f. (1873) and SCHUMANN (1891) placed it under the tribe *Psychotrieae*. RIDLEY (1923) reduced the genus to the section *Litosanthes* under the genus *Lasianthus*. BAKHUIZEN f. (1965) resurrected it to the generic status for reasons already stated. The genus is placed under the tribe *Psychotrieae* sensu stricto.

Range of morphological characters

The plants vary from under shrubs to small trees up to 6 m in height. Branchlets are glabrous in *L. dichotomus*, glabrous or puberulous in *L. longicauda*, pubescent in *L. cinereus*, ferruginous adpressed pubescent in *L. capitulatus* or strigose in *L. biermannii* and *L. micranthus*. Leaves are very variable in form in most of the species. They are glabrous on upper surface in all the Indian species and some of the Malayan species, but variously hairy on the undersurface; glabrous or sparsely pubescent in *L. venulosus*, pubescent in *L. capitulatus*, *L. cinereus*, puberulous in *L. longicauda*, strigose in *L. biermannii*, pubescent on midrib and nerves in *L. dichotomus*.

Inflorescence is much branched and largest (15-30 × 20-30 mm) in *L. biermannii*, smallest in *L. dichotomus*. Bracteoles are sessile or subsessile in *L. capitulatus*, *L. cinereus* and *L. venulosus*, stalked in *L. dichotomus*. Calyx is toothed or lobed; teeth linear-lanceolate in *L. micranthus*; lobes ovate in *L. capitulatus*, ovate-oblong in *L. biermannii*, ovate-lanceolate or narrowly oblong in *L. dichotomus*, linear-lanceolate in *L. cinereus*, linear in *L. venulosus*. Corolla lobes are 4 in most of the species, 5 in *L. micranthus* and 4-6 in *L. biermannii*, lobes are glabrous above, tomentose beneath in *L. longicauda*, puberulous on both sides in *L. venulosus*, strigose above and villous beneath in *L. biermannii*. Filaments are generally adnate below the midtube but above midtube in *L. biermannii*. Style is longest (12 mm) in *L. capitulatus*, glabrous or puberulous in *L. venulosus*; 4 stigma lobes are common, 5 in *L. biermannii* and *L. micranthus*, 2 or 3 in *L. cinereus*. Pyrenes are generally 4 but in *L. biermannii* and *L. micranthus* 5.

Litosanthes Bl., Bijdr. 994. 1826 & in Flora 8: 189. 1825 (*Lithosanthes*); DC., Prodr. 4: 465. 1830; Miq., Fl. Ind. Bat. 2: 314. 1855; Hook. f. in Benth. & Hook. f., Gen. Pl. 2: 131. 1873; Schumann in Engl. & Prantle, Nat. Pflanzenfam. IV. Rubiac. 121. 1891; Bakh. f. in Backer & Bakh. f., Fl. Java 2: 243. 1965.

Syn.: *Lithosanthes* A. Rich., Mem. Soc. Nat. Hist. Paris 5: 213. 1829. *Santia* Wt. & Arn., Prodr. 422. 1834. *Lasianthus* Sect. *Petiolatae* Hook. f., Fl. Brit. Ind. 3: 190. 1880. *Lasianthus* Sect. *Litosanthes* Ridley, Fl. Mal. Penins. 2: 151. 1923.

Undershrubs, shrubs or small trees, branching di or trichotomous, quadrangular when young, terete in age, glabrous, pubescent or strigose. Leaves opposite, decussate, petiolate, elliptic, elliptic-lanceolate, elliptic-ovate, ovate, oblong, orbicular or obovate, cuspidate or caudate-acuminate at apex, acute or subacute at base, glabrous above, strigose beneath; lateral nerves sub opposite, arcuate, forked, reticulate, inconspicuous above; stipules interpetiolar, more or less lobed or divided,

ciliate, pilose or setose, sometimes glabrous and fleshy. Inflorescence pedunculate, axillary cymes, bracteate and bracteolate, slender or stout, pubescent, puberulous or strigose, branching di or trichotomous or whorled. Flowers sessile or subsessile, bisexual, 4 merous; bracts minute, sometimes prominent. Hypanthium ovoid or obovoid. Calyx tube very short, oblong; teeth or lobes 4, short or long. Corolla urceolate, glabrous above, villous within, especially in throat; lobes 4-6, short, imbricate in bud. Stamens 4-6, adnate in throat; filaments minute; anthers dorsifixed, linear-oblong or oblong, included or subexserted. Disk fleshy, glabrous. Ovary 4-5 loculed; ovule one in each locule, basal, erect, anatropous; stigma 2-5 lobed, subclavate. Fruit a drupe, ovoid, obovoid or globose, crowned with persistent calyx lobes; pyrenes 4-5, ridged and furrowed, thick walled, crustaceous; with or without raphides, 1 seeded. Embryo albuminous, erect, incurved.

Type: *L. biflora* Bl.

Distribution. — About 20 species in India, Sri Lanka, Java, Malaya, Formosa and New Guinea.

Key to the Indian species

1. Branchlets glabrous in age; leaves glabrous or sparsely pubescent beneath 2
- 1a. Branchlets strigose or pubescent; leaves strigose or pubescent beneath 5
2. Fruits globose 4
- 2a. Fruits ovoid 3
3. Fruit glabrous; lateral nerves 6-9 pairs 7. *L. venulosus*
- 3a. Fruit puberulous; lateral nerves 7 pairs 4. *L. dichotomus*
4. Lateral nerves 3-4 pairs 3. *L. cinereus*
- 4a. Lateral nerves 6-10 pairs 5. *L. longicauda*
5. Leaves strigose beneath; pyrenes 4 6
- 5a. Leaves pubescent beneath; lateral nerves 7-10 pairs 2. *L. capitulatus*
6. Pyrenes 5; stigma 5 lobed 1. *L. biermannii*
- 6a. Pyrenes 5; stigma 4 lobed 6. *L. micranthus*

1. *Litosanthes biermannii* (King ex Hook. f.) Deb & Gang., comb. nov.

Syn.: *Lasianthus biermannii* King ex Hook. f., Fl. Brit. Ind. 3: 190. 1880 (Type: Yoksum, 1857, *T. Thomson s.n.* (K! selected as the lecto); Khasia, *J. D. Hooker & T. Thomson* Maphitidia (K!, CAL!); *Hooker & Thomson* 16 (E); Maphlong, *Griffith s.n.* E.I.C. 2931 & 2932 (K! CAL!); Brandis, Ind. Trees 398. 1906; Kanjilal & al., Fl. Assam 2: 93. 1939; Balakr., Fl. Jowai 1: 42. 1981.

Shrubs 3-6 m high; branchlets di or trichotomous, quadrangular when young, terete in age, strigose. Leaves petiolate, 6-22 × 2-5 cm, elliptic, elliptic-ovate or elliptic-lanceolate, caudate or cuspidate acuminate at apex, acute or subacute at base, greyish to blackish green or pale when dry, membranous, glabrous above, strigose beneath on midrib, nerves and sometimes nervules; midrib channelled, slender above; lateral nerves 5-7 pairs, subopposite, arcuate, inconspicuous above; nervules sub parallel, forked, reticulate; petioles 7-14 mm long, slender, strigose; stipules 3-5 × 2-2 mm, 3-5 lobed, triangular, linear-lanceolate or ovate-lanceolate, thick, dentate, mid lobe larger, acuminate, setose, strigose, with colleters at base inside. Inflorescence axillary cymes, 1.5-3 × 2-3 cm; peduncles 5-15 mm long, slender, strigose; branchlets short, di or trichotomous or whorled, sometimes strigose; bracts 6-10 mm long, linear, strigose; bracteoles 2-5 mm long, linear-lanceolate or triangular, strigose. Flowers congested, 8-9 mm long, sessile or subsessile; pedicels 0-0.5 mm long, strigose. Hypanthium 0.5-1 mm long, ovoid, strigose; calyx tube ± 0.5 × 1-1.5 mm, sparsely




Fig. 1. — *Litosanthes biermannii* (King ex Hook. f.) Deb & Gang.
A, habit; B, flower; C, flower split opened; D, fruit; E, seed; F, embryo.

strigose; lobes 4-6, white, ovate-oblong, acute at apex, strigose. Corolla tube 3-4 mm long, glabrous above, villous within; lobes 4-6, 2.5-3 mm long, oblong-ovate, strigose above, villous beneath. Stamens 4-6, inserted; filaments minute, adnate below throat; anthers 1-1.5 mm long, oblong, dorsally convex. Ovary \pm 0.5 mm long, ovoid, 5-loculed; style 3-4 mm long, slender, glabrous; stigma 5-lobed, \pm 0.3 mm long, linear, papillose; disk 0.5-1 mm across, flattened, smooth. Fruits sessile or subsessile, blue, 1-1.5 \times 1-1.2 cm, globose, crowned with persistent calyx lobes, strigose; pyrenes 5, acute at both ends, solitary ridged and grooved at back, thick walled, warty, with raphides, mesocarp pulpy. Seeds \pm 5 mm long, broadly stalked, mucronate at apex, dorsally convex; embryo \pm 3 mm long, radicle 2.5 mm long, obtuse at base; cotyledons ovate, acute at apex, thin.

Flowers and fruits. — Throughout the year.

Distribution. — India: Sikkim to Arunachal Pradesh, Assam, Meghalaya, Nagaland, Manipur and Mizoram, at 750-2500 m in altitude; Burma.

Exsiccata studied

Sikkim: Sinebur, *T. Anderson* 158 (CAL); Rinchingspong, *T. Anderson* 690, 935 (CAL); Sikkim, *G. King* 233, 690, 796 (CAL); Yoksam-Bakkim, *H. Hara & al.* 1241 (CAL); Pemayangtse, *J. D. A. Stainton* 5339 (BM); sureil, *G. H. Cave s.n.* (E); West Bengal; Darjeeling Dt., Rungbee, *C. B. Clarke* 8578, 13121B; *G. King s.n.* (CAL) & *Prain's collector s.n.* (BSI); Surumtong, *C. B. Clarke* 12636B (CAL); Darjeeling, *J. S. Gamble* 3789, 8356, 9810 (CAL). Tatkah, *J. H. Lace* 2203 (CAL, DD, E); Mawrim forest, *J. H. Lace* 2456 (E, CAL); Rambhinghora, *W. W. Smith* 331 (CAL); Sinchal, *S. Kurz s.n.* (CAL); Tangloo, *G. King s.n.* (CAL), Labha, *H. Santapau & S. K. Mukerjee* 123 (CAL); Dhobijhora, *J. S. Gamble* 10315 (CAL); Rungbee, *C. B. Clarke* 8658 (K); Burch Hill, *W. A. Talbot* 3365 (BSI). Arunachal Pradesh: Daphla Hills, Shengorh. *J. L. Lister* 295 & 298 (CAL) & *I. H. Burkhill* 36549 (CAL); Upper Renjing, *I. H. Burkhill* 36284 (CAL); Sissini, *G. Panigrahi* 6283 (CAL); Parila-Sissini, *G. Panigrahi* 7008 (CAL); Tirap Dt., Wakka, *D. B. Deb* 26481 (CAL). Subansiri Dt., Apa Tani valley, *Cox & Hutchinson* 388 (E). Meghalaya, *King's collector s.n.* (CAL) & *G. Mann s.n.* (CAL); Momluh, *N. L. Bor* 184 (DD); Khasia, *Griffith s.n.* (Kew Distrib. No. 2931, 2932) (CAL, K); Surareem, *C. B. Clarke* 45190 (CAL); Khasia, *B. Khan s.n.* (CAL); Oldham 8 (CAL); Jarain, *C. B. Clarke* 18319 (CAL); Khasia, *J. D. Hooker & T. Thomson* 16 Maphitidia (CAL); *G. A. Gammie* 416 (CAL); *G. A. Gammie s.n.* (CAL); Manipur: Japvo, *G. Watt* 6238 (BSIS, CAL); Sirohifur, *G. Watt* 6359 (BSIS, CAL, E). Nagaland: Kohima, *C. B. Clarke* 40968 (CAL). Mizoram: Aijal, *D. B. Deb* 30654 (CAL). Burma: Khaiyana, *F. Kingdon-ward* 17424 (BM)

2. *Litosanthes capitulatus* (Wt.) Deb & Gang., comb. nov.

Syn.: *Lasianthus capitulatus* Wt. in Calc. Journ. Nat. Hist. Soc. 6: 511. 1846 (Type: Nilgiri, Sispara, *Wight s.n.* (K!)); Hook. f., Fl. Brit. Ind. 3: 191. 1880; Brandis, Ind. Trees 398. 1906; Gamble, Fl. Pres. Madras 648. 1921.

Maphitidia capitula Walp., Ann. Bot. Syst. 2: 762. 1852.

Shrubs; branches dichotomous; branchlets slender, adpressed ferruginous pubescent. Leaves petiolate, 8-15 \times 2.5-6 cm, elliptic, ovate, oblong or rarely obovate, short acuminate at apex, acute or obtuse at base, slightly incurved at margin, glossy, glabrous above, pubescent beneath, brown or pale green when dry; midrib inconspicuous, shallow channelled above, terete beneath; lateral nerves 7-10 pairs, opposite or sub-opposite, subparallel, curved, ascending, conspicuous on both surfaces; nervules parallel, forked, conspicuous; petioles 3-12 mm long, stout, pubescent; stipules persistent, 1-3 \times 1-2 mm, triangular or ovate, narrowly or irregularly toothed or divided, setose, acute at apex, pubescent. Inflorescence axillary, peduncled cymes, 8-13 \times 10-20 mm, 3-4 flowered, bracteate, bracteoles 1-2 mm long, linear, puberulous. Flowers sessile or pedicellate, 12-15 mm long; pedicels 0-1 mm long, puberulous. Hypothecium \pm 1 mm long, ovoid, puberulous above; calyx tube \pm 0.5 \times 1.5-2.5 mm, puberulous above; lobes 4, 1.5-2 mm long, ovate, acute, glabrous, anastomosing conspicuously. Corolla tube \pm 10 mm long, puberulous above, pubescent at upper part beneath; lobes \pm 5 mm long, ovate-oblong, puberulous above, pubescent beneath. Stamens 4, included; filaments \pm 0.5 mm long, thin, adnate at mid tube; anthers \pm 1 mm long, oblong. Ovary \pm 0.5 mm long, ovoid, 4-loculed; style \pm 15 mm long, narrow, glabrous; stigma 4-lobed,




Fig. 2. — *Litosanthes capitulatus* (Wt.) Deb & Gang.
A, habit; B, bracteole; C, flower; D, flower split opened; E, fruit; F, seed; G, embryo.

horizontally spreading, \pm 0.5 mm long, ovate, papillose; disk 1-1.5 mm across, subglobose, smooth. Fruits sessile, \pm 6 \times 5 mm, ovoid, crowned with persistent calyx lobes, 4-shallow channelled; pericarp thin, warty; mesocarp thin; pyrenes 4, dorsally convex, ventrally angled, shallow channelled at back, ellipsoid, acute at both ends, without raphides. Seeds \pm 3 mm long, thin, flattened, funicle short, embryo 0.5 mm long; radicle minute, ovate, acute at apex.

Flowers: May-September. *Fruits:* November-December.

Distribution. — India: Tamilnadu and Kerala in evergreen forests upto 1800 m in alt.

Exsiccata studied

Tamilnadu: Nilgiri Dt., Sisparaghat, *R. H. Beddome* 4051, 4032 (BM) & *J. S. Gamble* 13374 (CAL) & 14273 (BM). Kerala: Palghat hills, *R. H. Beddome* 4035 (BM); Silent valley, *R. H. Beddome* 4023 (BM); Brahmagiri shola, *C. E. C. Fischer* 252 (CAL).

3. *Litosanthes cinereus* (Gamble) Deb & Gang., comb. nov.

Syn: *Lasianthus cinereus* Gamble in Kew Bull. 1920: 249 & Fl. Pres. Madras 648. 1921 (Type: Tamilnadu, Kannikatti to Kalivayalpil, 31 May 1901, *C. A. Barber* 3014 (K!, CAL)); Natirieal near Travancore, March 4, 1917, *K. Rangachari* 1450 (MH!).

Undershrubs or shrubs, 2-4 m high; branchlets slender, compressed, pubescent when young, terete, glabrous in age. Leaves petiolate, 7-10 \times 2-3.5 cm, obovate, elliptic or ovate, long acuminate at apex, acute at base, slightly incurved at margin, thin coriaceous, glabrous above, sparsely pubescent beneath on midrib and nerves; midrib slender, channelled above; lateral nerves 3-4 pairs, subopposite, arcuate; nervules parallel, conspicuous; petioles 5-7 mm long, slender, dense or sparsely pubescent; stipules 2-3 \times 1.5-5 mm, triangular, oblong or ovate-lanceolate, irregularly toothed or lobed, setose, acuminate, pilose inside and at margin, few colleters at base beneath. Inflorescence peduncled cymes, 1.5-2.5 \times 1.5-2 cm, di or trichotomously branching, pubescent; bracts and bracteoles 2-4 mm long, linear-lanceolate, pubescent. Flowers sessile or shortly pedicellate, 7-9 mm long, white, sometimes pinkish; pedicels 0.5-1 mm long, pubescent. Calyx pubescent; hypanthium 1-1.5 mm long, ovoid, pubescent; calyx tube \pm 0.5 \times 1-1.5 mm; lobes 4, 4-5 mm long, linear-lanceolate, pubescent above. Corolla 4-5 mm long, glabrous above, pubescent beneath at throat; lobes 4, 1.5-2 mm long, ovate, pubescent at tip, glabrous beneath. Stamens 4, inserted; filaments 0.5 mm, adnate at throat; anthers \pm 0.5 mm long, oblong. Ovary \pm 0.5 mm long, ovoid, 4-loculed; style 3-3.5 mm long, thin, glabrous; stigma 2 or 3 lobed, \pm 0.5 mm long, ovate-oblong, glabrous; disk 0.5-1 mm across, subglobose, smooth. Fruits shortly stalked, 5-6 \times 5-6 mm, globose, 4-furrowed, glabrous, crowned with persistent calyx lobes; stalks 1-2 mm long, pubescent or glabrous; pyrenes 4, ovoid, obtuse at apex, acuminate at base, dorsally convex, ventrally angled, thick walled, with raphides. Seeds 3-4 mm long, thin, flat, obtuse at apex, shortly stalked at base; embryo \pm 1.5 mm long; radicle \pm 1 mm long, obtuse at base, thin, flattened; cotyledons \pm 0.5 mm long, ovate, obtuse at apex, thin.

Flowers: March-May. *Fruits:* September-January.

Distribution. — India: Tamilnadu and Kerala at 850-1600 m in alt.

Exsiccata studied

Tamilnadu: Tirunelveli Dt., Agasthymalai, *A. N. Henry* 16293 (CAL); Naterikal, *K. C. Jacob* 13365 (MH); Sengelteri, *K. M. Mathew* 16285 (BM); Kanniyam Kumari Dt., Mathukuzhivayal, *A. N. Henry* 48627 (CAL); Vanamatti, *A. N. Henry* 70390 (CAL); Panagudi, *B. D. Sharma* 40028 (MH). Kerala: Trivandrum Dt., Bonaccord Estate, 800 m, *M. Mohanan* 59447 (CAL) & *J. Joseph* 44506 (MH); Adhimalai, *M. Mohanan* 666010 (MH).

4. *Litosanthes dichotomus* (Wt.) Deb & Gang., comb. nov.

Syn: *Lasianthus dichotomus* Wt. in Calc. Journ. Nat. Hist. Soc. 6: 508. 1845 (Type: Shevagherry hills, April 1836, *Wight* 1401 (K!) — selected as lecto; Herb. *Wight* s.n. (K!-para));




Fig. 3. — *Litosanthes cinereus* (Gamble) Deb & Gang.
A, habit; B, bracteole; C, flower; D, flower split opened; E, fruit; F, seed; G, embryo.




Fig. 4. — *Litosanthes dichotomus* (Wt.) Deb & Gang.
A, habit; B, flower; C, flower split opened; D, fruit; E, seed; F, embryo.

Bedd. Ic. Pl. Ind. Or. 1: t. 13. 1874; Hook. f., Fl. Brit. Ind. 3: 191. 1880; Brandis, Ind. Trees 398. 1906; Gamble, Fl. Pres. Madras 645. 1921.
Mephitidia dichotoma Walp., Ann. Bot. Syst. 2: 761. 1862.

Shrubs about 2 m high, branching dichotomous; branchlets quadrangular, pubescent when young, terete, glabrous in age; cork cells present. Leaves petiolate, 2.5-5.5 × 2-3.5 cm, ovate, ovate-orbicular, shortly acuminate at apex, slightly incurved at margin, cordate at base, thin coriaceous, glabrous above, pubescent on midrib and nerves beneath, pale green when dry, midrib channelled above; lateral nerves 7 pairs, subopposite, subparallel, arched; nervules slender, close, parallel; petioles ± 5 mm long, channelled, glabrous above, terete, pubescent beneath; stipules 5-6 × 2-3 mm, ovate-lanceolate, lobed or dentate, acuminate, pubescent above. Inflorescence axillary peduncled cymes, 2-3 flowered; peduncle 3-4 mm long, slender, pubescent; bracteoles 5-6 mm long, linear, pubescent. Flowers pedicellate, 9-10 mm long; pedicels 0.5-1 mm long, pubescent. Hypanthium 0.5-1 mm long, obovoid, glabrous or pubescent; calyx tube ± 0.5 × 1-1.5 mm, pubescent or glabrous above; lobes 4, 1.5 mm long, linear-oblong, or ovate-lanceolate, puberulous or pubescent above. Corolla tube 3-4 mm long, glabrous; lobes 4, ± 2.5 mm long, ovate-oblong, pubescent. Stamens 4, subexserted; filaments minute, adnate below throat; anthers 1-1.5 mm long, narrowly oblong. Ovary 0.5-1 mm long, obovoid, 4 loculed; style ± 3 mm long, thin, glabrous; stigma 4-lobed, lobes narrowly oblong, glabrous; disk ± 0.5 mm across, subglobose, smooth. Fruit stalked, 8-10 × 4-6 mm, ovoid, crowned with persistent calyx lobes, 4-5 shallow channelled, puberulous; stalks 5-10 mm long, puberulous; pyrenes 4, dorsally convex, ventrally angled, thin walled, with raphides. Seeds 3-3.5 mm long, semiterete, obtuse at apex, stalked at base; stalk flattened; embryo minute.

Flowers: November-January. *Fruits:* February.

Distribution. — India: Tamilnadu and Kerala at altitude upto 1550 m.

Exsiccata studied

Tirunelveli Dt., Tirunelveli hills, *R. H. Beddome* 4013 (BM). Kanniakumari Dt., Pangudi, *B. D. Sharma* 40033 (MH). Kerala, South Travancore, 1500 m, *R. H. Beddome* 4020 (BM).

5. *Litosanthes longicauda* (Hook. f.) Deb & Gang., comb. nov.

Syn.: *Lasianthus longicauda* Hook. f., Fl. Brit. Ind. 3: 190. 1880 (Type: Khasia, *J. D. Hooker & T. Thomson* (K!, E!, L!); East Bengal, *Griffith* s.n. E. I. C. 2914 (K-para.)); Brandis, Ind. Trees 398. 1906; Kanjilal & al., Fl. Assam 3: 93. 1939.

Shrubs 3-3.5 m high; branchlets quadrangular when young, terete in age, slender, puberulous or glabrous, blackish brown when dry. Leaves petiolate, 7-18 × 1.5-3 cm, ovate-lanceolate, or oblong-lanceolate, caudate acuminate at apex, acute at base, slightly incurved at margin, thin coriaceous, glabrous above, puberulous beneath, brown, pale green or black when dry; midrib slender, channelled above; lateral nerves 6-10 pairs, opposite or subopposite, parallel, slender, faint, intramarginal nerves join them; nervules parallel, forked, inconspicuous; petioles 8-13 mm long, slender, glabrous above, glabrous or sparsely puberulous beneath; stipules 1-2 × 1-2 mm, triangular, lobed or toothed, acute or acuminate, pilose, leaving no hairs when fall off. Inflorescence axillary, short peduncled. Flowers sessile, 7-8 mm long, blue purple. Hypanthium ± 1 mm long, obovoid, glabrous or sparsely puberulous; calyx tube 1 × 1.5 mm, cupular, puberulous above; teeth 4, triangular, acute, puberulous. Corolla tube 4-5 mm long; lobes ± 2 mm long, ovate, glabrous above, tomentose beneath. Stamens 4, inserted; filaments minute, adnate inside the tube; anthers 1-1.5 mm long, oblong. Ovary ± 0.5 mm long, obovoid, 4-loculed; style 4-5 mm long, narrow, glabrous; stigma ± 0.5 mm long, 4-lobed, lobes linear, glabrous, disk ± 1 mm across, subglobose, smooth. Fruit ± 4 × 5 mm, globose, short stalked, crowned with persistent calyx teeth, 4-furrowed; pericarp thin, glabrous; mesocarp thin; pyrenes 4, dorsally convex, shallow longitudinally channelled, ventrally angled, ellipsoid, acute at both ends, thick walled. Seeds ± 4 mm long, oblong, acute at apex, short stalked at base, shallow channelled; embryo ± 3 mm long; radicle ± 2.5 mm long, semiterete, obtuse at base; cotyledons ovate, acute at apex, thin.




Fig. 5. — *Litosanthes longicauda* (Hook. f.) Deb & Gang.
A, habit; B, flower; C, flower split opened; D, fruit; E, seed; F, embryo.

Flowers: May-July. *Fruits:* August-November.

Distribution. — India: Arunachal Pradesh, Meghalaya at 1200-2100 m in altitude.

Exsiccata studied

Arunachal Pradesh: Subansiri Dt., Hapoli, *G. Panigrahi* 19813 (CAL). Meghalaya: Khasi, Maflong, *J. D. Hooker & T. Thomson* 1988 & 18 Mephididia (E, CAL); Surareem, *C. B. Clarke* 43890A (CAL); Maflong, *C. B. Clarke* 44237A (BM), 44237B (CAL), 44237G (CAL); Syndoh forest, *U. N. Kanjilal* 2497 (CAL); Rambrai, *C. B. Clarke* 15318 (CAL); Shillong, *D. Prain* s.n. (BSIS, CAL).

6. Litosanthes micranthus (Hook. f.) Deb & Gang., comb. nov.

Syn.: *Lasianthus micranthus* Hook. f., Fl. Brit. Ind. 3: 190. 1880 (Type: Arunachal, Mishmi hills, *Griffith* s.n. EIC 2930 (K!, iso. CAL!); Brandis, Ind. Trees 398. 1906.

Shrubs; branchlets slender, glabrous; very young short strigose. Leaves short petiolate, 7-12 × 3-4 cm, ovate, elliptic-oblong, or obovate, acuminate at apex, rounded or subacute at base, membranous, glabrous above, strigose on midrib, nerves and nervules beneath, pale green when dry; midrib slender, shallow channelled above; lateral nerves 5-6 pairs, subopposite, arcuate, slender, inconspicuous above; nervules slender, parallel; petioles 4-9 mm long, slender, strigose; stipules 1.5-2 × 2-3 mm, triangular, acute, strigose above. Inflorescence short peduncled cymes, 4-5 flowered; peduncle 3-6 mm long, slender, strigose; bracteoles minute, triangular, acute, pubescent or strigose. Flowers sessile, ± 4 mm long, short tubular. Hypanthium ± 1 mm long, obovoid, strigose or puberulous; calyx tube ± 1 × 1.5 mm; teeth 5, linear-lanceolate, puberulous or strigose above. Corolla tube 1.5-2 mm long, glabrous above, villous within above; lobes 5, ± 1.5 mm long, ovate-lanceolate, puberulous or strigose above, glabrous beneath. Stamens 5, included; filaments minute, adnate at mid tube; anthers hardly 1 mm long, oblong. Ovary ± 0.5 mm long, obovoid, 5-loculed; style ± 2 mm long, filiform, glabrous; branches 5, minute; disk ± 0.5 mm across, subglobose, smooth. Fruit 5-6 × 5-6 mm, globose, crowned with persistent calyx lobes, 5-shallow grooved; pericarp thin, glabrous; mesocarp thin; pyrenes 5, dorsally convex, shallow ridged, furrowed, ventrally angled, obtuse at apex and base, thin, hard walled. Seeds ± 2 mm long, curved, terete, obtuse at apex, short stalked at base; embryo ± 1.5 mm long, bent; radicle semi-terete, narrow, obtuse at base; cotyledons ovate, obtuse at apex.

Distribution. — India: Assam, Meghalaya, Arunachal Pradesh.

Exsiccata studied

Assam, *Jenkins* s.n. (CAL); Khasia, *Masters* s.n. (CAL); Khasia, *Masters* s.n. (CAL). Arunachal Pradesh: Baligaon, *Prain's collector* 178 (CAL).

7. Litosanthes venulosus (Wt. & Arn.) Deb & Gang., comb. nov.

Syn.: *Sanitia venulosa* Wt. & Arn., Prodr. 422. 1834 (Type: *Wight* 1353 (E holo)).
Lasianthus venulosus (Wt. & Arn.) Wt. in Calc. Journ. Nat. Hist. Soc. 6: 508. 1845;
 Hook. f., Fl. Brit. Ind. 3: 191. 1880; Wt. Ic. t. 1032. 1845; Bedd., Fl. Sylv. Anal. Gen. t. 17. f. 5. 1869; Cooke, Fl. Pres. Bombay 1: 621. 1903; Brandis, Ind. Trees 398. 1906;
 Fyson, Fl. Nilgiri Pulney Hill Tops 1: 198. 1915; Gamble, Fl. Pres. Madras 648. 1921.
Mephididia venulosa Walp., Ann. Bot. Syst. 2: 761. 1852.

Undershrubs, 1-2 m high; branches and branchlets di or tri-chotomous, glabrous, smooth, blackish chocolate, internode short. Leaves petiolate, 2-12 × 1-5.5 cm, ovate, ovate-oblong, oblong, elliptic or rarely obovate, acute, acuminate or mucronate at apex, rounded or subacute at base, slightly incurved at margin, thin, membranous, glabrous above, glabrous or sparsely pubescent on midrib and nerves beneath, reddish, black or brown when dry; midrib inconspicuous above, stout beneath; lateral nerves 6-9 pairs, opposite or subopposite, subparallel, arcuate, conspicuous; nervules very close, reticulate, conspicuous; petioles 5-10 mm long, stout, glabrous, rarely sparsely pubescent above; stipules 2-4 × 1.5-3 mm, triangular or lanceolate, divided or lobed, acute at apex, pilose at margin and beneath or throughout and few colleters present at base beneath. Inflorescence




Fig. 6. — *Litosanthes venulosus* (Wt. & Arn.) Deb & Gang.
A, habit; B, flower; C, flower split opened; D, fruit; E, seed; F, embryo.

peduncled cymes, bracteate, bracteolate, 3-4 flowered, subracemose; peduncle 2-8 mm long, glabrous or strigose; branchlets short or absent, strigose or pubescent when present; bracts 6-9 mm long, subulate, pubescent; bracteoles 1-2 mm long, triangular, pubescent. Flowers subsessile or pedicellate, 9-11 mm long, tubular, white; pedicels 0.5-1 mm long, glabrous or sparsely pubescent; hypanthium \pm 1 mm long, obovoid, glabrous or sparsely pubescent; calyx tube \pm 0.5-1.5-2 mm, glabrous or sparsely pubescent above; lobes 4, 2.5-5 mm long, linear, acute or obtuse at apex, puberulous or pubescent. Corolla tube 5.5-6.5 mm long, glabrous above, pubescent within at upper part; lobes 4, \pm 2 mm long, ovate, ovate-lanceolate, puberulous sometimes ciliate at apex. Stamens 4, included or subexserted; filaments 0.5-1 mm long, thin, narrow, glabrous, adnate below throat; anthers \pm 1 mm long, oblong. Ovary \pm 0.5 mm long, obovoid, 4-loculed; style 3-6 mm long, thin, narrow, glabrous or puberulous above; lobes 4, \pm 0.5 mm long, ovate-oblong, obtuse, glabrous; disk \pm 1 mm across, subglobose, smooth. Fruit sessile or short stalked, 6-7 \times 5-6 mm, ovoid, with persistent calyx lobes, 4-shallow channelled, blackish green; pericarp thin, glabrous; mesocarp thin; pyrenes 4, planoconvex, acute at both ends, wall wrinkled. Seeds \pm 4 mm long, narrowly oblong, wrinkled; embryo \pm 3 mm long; radicle 2.5 mm long, blunt, semiterete; cotyledons ovate-oblong, obtuse.

Flowers and fruits: throughout the year.

Distribution. — India: Tamilnadu in evergreen forest at 1500-2500 m in altitude.

Exsiccata studied

Coimbatore Dt., Anamallai, *R. H. Beddome* 4033 (BM). Madurai Dt., Pulney ghat, 1800 m, *R. H. Beddome* 4034 (BM); Glen falls shola, Bourne 273 (CAL). Kodaikanal, *D. A. Barber* 7593 (MH), *S. Munch* 10 (CAL). Nilgiri Dt., *R. H. Beddome* 4014 (BM); *Perrotet* 509, 510 (BM); *Schmidt* s.n. (CAL, E); Coonoor, 2100 m, *C. B. Clarke* II058A (BM); II058B, II058D (CAL); II063A (BM), II063B (CAL); Ooty, *C. B. Clarke* II455A, II455B (CAL); *G. King* s.n. (CAL); *D. Brandis* s.n. (CAL) & *Shethi & Nagi* 26171 (DD); Dodabetta, *J. S. Gamble* 17461 (BSI, CAL) & *N. C. Rathakrishnan* 39034 (MH); Lamb's rock, *J. S. Gamble* II596, II597, II867, I3115, I3368 (CAL) & *G. A. Gammie* 632 (BSI). Sisparaghat, *J. S. Gamble* I3373 (CAL); Dodabetta, *J. S. Gamble* II859 & I4471 (CAL); *M. Gangopadhyay & D. C. Mondal* 2161 (CAL); Ooty, 2100 m, *J. S. Gamble* 17461 (BSI, CAL); Brahmagiri sholas, *C. E. C. Fischer* 252 (CAL); Shembaganur, 1800 m, *Sauliere* 333 (CAL); Bikapathimund, *C. E. C. Fischer* 4590 (CAL); Pakasurmalai, *K. M. Sebastian* 3213 & 3940 (CAL).

New combinations in Malayan and Sri Lankan *Litosanthes* Bl.

1. ***Litosanthes filiformis* (King & Gamble) Deb & Gang., comb. nov.**. Basionym: *Lasianthus filiformis* King & Gamble in Journ. Asiat. Soc. 73: 131. 1904 (Type: Malaya, Pahang, *Scortechini* 29! CAL selected as the lectotype; W. Singa Ryah, *King's collector* 851! (CAL).
2. ***L. filiformis* var. *bracteata* (King & Gamble) Deb & Gang., comb. nov.**. Basionym: *Lasianthus filiformis* var. *bracteata* King & Gamble in Journ. Asiat. Soc. 73: 131. 1904 (Type: Pahang, *W. N. Ridley* 8578! CAL selected as the lectotype; Pahang, *Scortechini* 180 & 190!, CAL).
3. ***L. pendulus* (Ridley) Deb & Gang., comb. nov.**. Basionym: *Lasianthus pendulus* Ridley in Journ. Fed. Mal. States Mus. 4: 67. 1909 (Type: Malaya, Kahang, *Ridley* not seen).
4. ***L. pseudolucidus* (King) Deb & Gang., comb. nov.**. Basionym: *Lasianthus pseudo-lucidus* King in Journ. Asiat. Soc. 73: 128. 1904 (Type: Perak, *L. Wray* 4109! CAL).
5. ***L. robinsonii* (Ridley) Deb & Gang., comb. nov.**. Basionym: *Lasianthus robinsonii* Ridley in Journ. Fed. Mal. States Mus. 4: 37. 1909 (Type: Pahang, *Ridley* not seen).
6. ***L. scalariformis* (King & Gamble) Deb & Gang., comb. nov.**. Basionym: *Lasianthus scalariformis* King & Gamble in Journ. Asiat. Soc. 73: 131. 1904 (Type: Perak, *Scortechini* 411 ! CAL; *L. Wray* 332 ! CAL).

7. **L. subspicatus** (King & Gamble) Deb & Gang., **comb. nov.**. Basionym: *Lasianthus subspicatus* King & Gamble in Journ. Asiat. Soc. 73: 118. 1904 (Type: Perak, *L. Wray* 863; Penang, *Curtis* 2476).
8. **L. varians** (Hook. f.) Deb & Gang., **comb. nov.**. Basionym: *Lasianthus varians* Hook. f., Fl. Brit. Ind. 3: 191. 1880 (Type: Sri Lanka, C. P., Galagama, *Thwaites*). *Maphitidia varians* Thw. p.p.
9. **L. villosus** (Ridley) Deb & Gang., **comb. nov.**. Basionym: *Lasianthus villosus* Ridley in Journ. Roy. Asiat. Soc. Str. Br. 79: 88. 1915 (Type: Perak, Tapa Words, *Ridley* ...).

Note. — *Lasianthus moonii* Hook. f., Fl. Brit. Ind. 3: 190. 1880 also most probably represents *Litosanthes* but due to non availability of flowering material it could not be decided properly.

REFERENCES

- BAKHUIZEN VAN DEN BRINK, R. C. (1965). In: BACKER, C. A. & R. C. BAKHUIZEN VAN DEN BRINK, *Flora of Java*. Vol. 2, N.V.P. Noordhofgrinogen, The Netherlands.
- BEDDOME, R. H. (1869-1873). *Flora Sylvatica for South India*. 2 vols. Madras.
- BEDDOME, R. H. (1869-1874). *Icones plantarum indiae orientale* or plates and descriptions of new and rare plants chiefly from South India and Ceylon, Madras.
- BLUME, C. (1825). *Flora Botanische Zeitung* 8: 189.
- BLUME, C. (1825-1826). *Bijdragen tot de Flora van Nederlandsch Indie*. Batavia. 2 vols.
- CANDOLLE DE, A. P. (1830). *Prodromus Systematis Naturalis Regni vegetabilis* 4: 465.
- COOKE, T. (1903). *The Flora of the Presidency of Bombay*. Vol. 1. London.
- FYSON, P. F. (1915). *The Flora of Nilgiri and Pulney Hill tops*.
- GAMBLE, J. S. (1920). Notes on the Flora of Madras. *Bull. Miscellaneous Information Kew* 1920: 49-57.
- GAMBLE, J. S. (1921). *The Flora of the Presidency of Madras*. London.
- HOOKER, J. D. (1873). In: BENTHAM, G. & J. D. HOOKER, *Genera Plantarum*: 2. London.
- HOOKER, J. D. (1880). *The flora of British India*: 3. London.
- MIQUEL, F. A. W. (1855). *Flora van Nederlandsch Indie* 2: 314. Amsterdam & Utrecht.
- RICHARD, A. (1829). *Mémoire sur la famille des Rubiacées contenant les caractères des genres de cette famille et d'un grand nombre d'espèces nouvelles*. Paris.
- RIDLEY, H. N. (1909). The Flora of the Telom and Batang Padang valleys. *Journ. Fed. May States Museum* 4: 67.
- RIDLEY, H. N. (1915). Material for a Flora of the Malayan Peninsula. *Journ. Royal Asiatic Society straites Branch* 19: 88.
- RIDLEY, H. N. (1923). *Flora of Malayan Peninsula*. Vol. 2. London.
- SCHUMANN, K. (1891). In: ENGLER, A. & K. PRANTL, *Die natürlichen Pflanzenfamilien. IV. Rubiaceae*: pp. 110 & 121.
- WALPERS, W. G. (1852). *Annales botanices systematicae*. Vol. 3.
- WIGHT, R. (1835-1853). *Icones plantarum Indiae Orientalis*.
- WIGHT, R. (1845). *Calcutta J. Nat. Hist. Soc.* 6: 511.
- WIGHT, R. & A. WALKER-ARNOTT (1834). *Prodromus Florae Peninsulae Indiae Orientalis*. Vol. 1. London.

