

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	46 (1991)
Heft:	2
Artikel:	Grewia indandamanica Ellis & Ray : a new species from Andaman Islands in the Bay of Bengal, India
Autor:	Ellis, J.L. / Ray, L.N.
DOI:	https://doi.org/10.5169/seals-879831

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Grewia indandamanica Ellis & Ray — a new species from Andaman Islands in the Bay of Bengal, India

J. L. ELLIS
&
L. N. RAY

ABSTRACT

ELLIS, J. L. & L. N. RAY (1991). *Grewia indandamanica* Ellis & Ray — a new species from Andaman Islands in the Bay of Bengal, India. *Candollea* 46: 341-343. In English, English and French abstracts.

A new species, *Grewia indandamanica*, has been described from Saddle Peak in North Andaman Islands, Andaman Islands.

RÉSUMÉ

ELLIS, J. L. & L. N. RAY (1991). *Grewia indandamanica* Ellis & Ray — espèce nouvelle des Iles Andaman, Golfe du Bengale, Indes. *Candollea* 46: 341-343. En anglais, résumés anglais et français.

Une espèce nouvelle, *Grewia indandamanica*, est décrite de Saddle Peak, Ile Andaman Nord, Iles Andaman.

Airy Shaw (Willis, Dict. Fl. Plants, 1973) has given 150 species of *Grewia* L. as occurring in Africa, Asia and Australia, especially in the tropical belt. Masters (in Hook. f., Fl. Brit. Ind. 1: 383-393, 1872) has accounted for 36 species and varieties from India and surrounding areas. Parkinson (For. Fl. Andaman Is., pp. 104-105, 1923) records three species from the Andaman Islands. He gives doubtfully another plant saying that it is a plant from a mountainous region in Middle Andaman, "...with leaves 1-1.5 in. long, elliptic, crenate-serrate and flowers in the leaf axils".

The presently described species has been collected from Saddle Peak — the highest peak in these Bay of Bengal islands (726 m msl.). This region supports a stunted, evergreen-type of forest which may be called Hill-top Forest. The close proximity of the Andaman and Nicobar groups of islands with Equator and the direct influence of both south-west monsoon and north-east monsoon and high humidity have contributed very significantly to the development of humid tropics in these islands. The resultant diversity in the types of vegetation is mainly owing to the influence of all these factors in addition to edaphic factors and availability of ground water. The Saddle Peak again is affected by the altitudinal gradation, in addition to the solar insolation and wind velocity. As the peak is commonly covered by thick clouds, the variation in temperature from the surrounding areas is very marked, thus encouraging plant diversity. This is very much evident by the climbing bamboo, *Dionochloa andamanica* Kurz, making an impenetrable thicket on the Saddle Peak, amidst which some plants typical of arid zone occur like *Phoenix sylvestris* Roxb., *Rostellutaria procumbens* (L.) Nees, etc.

***Grewia indandamanica* Ellis & Ray, spec. nov. (Fig. 1).**

Holotypus Ellis ANC 12775-A (Saddle Peak, Andaman borealis, die 18.10.1987), positus in CAL, Isoty whole Ellis ANC 12775 B-D positi in PBL.




Fig. 1. — *Grewia indandamanica* Ellis & Ray
1, habit sketch; 2, flower with a sepal removed; 3, sepal; 4, petal showing the gland; 5, stamen; 6, ovary; 7, capsule.

G. laevigatae Vahl affinis, sed differt, (i) floribus solitariis, (ii) pedicellis pedunculisque per-brevioribus, (iii) foliis elliptic-ovatis, parum acuminatis, basi rotundatis ad parum cordatis, non attenuatis, perparvioribus, ad marginem crenato-serratis; stipulis perparvioribus, et (iv) fructibus sparsim puberulis, laviter angulatis, sulcatis.

Allied to *Grewia laevigata* Vahl, but differs from it in having — (i) flowers single, (ii) pedicels and peduncles much shorter; (iii) leaves elliptic-ovate, slightly acuminate, and at base rounded to slightly cordate and not attenuate, much smaller, margin crenate-serrate; stipules much smaller, and (iv) fruits sparsely puberulous, faintly angled, grooved.

Frutex fruticulosus ramosus, usque ad 1 m altus, rami ravidii, junioros sparsim stellate-pilosi, cortex rocosa. Folia usque ad 7.0×2.8 cm, ovato-elliptica, breviter acuminata, ad basin rotundata et cordata, crenato-serrata, chartacea ad parum coriacea; nervi laterales 5, arcuati, reticulationes saepe parallelae, nervi nervulisque prominenti in superficie bambabus; petiolus fuscatus, 0.5 cm longus, junior pubescens; stipuli 0.1 cm longi. Flores lutei, 1 cm longi, solitarii axillares, alabastra 0.5 cm longa laeviter stellate tomentosa; pedicelli usque ad 0.9 cm longi, sparsim puberuli; pedunculi usque ad 1 cm longi, ad medium articulati et parum incrassati; bracteae 0.2 cm longae, bracteolae 0.1 cm longae; sepala 1.0×0.15 cm, linear-ovata, spatulate, cucullata, ad basin obtusa ad truncata, extus puberula, intus brunnea glabraque, ad marginem incurva, stellate-tomentosa; petala 0.25×0.1 cm, ovato-oblonga, integra, extus glabra, intus dense pubescentia secus marginem, usque ad 3/4 longitudinis et glandem cingens, gland plus quam 1/3 petali, glabra; stamina numerosa, filamenta 0.6 cm longa, filiformia, glabra, anthera innata, reniformis; torus 0.2 cm longus, dimidium inferius glabrus et dimidium superius dense stellate-pubescent; ovarium globosum, dense pubescent, 4-loculare, ovule in quoque loculo solitario, stylus 0.6 cm longus, gracilis, puberulose basi usque ad 2/3 longitudinis, sursum glaber, stigma parum expansum, 4-angulatum. Drupa nigra, 1 cm diam., bilobata, plerumque inaequaliter bilobata, globosa vel parum complanata vel angulata, rugosa, sparsim pilosa, lignosa; semen unicum in quoque loculo, glabrum.

A much-branched shrub or an undershrub, up to 1 m high; branches greyish, sparsely stellate-hairy when young, bark wrinkled. Leaves up to 7.0×2.8 cm, ovate-elliptic, shortly acuminate, rounded and cordate at base, crenate-serrate, chartaceous to slightly coriaceous; lateral nerves 5, arcuate, reticulations often parallel, nerves and nervules prominent on both the surfaces; petiole darkish in colour, 0.5 cm long, pubescent when young; stipules 0.1 cm long; flowers yellow, 1 cm long, solitary, axillary, flowering buds 0.5 cm long, faintly stellate-tomentose; pedicels up to 0.9 cm long, sparsely puberulous; peduncles up to 1 cm long, jointed in the middle and slightly thickened; bracts 0.2 cm long; bracteoles 0.1 cm long; sepals 1.0×0.15 cm, linear-ovate, spatulate, cucullate, obtuse to truncate at base, puberulous without, brown and glabrous within, margins incurved, stellate-tomentose; petals 0.25×0.1 cm, ovate-oblong, entire, glabrous without, densely pubescent along margin up to 3/4th the length and in a circle round the gland within, gland more than 1/3rd the petal, glabrous; stamens numerous, filaments 0.6 cm long, filiform, glabrous, anthers innate, reniform; torus 0.2 cm long, lower half glabrous and upper half densely stellate pubescent; ovary globose, densely pubescent, 4-celled with 1 ovule in each, style 0.6 cm long, slender, puberulous from base up to 2/3rd the length, glabrous above, stigma slightly expanded and 4-angled. Drupe black, 1 cm across, bilobed, generally unequally so, globose or slightly flattened or angled, wrinkled, sparsely hairy, woody; seed one in each locule, glabrous.

ACKNOWLEDGEMENTS

Thanks are due to the Director, Botanical Survey of India, for facilities. We are thankful Dr. S. K. Srivastava, Botanical Survey of India, Port Blair, for the line drawings. We are grateful to Dr. N. C. Majumdar, Red. Scientist "SE", for Latin translation of the protologue.

