

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 47 (1992)

Heft: 1

Artikel: An annotated check-list of the Balearic vascular Flora : I. Pteridophyta-Coniferophytina

Autor: Rosselló, J.A. / Cubas, P. / Torres, N.

DOI: <https://doi.org/10.5169/seals-879563>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An annotated check-list of the Balearic vascular Flora. I. Pteridophyta-Coniferophytina

J. A. ROSSELLÓ
P. CUBAS
&
N. TORRES

ABSTRACT

ROSSELLÓ, J. A., P. CUBAS & N. TORRES (1992). An annotated check-list of the Balearic vascular Flora. I. Pteridophyta-Coniferophytina. *Candollea* 47: 61-69. In English, English and Spanish abstracts.

First part of a series covering the revised check-list of the vascular flora of the Balearic Islands. Species, subspecies and hybrid taxa have been included. Their insular distribution in the larger islands, i.e. Mallorca (Ma), Menorca (Me), Eivissa (Ei), Formentera (Fo) and Cabrera (Ca) is provided. The combination *Pinus halepensis* Mill. var. *ceciliae* (A. Llorens & L. Llorens) L. Llorens is validated.

RESUMEN

ROSSELLÓ, J. A., P. CUBAS & N. TORRES (1992). Catálogo anotado de la Flora Vascular de las islas Baleares. I. Pteridophyta-Coniferophytina. *Candollea* 47: 61-69. En inglés, resúmenes en inglés y en español.

Se presenta la primera parte del catálogo revisado de la flora vascular de las Islas Baleares. En este catálogo se incluyen las plantas nativas y naturalizadas con rango específico, subespecífico y nothotáxones, indicándose su distribución en las islas mayores, i.e. Mallorca (Ma), Menorca (Me), Eivissa (Ei), Formentera (Fo) and Cabrera (Ca). Se valida una nueva combinación: *Pinus halepensis* Mill. var. *ceciliae* (A. Llorens & L. Llorens) L. Llorens.

For the last 170 years, the vascular plants of the Balearic Islands have attracted the attention of many European botanists and have been the subject of many papers. However, compilations of the floristic knowledge of the islands are scanty. Some of the most important early contributions covering the Balearic Flora were those of BARCELO (1879-1881), MARES & VIGINEIX (1880), RODRÍGUEZ (1904) and KNOCHE (1921-1922). BARCELO (1879-1881) and KNOCHE (1921-1922) have been followed as basic floras for more comprehensive works such as "Flora Europaea" (TUTIN & al., 1964-1980) and "Flora Ibérica" (CASTROVIEJO & al., 1986-1990). Unfortunately, these basic floras of Baleares have become outdated due to the amount of information gained during the last 40 years.

More recently, DUVIGNEAUD (1974, 1979) published a check-list of the Balearic vascular Flora. This valuable contribution lacks, however, explanatory notes on the insular distribution of the taxa, and no bibliographic references are given for the most doubtful or puzzling records. Besides, the precise location, and references, to published data or to the herbarium specimens are not provided for the new reported taxa. Moreover, old misidentifications and imprecise determinations have been included.

Later, BONAFE (1980) published a check-list of the Balearic Flora which, unfortunately, did not improve Duvigneaud's work.

Check-lists of territories including the Balearic Islands (GREUTER & al., 1984-1989; BOLÓS & VIGO, 1984-1990; CASTROVIEJO & al., 1986-1990) have not been at present finished and the scope of some of them is so great that not equal accuracy can be claimed for all the countries included. The work of SMYTHIES (1984, 1986) has paid little attention to the problematic Balearic flora and his results come from earlier bibliographic works.

The lack of adequate synthetic work concerning the whole Balearic flora has moved us to present this check-list, which is the result of extensive field work through all the islands, herbarium revision and a thorough bibliographic compilation. Only native and naturalized plants, including hybrid taxa, have been added to the check-list, whereas cultivated plants and aliens have been excluded from the main text. However, the most important ones are included at the notes. Taxa pointed out in the Balearic or insular floras and believed to be misidentifications are reported in the notes. Insular distributions are reported for the larger islands, i.e. Mallorca (Ma), Menorca (Me), Eivissa (Ei), Formentera (Fo) and Cabrera (Ca). The flora of Dragonera, Vedrà and those smaller islets surrounding the five major islands is, as a rule, not comprehensively detailed, however, details may be provided if requested to the authors. Only the distribution of those taxa occurring in the islets and with a restricted distribution or having biogeographical, taxonomical or ecological interest are presented. Arrangement of the families follows CASTROVIEJO & al. (1986) for Pteridophyta and Coniferophytina and CRONQUIST (1988) for Magnoliophytina. Nomenclature for Pteridophytes is that of DERRICK & al. (1987), whereas TUTIN & al. (1964-1980), GREUTER & al. (1984, 1986, 1989) and CASTROVIEJO & al. (1986-1990) are followed for Spermatophyta. However, some taxonomic or nomenclatural inconsistencies between these authorities and our point of view exist, so these will be pointed out in the notes.

Comments, mainly taxonomic and chorological, are presented for each taxon when needed after the taxonomic part of each paper. Only specific and subspecific ranks are used in this check-list. Taxa below these ranks, if worth of mention, are reported in the notes. The following symbols will be used: (●) native plants, (○) naturalized plants, (□) plants whose presence is doubtful.

For the sake of convenience this Balearic catalogue is divided in four parts, the first of which is here published and cover Pteridophyta and Coniferophytina. The other three parts will include *Magnoliidae-Caryophyllidae*, *Dilleniidae-Rosidae*, and *Asteridae-Liliidae*.

PTERIDOPHYTA (1)

	Ma	Me	Ei	Fo	Ca
Fam. <i>Selaginellaceae</i>					
<i>Selaginella denticulata</i> (L.) Spring	●	●	●	●	
Fam. <i>Isoetaceae</i>					
<i>Isoetes durieui</i> Bory (2)		●			
<i>Isoetes histrix</i> Bory (3)	●	●			
<i>Isoetes velatum</i> A. Br. in Bory & Durieu (4).....		●			
Fam. <i>Equisetaceae</i>					
<i>Equisetum arvense</i> L. (5)	●				
<i>Equisetum ramossissimum</i> Desf. (6).....	●	●	●		
<i>Equisetum telmateia</i> Ehrh	●	●			

Fam. *Ophioglossaceae*

Ophioglossum lusitanicum L. ● ● ● ●

Fam. *Polypodiaceae*

Polypodium cambricum L. subsp. *serrulatum* (Sch. ex Arcangeli) Pichi-Serm. (7) ● ● ● ● ●

Fam. *Sinopteridaceae*

Cheilanthes acrosticha (Balbis) Tod. (8) ● ● ●

Fam. *Pteridaceae*

Pteris vittata L. (9) ●

Fam. *Adiantaceae*

Adiantum capillus-veneris L. ● ● ● ● ●

Fam. *Hemionitidaceae*

Anogramma leptophylla (L.) Link (10) ● ● ● ● ●
Cosentinia vellea (Ait.) Tod. (11) ● ● ● ● ●

Fam. *Marsileaceae*

Marsilea strigosa Willd. (12) ● ●
Pilularia minuta Durieu in Bory & Durieu (13) ●

Fam. *Hypolepidaceae*

Pteridium aquilinum (L.) Kuhn in Kersten (14) ● ● ●

Fam. *Aspleniaceae*

Asplenium azomanes Rosselló, Cubas & Rebassa (15) .. ● ● ● ●
Asplenium balearicum Shivas (16) ● ● ● ● ●
Asplenium ceterach L. ● ● ● ● ●
Asplenium fontanum (L.) Bernh. (17) ● ● ● ● ●
Asplenium marinum L. ● ● ● ● ●
Asplenium majoricum Litard. (18) ● ● ● ● ●
Asplenium onopteris L. (19) ● ● ● ● ●
Asplenium petrarckae (Guérin) DC. in Lam. & DC.
 subsp. *petrarckae* (20) ● ● ● ● ●
 subsp. *bivalens* (D. E. Meyer) Lovis & Reichst. (21) ● ● ● ● ●
Asplenium ruta-muraria L. (22) ● ● ● ● ●
Asplenium sagittatum (DC.) A. J. Bange (23) ● ● ● ● ●
Asplenium scolopendrium L. ● ● ● ● ●
Asplenium trichomanes L.
 subsp. *inexpectans* Lovis (24) ● ● ● ● ●

subsp. <i>quadrivalens</i> D. E. Meyer	●	●	●
nothosubsp. <i>lucanum</i> Cubas, Rosselló & Pangua (25)	●		
<i>Asplenium</i> × <i>artanense</i> Rosselló, Cubas, Gradaille & Sastre (26)	●		
<i>Asplenium</i> × <i>barrancense</i> (Bennert & D. E. Meyer) Pericás & Rosselló (27)	●		
<i>Asplenium</i> × <i>helii</i> Lusina nothosubsp. <i>helii</i> (28)	●		
nothosubsp. <i>calobraense</i> Bennert, H. Rasbach & K. Rasbach (29)	●		
<i>Asplenium</i> × <i>orellii</i> Lovis & Reichst. (30)	●		
<i>Asplenium</i> × <i>reichstenii</i> Bennert, H. Rasbach & K. Rasbach (31)	●		
<i>Asplenium</i> × <i>sollerense</i> Lovis, Sleep & Reichst. (32) ..	●		
<i>Asplenium</i> × <i>tubalense</i> Rosselló, Cubas & Rebassa (33)	●		
<i>Asplenium</i> × <i>tyrrhenicum</i> Cubas, Pangua & Rosselló (34)		●	

Fam. *Athyriaceae*

<i>Athyrium filix-femina</i> (L.) Roth (35)	●
<i>Cystopteris fragilis</i> (L.) Bernh. (36)	●

Fam. *Aspidiaceae*

<i>Dryopteris filix-mas</i> (L.) Schott (37)	●
<i>Dryopteris pallida</i> (Bory) C. Chr. ex Maire & Petitmengin subsp. <i>balearica</i> (Litard.) Fraser-Jenkins (38)	●
<i>Dryopteris tyrrhena</i> Fraser-Jenkins & Reichst. (39)	●
<i>Polystichum aculeatum</i> (L.) Roth (40)	●
<i>Polystichum setiferum</i> (Forsskål) Woyнар	●
<i>Polystichum</i> × <i>bicknellii</i> (Christ) Hahne (41)	●

CONIFEROPHYTINA

Fam. *Cupressaceae*

<i>Juniperus oxycedrus</i> L. subsp. <i>oxycedrus</i>	●	●	●	●	●
subsp. <i>macrocarpa</i> (Sm.) Ball	●				
<i>Juniperus phoenicea</i> L. (42)	●	●	●	●	●

Fam. *Ephedraceae*

<i>Ephedra fragilis</i> L. (43)	●	●			●
---------------------------------------	---	---	--	--	---

Fam. *Pinaceae*

<i>Pinus halepensis</i> Miller (44)	●	●	●	●	●
<i>Pinus pinea</i> L. (45)	○	○	○	○	

Fam. *Taxaceae*

<i>Taxus baccata</i> L. (46)	●
------------------------------------	---

NOTES

¹GANDOGGER (1891) published several illegitimate taxa from the Balearic flora, all of them without taxonomic relevance: *Selaginella balearica* (Mallorca), *S. platyclada* (Menorca), *Isoetes balearica* (Menorca), *I. tenuicola* (Menorca), *Asplenium balearicum* (Menorca), *A. tenerarium* (Menorca), *Polypodium balearicum* (Menorca) and *Polystichum balearicum* (Mallorca).

²Even if early reports had indicated the existence of *I. durieui* in Mallorca (DUVIGNEAUD, 1979), no evidence has been provided to support the actual presence of this taxon in the island. In fact, all the herbarium specimens previously ascribed to *I. durieui* have proved to be *I. histrix*.

³For more details regarding the distribution of this taxon in Mallorca and Menorca see ROSSELLÓ & ALOMAR (1987) and BALLESTEROS (1989).

⁴In the Balearic Islands these plants have been identified as subspecies *velatum* (PRADA, 1983; 1986). Recently, SMYTHIES (1984a) and DERRICK & al. (1987) have ascribed them to the subspecies *tenuissima* (no details are given), suggesting that a revision of the *I. velatum* aggr. is needed. Furthermore, plants from Menorca display some differences (morphology and spore characteristics) when compared to the typical subsp. *velatum*.

⁵A doubtful first indication of this taxon for Menorca (RODRIGUEZ, 1874), it was later referred as *E. telmateia* (RODRIGUEZ, 1904). At present, its existence has only been confirmed for Mallorca.

⁶Previous reports indicating the existence of *E. fluviatile* L. (sub *E. limosum*) in Eivissa (CAMBESSEDES, 1827) and in Mallorca (GARCÍAS, 1919) are in fact misidentifications of *E. ramossissimum*. Contrary to SALVO & al. (1984), there is not any record of *E. ramossissimum* from Formentera.

⁷Contrary to the comments of FONT QUER (1919), *P. vulgare* L. is not present in the Balearic Islands and the old records are misidentifications of *P. cambricum*. Plants collected in the neighbourhood of Valldemossa (Mallorca) have been ascribed to *P. cambricum* L. subsp. *cambricum* (PALAU, 1954; BONAFE, 1977). They show lacinated pinnae and numerous and well developed sporangia. Despite their anomalous morphology they must be included within *P. cambricum* subsp. *serrulatum*.

⁸The record of *Ch. hispanica* from Mallorca (JAQUOTOT & ORELL, 1968) is erroneous; the plants are in fact *Ch. acrosticha*. *Ch. pteridioides* (Reichard) C. Chr. (= *Ch. maderensis* Lowe) has been cited by DERRICK & al. (1987), however no references are given. Furthermore, it should be noted that the ecological requirements of this taxon make its presence in the Balearic Islands very improbable. We consider that the only one species of *Cheilanthes* growing in Balears is *Ch. acrosticha*. MORALES & FERNANDEZ CASAS (1989) erroneously consider that all the Balearic *Cheilanthes* plants are *Ch. maderensis*.

⁹The indication of *Blechnum spicant* L. in Mallorca (BARCELO, 1867) is an erroneous identification of *P. vittata*. *B. spicant* L. is not to be regarded as extinguished in the islands as reported (ORMONDE, 1986), in fact, it never existed in the Balearic Islands. Regarding *P. vittata*, the authors bear in mind some doubts regarding the autochthony of the taxon in Mallorca.

¹⁰Cited for Eivissa from one locality (Puig de Sa Fita, 29.4.1948, P. Font Quer, BC 79094). It has not been found again.

¹¹Reported from Menorca by LLORENS (1979). The Balearic populations of *C. vellea* might belong to the tetraploid cytotype (subsp. *vellea*), however, cytological confirmation is required.

¹²Recently reported from Menorca (ALOMAR & al., 1991).

¹³Found in Menorca by RITA (1987).

¹⁴Known in only one locality of Eivissa (Puig de Ses Roques, Sant Joan, N. Torres; pers. herb.). SHEFFIELD & al. (1986) suggested that the Mallorcan populations, living in calcareous soils, were diploid (*Pteridium herediae* (Colmeiro) Löve & Kjellqvist). However, chromosome counts in gametophytic tissues and complementary observations have established that they are tetraploid (SHEFFIELD & al., 1989) i.e. *P. aquilinum* subsp. *aquilinum*.

¹⁵Considered as a subspecies of *A. trichomanes* (*A. trichomanes* subsp. *coriaceifolium*) by RASBACH & al. (1990). Taxonomic and chorological details are shown in ROSSELLÓ & al. (1991).

¹⁶All the records of *A. obovatum* Viv. subsp. *obovatum* and *A. obovatum* subsp. *lanceolatum* P. Silva (RODRIGUEZ, 1904; KNOCHE, 1921; MONTSERRAT, 1974; SMYTHIES, 1984; BOLÓS & VIGO, 1984) in Menorca are erroneous and those plants must be ascribed to *A. balearicum*. The record of *A. obovatum* s.l. in the southeast of Mallorca (MAHEU, 1912) is also erroneous. The existence of *A. balearicum* in Mallorca (JALAS & SUOMINEN, 1972; SALVO & al., 1984) is not supported by any evidence (ROSSELLÓ & al., 1986; ROSSELLÓ & SERRA, 1987; CUBAS & al., 1988).

¹⁷Reported from Sóller (BIANOR, 1917), this taxon has not been found again in this locality, where it has probably been extinguished due to an excess of collecting. At present, only one alive specimen is known in Mallorca (ROSSELLÓ & al., 1986).

¹⁸Until now regarded as a Balearic endemism; recently found in Continental Spain (C. Prada & E. Pangua, pers. comm.).

¹⁹First finding in Formentera (Torrent des Gat, 31.3.1988, Rosselló & Torres; pers. herb.). Previously mistaken with *A. adiantum-nigrum* L. and *A. cuneifolium* Viv. (MARES & VIGINEIX, 1880; JALAS & SUOMINEN, 1972) due to its morphological variability in the Balearic Islands. *A. adiantum-nigrum* and *A. cuneifolium* have not been found in the islands.

²⁰Chromosome counts available only from Mallorca (LOVIS & al., 1970; CUBAS, unpublished data). Lacking adequate chromosome data from other places, the distribution of both subspecies has to be regarded as provisional. Not reported from Formentera, however, its presence in the island is probable.

²¹The diploid cytotype (subsp. *bivalens*) has been unequivocally identified in Sa Calobra (BENNERT & al., 1990).

²²The old records from Menorca have never been confirmed and they are probably erroneous. There is a herbarium sheet of this species at the Institut Botanic of Barcelona (Eivissa, Els Puntals, 25.6.1920, Gros, BC 77592). However, since the locality name (Els Puntals) does not actually exist in Eivissa but is common in Mallorca, this report is probably a labeling error.

²³The plants recorded as *A. hybridum* from Mallorca (TRABUT, 1917) are here considered to be a lobed form of *A. sagittatum*. This theratological form has been sporadically found growing among normal populations of *A. sagittatum*, and does not, in our opinion, deserve taxonomic recognition.

²⁴*A. trichomanes* subsp. *trichomanes* is not present in the Balearic Islands. The records of this subspecies (GREUTER & al., 1984; SMYTHIES, 1984a; SALVO & al., 1984; MORALES & FERNANDEZ CASAS, 1989) have to be referred to subsp. *inexpectans* or to subsp. *quadrivalens*. For details on the distribution of the *A. trichomanes* complex in the Balearic Islands see ROSSELLÓ (1989).

²⁵Intraspecific cross between the subspecies *quadrivalens* and *inexpectans* (CUBAS & al., 1989). The type material is from Mallorca, even though this taxon is also found in other European localities. *A. trichomanes* nothosubsp. *menzeri* (LOVIS & al., 1989) is a later synonym of nothosubsp. *lucanum*.

²⁶Diploid hybrid between *A. sagittatum* and *A. trichomanes* subsp. *inexpectans*. See ROSSELLÓ & al. (1990) for details regarding its origin, morphology and distribution.

²⁷Tetraploid hybrid between *A. ceterach* subsp. *ceterach* and *A. majoricum*. Described as \times *Asplenoceterach barrancense* (BENNERT & MEYER, 1972) and later combined by ROSSELLÓ & al. (1986). See MUS & al. (1990) for the taxonomical position of \times *Asplenoceterach* D. E. Meyer in *Asplenium* L. Cytological data can be obtained from RASBACH & al. (1987).

²⁸Triploid hybrid between *A. petrarchae* subsp. *petrarchae* and *A. trichomanes* subsp. *inexpectans*. *A. \times litardierei* Bennert & Meyer, the triploid hybrid described from Mallorca (BENNERT & MEYER, 1974) is to be regarded as a synonym of *A. \times helii* nothosubsp. *helii*. See BENNERT & al. (1988, 1989) for the nomenclature of the hybrids between *A. petrarchae* and *A. trichomanes*.

²⁹Triploid hybrid between *A. petrarchae* subsp. *bivalens* and *A. trichomanes* subsp. *quadrivalens* (BENNERT & al., 1990).

³⁰Tetraploid hybrid between *A. majoricum* and *A. trichomanes* subsp. *quadrivalens* (LOVIS & REICHSTEIN, 1970).

³¹Triploid hybrid between *A. fontanum* and *A. majoricum* (BENNERT & al., 1987).

³²Tetraploid hybrid between *A. majoricum* and *A. petrarchae* subsp. *petrarchae* (LOVIS & al., 1970).

³³Tetraploid hybrid between *A. azomanes* and *A. trichomanes* subsp. *quadrivalens* (ROSSELLÓ & al., 1991). *A. trichomanes* nothosubsp. *barreraense* (RASBACH & al., 1990) is a synonym of *A. \times tubalense*.

³⁴Triploid hybrid between *A. balearicum* and *A. onopteris*. Known only from Menorca (CUBAS & al., 1987, 1989; BENNERT & al., 1988), from where it was described. It could probably be found in Sardinia and other Italian islands where its parents grow (PICHI-SERMOLLI, 1988). *A. \times rosselloi* (BENNERT & al., 1988) is a later synonym of *A. \times tyrrhenicum*.

³⁵Reported for the Balearic flora by SALVO & al. (1984) based on a single frond included within a sheet also containing fronds of *D. pallida* subsp. *balearica* (Barranco de Sóller, E. Bourgeau; MA 236455). *A. filix-femina* has not been found again in the islands and perhaps it is a labeling error.

³⁶Very rare in Mallorca and showing a slightly different morphology when compared to *C. fragilis* subsp. *fragilis*. The scarcity of herbarium specimens and the difficult taxonomy of the *Cystopteris fragilis* complex prevent a more accurate identification.

³⁷Recently reported in the Balearic flora (ROSSELLÓ & al., unpubl. data).

³⁸FRASER-JENKINS (1982) reports the presence of *D. pallida* subsp. *pallida* in the Balearic flora. However, in our opinion, the specimen studied by this author (MPU) is only a peculiar extreme form of *D. pallida* subsp. *balearica*, approaching in some morphological details to subsp. *pallida* but, nevertheless, falling within the range of variation of subsp. *balearica*. Herbarium specimens with similar characteristics have been collected from scattered localities in Mallorca, which in any case, are well within the distribution area of subsp. *balearica*. Similarly, even if some plants from Sardinia (MPU) are morphologically approaching to subsp. *balearica* (DERRICK & al., 1987) they are to be regarded as morphological variations of subsp. *pallida*. The great polymorphism shown by *D. pallida* in the Balearic Islands is far away from the limits indicated by FRASER-JENKINS (1982) and does not allow us to recognize more than one taxon. Therefore, only *D. pallida* subsp. *balearica* would be present in the Balearic flora. Other taxonomic comments about *D. pallida* subsp. *balearica* can be found in NARDI (1976) and VIANE (1990).

³⁹The discovery of this species in Mallorca was reported by ROSSELLÓ & al. (1989).

⁴⁰See ROSSELLÓ & ALOMAR (1987) for details on its presence in Mallorca.

⁴¹A single plant with meiotic irregularities and growing in the neighbouring of *P. aculeatum* and *P. setiferum* has been referred to this hybrid (CUBAS & ROSSELLÓ, unpubl. data).

⁴²The taxonomy of the infraspecific taxa of *J. phoenicea* has not been yet adequately solved. The subspecies *phoenicea*, *turbinata* (Guss.) Nyman and *eumediterranea* Lebreton & Thivend are reported in the Balearic flora (LEBRETON & THIVEND, 1981; GREUTER & al., 1984; AMARAL FRANCO, 1986). We have not found any clear morphological discontinuity among the Balearic populations, and in our opinion, only one polymorphic taxon of *J. phoenicea* exists in the Balearic Islands.

⁴³*E. fragilis* is not present in Eivissa and Formentera. The unlocated record from Eivissa of BARCELO (1879-1881), later expanded to Formentera (BOLOS & VIGO, 1984) has never been confirmed.

⁴⁴*P. ceciliae* A. Llorens & L. Llorens was described from specimens of southern Mallorca (LLORENS & LLORENS, 1972; LLORENS, 1984) and its distribution was later expanded to Menorca and Eivissa (LLORENS, 1979). The characters given by LLORENS & LLORENS (1972) to differentiate *P. ceciliae* from *P. halepensis* are poor and intermediate forms between them are known. Also, most seeds sown from wild *P. ceciliae* cones develop as normal *P. halepensis* (J. L. Gradaille, pers. com.). Populations of *P. ceciliae* are always small and they are found mixed with specimens of *P. halepensis*. From the above statements

we think that *P. ceciliae* is only a local mutation of *P. halepensis*, however, other possibilities cannot be, at present, excluded. Degenerating genetic forms of other pine species from islands of eastern Mediterranean have also been documented (PANETSOS, 1989). If the *P. ceciliae* specimens are to be considered as a different taxon, we think that a varietal rank would be the most adequate. We propose the following nomenclatural change:

Pinus halepensis* Miller var. *ceciliae* (A. Llorens & L. Llorens) L. Llorens, **comb. nov.*

Basionym: *Pinus ceciliae* A. Llorens & L. Llorens, Bol. Soc. Hist. Nat. Bal. 17: 51 (1972); Folia Bot. Misc. 4: 55 (1984).

Synonym: *P. halepensis* var. *ceciliae* (A. Llorens & L. Llorens) L. Llorens, Mediterranea 3: 116 (1979), **comb. inval.**

⁴⁵Contrary to the statements of BONAFE (1980) and GREUTER & al. (1984), *P. pinea* is not native in the Balearic flora. All the populations are naturalized.

⁴⁶Distribution given in ALOMAR (1982).

REFERENCES

- ALOMAR, G. (1982). Distribució del Teix (*Taxus baccata* L., 1753) a Mallorca. *Est. Bal.* 7: 37-51.
- ALOMAR, G., J. RITA & J. A. ROSSELLÓ (1991). Notas florísticas de las Islas Baleares (3). *Boll. Soc. Nat. Hist. Nat. Balears* 31: 141-144.
- AMARAL FRANCO, J. (1986). Juniperus L. In: CASTROVIEJO, S. & al. (Eds.), *Flora Ibérica* 1: 181-188.
- BALLESTEROS, E. (1989). Isoetes histrix Bory a Mallorca i Menorca. *Buttl. Inst. Cat. Hist. Nat.* 57: 107.
- BARCELO, F. (1867). Apuntes para una flora de las islas Baleares o Catálogo metódico de las plantas observadas en esta región que no se hallan mencionadas en el Enumeratio de Cambessedes. *Rev. Progr. Ciencias* 17(5): 294-317; 17(6): 345-369.
- BARCELO, F. (1879-1881). *Flora de las Islas Baleares*. Impr. J. Gelabert, Palma de Mallorca. 654 pp.
- BENNERT, W. & D. E. MEYER (1972). Der Gattungsbastard \times Asplenoceterach barrancense hybr. nov. (*Asplenium majoricum* Lit. \times *Ceterach officinarum* Lam. et DC.). *Willdenowia* 6: 461-470.
- BENNERT, W. & D. E. MEYER (1974). Ein neuer *Asplenium*-Bastard von Mallorca, *Asplenium* \times *litardierei* hybr. nov. *Ber. Deutsch. Bot. Ges.* 87: 21-28.
- BENNERT, W., R. E. G. PICHI SERMOLLI, H. RASBACH, K. RASBACH & T. REICHSTEIN (1988). *Asplenium* \times *helii* Lusina the valid name for the hybrid between *A. petrarchae* (Guérin) DC. and *A. trichomanes* L. (*Aspleniaceae*, *Pteridophyta*) I. Nomenclatural notes. *Bauhinia* 9: 103-106.
- BENNERT, W., R. E. G. PICHI SERMOLLI, H. RASBACH, K. RASBACH & T. REICHSTEIN (1989). *Asplenium* \times *helii* Lusina the valid name for the hybrid between *A. petrarchae* (Guérin) DC. and *A. trichomanes* L. (*Aspleniaceae*, *Pteridophyta*) II. Detailed description and illustrations. *Webbia* 43: 311-337.
- BENNERT, W., H. RASBACH & K. RASBACH (1987). *Asplenium* \times *reichstenii* (= *Asplenium fontanum* \times *A. majoricum*; *Aspleniaceae*, *Pteridophyta*), a new endemic hybrid from Mallorca, Balearic Islands. *Fern Gaz.* 13: 133-141.
- BENNERT, W., H. RASBACH & K. RASBACH (1990). *Asplenium petrarchae* (Guérin) DC. subsp. *bivalens* und *Asplenium* \times *helii* nothosubsp. *calabraense* — Neufunde auf der Insel Mallorca. *Farnblätter* 21: 15-26.
- BENNERT, W., H. RASBACH, K. RASBACH & T. REICHSTEIN (1988). *Asplenium* \times *rosselloi* (= *A. balearicum* \times *A. onopteris*; *Aspleniaceae*), a new fern hybrid from Menorca, Balearic Islands. *Willdenowia* 17: 181-192.
- BIANOR, F. (1917). Plantes de Mallorca. *Buttl. Inst. Cat. Hist. Nat.* 17: 133-152.
- BOLOS, O. & J. VIGO (1984-1990). *Flora dels Països Catalans*. Vol. 1 (1984: 736 pp.). Vol. 2 (1990: 921 pp.). Ed. Barcino, Barcelona.
- BONAFE, F. (1977-1980). *Flora de Mallorca*. Vol. 1 (1977: 363 pp.). Vol. 2 (1978: 378 pp.). Vol. 3 (1979: 380 pp.). Vol. 4 (1980: 449 pp.). Ed. Moll, Palma de Mallorca.
- CAMBESSEDES, J. (1827). Enumeratio plantarum, quas in insulis Balearibus collegit J. Cambessedes, earumque circa Mare Mediterraneum distributio geographica. *Mém. Mus. Hist. Nat. Paris* 14: 173-335.
- CASTROVIEJO, S., M. LAINZ, G. LOPEZ GONZALEZ, P. MONTSERRAT, F. MUÑOZ GARMENDIA, J. PAIVA & L. VILLAR (Eds.) (1986-1990). *Flora Ibérica. Plantas vasculares de la Península Ibérica e Islas Baleares*. Vol. 1 (1986: Lycopodiaceae-Papaveraceae, liv + 575 pp.). Vol. 2 (1990: Platanaceae-Plumbaginaceae (partim), lii + 897 pp. Real Jardín Botánico, CSIC, Madrid.
- CRONQUIST, A. (1988). *The evolution and classification of flowering plants*. New York Botanical Garden. 555 pp.
- CUBAS, P., E. PANGUA & J. A. ROSSELLÓ (1987). Un nuevo híbrido balear del género *Asplenium*. *Anales Jard. Bot. Madrid* 44: 534-536.
- CUBAS, P., J. A. ROSSELLÓ & E. PANGUA (1988). Comparative study of *Asplenium balearicum*, *A. onopteris* and their spontaneous hybrid, *A.* \times *tyrrhenicum*. *Anales Jard. Bot. Madrid* 45: 75-92.
- CUBAS, P., J. A. ROSSELLÓ & E. PANGUA (1989). A new triploid hybrid in the *Asplenium trichomanes* complex: *Asplenium trichomanes* nothosubsp. *lucanum* (*A. trichomanes* subsp. *inexpectans* \times *A. trichomanes* subsp. *quadri-valens*) nothosubsp. *nova*. *Candollea* 44: 181-190.
- DERRICK, L. N., A. C. JERMY & A. M. PAUL (1987). Check-list of European Pteridophytes. *Sommerfeltia* 6: 1-94.
- DUVIGNEAUD, J. (1974). Excursion du 21 au 28 juin 1974 à Majorque. Catalogue provisoire de la Flore des Baléares. *Dép. Bot. Univ. Liège*: 44 pp.

- DUVIGNEAUD, J. (1979). Catalogue provisoire de la flore des Baléares. 2^{me} éd. *Soc. Ech. Pl. Vasc. Eur. Occ. et Bass. Médit.* 17 (suppl.): 43 pp.
- FONT QUER, P. (1919). Pteridófitas de las Pitiusas. *Bol. R. Soc. Esp. Hist. Nat.* 19: 505-511.
- FRASER-JENKINS, C. (1982). Dryopteris in Spain, Portugal and Macaronesia. *Bol. Soc. Brot.*, ser. 2, 55: 175-336.
- GANDOGGER, M. (1891). *Flora Europae terrarumque adjacentium*. Vol. 27: 314 pp.
- GARCÍAS FONT, L. (1919). Contribució a la Flora Balear V. Plantes dels voltants d'Artá i Capdepera. *Bull. Inst. Cat. Hist. Nat.* 1919: 110-117.
- GREUTER, W., H. M. BURDET & G. LONG (1984). *Med-Checklist 1: Pteridophyta (ed. 2). Gymnospermae. Dicotyledones (Acanthaceae-Cneoraceae)*. Ed. Conserv. Jard. Bot. Genève, 330 pp. + 100 (appendix).
- GREUTER, W., H. M. BURDET & G. LONG (1986). *Med-Checklist 3: Dicotyledones (Convolvulaceae-Labiatae)*. Ed. Conserv. Jard. Bot. Genève, 395 pp. + 129 (appendix).
- GREUTER, W., H. M. BURDET & G. LONG (1989). *Med-Checklist 4: Dicotyledones (Lauraceae-Rhamnaceae)*. Ed. Conserv. Jard. Bot. Genève, 458 pp. + 129 (appendix).
- JALAS, J. & J. SUOMINEN (1972). *Atlas Florae Europaeae*. Vol. 1. Helsinki.
- JAQUOTOT, M. C. & J. ORELL (1968). *Asplenium majoricum* R. Litardièrre, su área de expansión en la Sierra Norte de Mallorca. *Collect. Bot.* 7: 559-571.
- KNOCHÉ, H. (1921-1922). *Flora Balearica. Etude phytogéographique sur les îles Baléares*. Vols. 1-4. Ed. Imp. Roumégous & Déhen, Montpellier.
- LEBRETON, P. & S. THIVEND (1981). Sur une sous-espèce du Genévrier de Phénicie (*Juniperus phoenicea* L.), définie à partir de critères biochimiques. *Naturalia Monspel., Sér. Bot.* 52: 125-134.
- LLORENS, A. & L. LLORENS (1972). Contribució al estudio de la flora balear. *Bol. Soc. Hist. Nat. Bal.* 17: 51-54.
- LLORENS, L. (1979). Nueva contribució al conocimiento de la flora balear. *Mediterranea* 3: 101-122.
- LLORENS, L. (1984). Notas florísticas baleáricas. *Fol. Bot. Misc.* 4: 55-58.
- LOVIS, J., H. RASBACH & T. REICHSTEIN (1989). *Asplenium trichomanes* L. nothosubsp. *melzeri* nothosubsp. nov. The triploid hybrid between *A. trichomanes* subsp. *inexpectans* and subsp. *quadrivalens*. *Candollea* 44: 543-553.
- LOVIS, J. & T. REICHSTEIN (1970). *Asplenium* × *orellii* hybr. nov. = *A. majoricum* Litard. × *A. trichomanes* L. subsp. *quadrivalens* D. E. Meyer. *Ber. Schweiz. Bot. Ges.* 79: 335-345.
- LOVIS, J., A. SLEEP & T. REICHSTEIN (1970). Der Farnbastard *Asplenium* × *sollerense* hybr. nov. = *A. majoricum* Litard. × *A. petrarchae* (Guérin) DC. subsp. *petrarchae*. *Ber. Schweiz. Bot. Ges.* 79: 369-376.
- MAHEU, J. (1912). Exploration et flore souterraine des cavernes de Catalogne et des Iles Baléares. *Spelunca* 8: 361-465.
- MARES, P. & G. VIGINEIX (1880). Catalogue raisonné des plantes vasculaires des Iles Baléares. Lib. G. Masson, Paris. 418 pp.
- MONTERRAT, P. (1974). Pteridófitas del herbario JACA. *Anal. Inst. Bot. A. J. Cavanilles* 31: 55-70.
- MORALES, M. J. & J. FERNÁNDEZ CASAS (1989). Mapa 290. In: FERNÁNDEZ CASAS, J. (Ed.), Asientos para un atlas corológico de la flora occidental, 14. *Fontqueria* 25: 1-201.
- MUS, M., J. A. ROSSELLÓ & N. TORRES (1990). De flora balearica adnotaciones (6-8). *Candollea* 45: 75-80.
- NARDI, E. (1976). La distribuzione italiana di "Dryopteris pallida" (Bory) Fomin. *Webbia* 30: 3-32.
- ORMONDE, J. (1986). *Blechnum* L. In: CASTROVIEJO, S. & al. (Eds.), *Flora Iberica*. Vol. 1: 151-153. Real Jardín Botánico, CSIC, Madrid.
- PALAU, P. (1954). Nuevas estirpes para la flora de Baleares (2^a serie). *Anal. Inst. Bot. A. J. Cavanilles* 12: 293-303.
- PANETSOS, K. P. (1989). Degradación genética del pinar de la isla de Lesbos. In: *Degradación de zonas áridas en el entorno mediterráneo. Monografías de la Dirección General del Medio Ambiente*. MOPU.
- PICHI SERMOLLI, R. E. G. (1988). *Asplenium balearicum* Shivas in Sardegna. *Bull. Soc. Sarda Sci. Nat.* 24: 149-162.
- PRADA, C. (1983). El género *Isoetes* en la Península Ibérica. *Acta Bot. Malacitana* 8: 73-100.
- PRADA, C. (1986). *Isoetes* L. In: CASTROVIEJO, S. & al. (Eds.), *Flora Iberica*. Vol. 1: 15-20. Real Jardín Botánico, CSIC, Madrid.
- RASBACH, H., K. RASBACH & H. W. BENNERT (1987). × *Asplenoceterach barrancense* Bennert & Meyer (*Aspleniaceae*, Pteridophyta) — Neufunde und zytologische Untersuchungen. *Farnblätter* 17: 3-16.
- RASBACH, H., K. RASBACH, T. REICHSTEIN & H. W. BENNERT (1990). *Asplenium trichomanes* subsp. *coriaceifolium*, a new subspecies and two new intraspecific hybrids of the *A. trichomanes* complex (*Aspleniaceae*, Pteridophyta). I. Nomenclature and typification. *Willdenowia* 19: 471-474.
- RITA, J. (1987). *Pilularia minuta* Durieu (*Marsileaceae*) en las islas Baleares. *Acta Bot. Malacitana* 12: 249-252.
- RÓDRIGUEZ, J. J. (1874). Suplemento al Catálogo razonado de las plantas vasculares de Menorca. *Anal. Soc. Esp. Hist. Nat.* 3: 5-68.
- RÓDRIGUEZ, J. J. (1904). *Flórlula de Menorca*. Imp. Fàbregues, Mahón. 198 pp.
- ROSSELLÓ, J. A. (1989). *La Pteridoflora de las Islas Baleares*. Unpubl. Ph. D. University of the Balearic Islands.
- ROSSELLÓ, J. A. & G. ALOMAR (1987). Dos pteridófitos nuevos para la flora balear. *Acta Bot. Malacitana* 12: 256.
- ROSSELLÓ, J. A., P. CUBAS, J. L. GRADAILLE & B. SASTRE (1990). *Asplenium* × *artanense* (*Aspleniaceae*: Pteridophyta) a new diploid hybrid from Mallorca, Spain. *Fern Gaz.* 13: 349-355.

- ROSSELLÓ, J. A., P. CUBAS & A. REBASSA (1991). Two new *Asplenium* taxa from the Balearic Islands: *A. azomanes* and *A. × tubalense* (= *A. azomanes* × *A. trichomanes* subsp. *quadri-valens*). *Rivasgodaya* 6: 115-128.
- ROSSELLÓ, J. A., J. L. GRADAILLE, B. SASTRE, J. VICENS & A. E. SALVO (1989). *Dryopteris tyrhena* Fraser-Jenkins & Reichstein en Baleares. *Acta Bot. Malacitana* 14: 255-258.
- ROSSELLÓ, J. A., J. PERICAS, G. ALOMAR & N. TORRES (1986). Atlas Pteridológico de las islas Baleares. *Acta Bot. Malacitana* 11: 294-302.
- ROSSELLÓ, J. A. & M. SERRA (1987). *Asplenium balearicum* Shivas en la isla de Menorca. *Acta Bot. Malacitana* 12: 95-102.
- SALVO, A. E., B. CABEZUDO & L. ESPAÑA (1984). Atlas de la pteridoflora Ibérica y Balear. *Acta Bot. Malacitana* 9: 105-128.
- SHEFFIELD, E., P. G. WOLF, C. H. HAUFLER & J. M. S. JUBRAEL (1986). *Pteridium herediae*: A dangerous diploid bracken? In: SMITH, R. T. & al. (Eds.), *Bracken, Ecology, land use and technology*: 301-307.
- SHEFFIELD, E., P. G. WOLF, C. H. HAUFLER, C. H. RANKER & A. C. JERMY (1989). A re-evaluation of plant referred to as *Pteridium herediae* (Colmeiro) Löve et Kjellqvist. *Bot. J. Linn. Soc.* 99: 377-386.
- SMYTHIES, B. E. (1984a). Flora of Spain and the Balearic Islands. Checklist of vascular plants I. Pteridophyta, Gymnospermae, Acanthaceae-Crassulaceae. *Englera* 3: 1-212.
- SMYTHIES, B. E. (1984b). Flora of Spain and Balearic Islands. Checklist of vascular plants II. Cruciferae-Rutaceae. *Englera* 3: 213-486.
- SMYTHIES, B. E. (1986). Flora of Spain and Balearic Islands. Checklist of vascular plants III. Salicaceae-Zygophyllaceae, Agavaceae-Zannichelliaceae. *Englera* 3: 487-880.
- TRABUT, L. (1917). *Scolopendrium hybridum* Milde à Majorque. *Bull. Soc. Hist. Nat. Afrique Nord* 8: 95-96.
- TUTIN, T. G. & al. (Eds.) (1964-1980). *Flora Europaea*. Vol. 1 (1964: 464 pp.). Vol. 2 (1968: 455 pp.). Vol. 3 (1972: 370 pp.). Vol. 4 (1976: 505 pp.). Vol. 5 (1980: 452 pp.). Cambridge Univ. Press, Cambridge.
- VIANE, R. (1990). Epidermology of european ferns. In: RITA, J. (Ed.), *Taxonomía, biogeografía y conservación de pteridófitos*: 69-89. Soc. Hist. Nat. Bal. & Inst. Men. Est., Palma de Mallorca.

Addresses of the authors: J. A. R.: Botánica, Dept. Biología Vegetal, Fac. Ciencias Biológica, Universidad de Valencia, E-46071 Burjassot, Spain.

P. C.: Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense, E-28040 Madrid, Spain.

N. T.: Apartado de Correos 64, E-07800 Eivissa, Islas Baleares, Spain.

