

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	48 (1993)
<b>Heft:</b>	1
<b>Artikel:</b>	Callicarpa moldenkeiana (Verbenaceae) : a new species from north-eastern India
<b>Autor:</b>	Rajendran, A. / Daniel, P.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-879641">https://doi.org/10.5169/seals-879641</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# **Callicarpa moldenkeiana (Verbenaceae), a new species from north-eastern India**

A. RAJENDRAN  
&  
P. DANIEL

## **ABSTRACT**

RAJENDRAN, A. & P. DANIEL (1993). *Callicarpa moldenkeiana* (Verbenaceae), a new species from northeastern India. *Candollea* 48: 111-113. In English, French and English abstracts.

A new species, *Callicarpa moldenkeiana* Rajendran & Daniel from northeastern India, is described and illustrated.

## **RÉSUMÉ**

RAJENDRAN, A. & P. DANIEL (1993). *Callicarpa moldenkeiana* (Verbenaceae), nouvelles espèce du nord-est de l'Inde. *Candollea* 48: 111-113. En anglais, résumés anglais et français.

Une nouvelle espèce, *Callicarpa moldenkeiana* Rajendran & Daniel du nord-est de l'Inde est décrite et illustrée.

**KEY-WOROS:** *Callicarpa moldenkeiana* — Meghalaya — India.

Specimens of the genus *Callicarpa* L. in Indian herbaria were studied for a revision of the Indian Verbenaceae. Some specimens from Meghalaya in northeastern India at ASSAM were found to have been wrongly determined as *Callicarpa arborea* Roxb. Though they apparently resembled *C. arborea*, a critical study proved that they belonged to a new species which is described here with an illustration. A comparison is made with the closely related *C. vestita*.

## **Callicarpa moldenkeiana Rajendran & Daniel, spec. nov. (Fig. 1).**

**Typus:** India: Meghalaya, Punglong, 4.11.1938, R. N. De 18877 (holo: ASSAM); **Paratypi:** Meghalaya, Khasia & Jaintia hills, Punglong forest, 30.8.1935, S. R. Sharma 12154; Meghalaya, Khasia & Jaintia hills, Jarain to Nongthala, 25 km from Jarain on Dawki road, 900 m, 15.11.1968, N. P. Balakrishnan 49852 (ASSAM).

*C. vestita* affinis sed habitus fruticoso; partibus junioribus stellato-tomentosis; foliis angustiore lanceolatis ad marginem intergris ad laeviter undulatis; inflorescentia duplo vel triplo ramosa; pedunculis petiolis curtioribus differt.

Differs from *C. vestita* in being a shrub, young parts stellate-tomentose, leaves lanceolate, acute or obtuse at base, entire to slightly undulate at margins, long-acuminate at apex, 10-23 × 4-6 cm,



Fig. 1. — *Callicarpa moldenkeiana* Rajendran & Daniel, spec. nov.  
A, habit; B, calyx; C, flower.

densely matted with stellate-tomentum beneath, petioles up to 4.5 cm long, inflorescence 2-3 times branched, peduncles 1.5-2.5 cm long, shorter than petioles, corolla lobes oblong and ovary oblong.

Shrub, 3-6 m high; branches and branchlets subterete; young parts densely stellate-tomentose, glabrous when mature; bark brownish grey; nodes annulate; internodes 2.5-10 cm long. Leaves lanceolate, acute or obtuse at base, entire to slightly undulate at margins, long-acuminate at apex, 10-23 × 4-6 cm, subcoriaceous, dark green, glabrous above, paler, densely matted, stellate-tomentose with minute rounded yellow glands beneath; lateral nerves 6-12 pairs, ascending, arcuate at margins, obscure above, distinct beneath; petioles cylindric, stout, 3-4.5 cm long, densely stellate-tomentose. Cymes axillary, 2-3 times dichotomously branched; peduncles slender, terete, shorter than petioles, 1.5-2.5 cm long, stellate-tomentose; bracts linear-lanceolate, ± 5 × 2 mm. Flowers numerous, purple. Calyx cupular, ± 1 × 1 mm, 4-toothed; teeth acute, stellate-tomentose outside. Corolla infundibular, 4-lobed; lobes oblong, obtuse, ± 1 × 0.5 mm, glabrous; tube narrow, ± 1.5 × 1 mm, glabrous. Stamens 4, inserted below mouth of corolla tube; filaments filiform, 1.5-2 mm long, glabrous; anthers oblong, ± 1 × 0.1 mm, 2-celled, divaricate, basifixied. Ovary oblong, ± 1 × 0.3 mm, glabrous; style slender, ± 4 mm long; stigma capitate. Drupes globose, ± 2 × 2 mm, purple black when mature; fruiting-calyx platelliform, ± 0.2 × 1 mm, membranous, pubescent.

*Flowering and fruiting.* — August-November.

*Habitat.* — In secondary forests, mostly on hill slopes of evergreen forests, at about 900 m.

*Distribution.* — Meghalaya, Punglong forests in Khasia and Jaintia hills.

*Note.* — This species is named after Dr. Harold N. Moldenke, a lifetime student of the Verbenaceae who has contributed enormously to the family.

#### ACKNOWLEDGMENTS

We are thankful to Dr. N. P. Balakrishnan, Joint Director, Botanical Survey of India, Coimbatore, for facilities, and to Dr. V. J. Nair, Scientist-SD, Botanical Survey of India, Shillong, for the Latin diagnosis. A. R. is thankful to the Director, Botanical Survey of India, for a research fellowship. He is also thankful to the officer-in-charge of ASSAM for permission to study the specimens there.

