

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	48 (1993)
<b>Heft:</b>	2
<b>Artikel:</b>	Clerodendrum nicolsonii (Verbenaceae) : a new species from north-eastern India
<b>Autor:</b>	Rajendran, A. / Daniel, P.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-879663">https://doi.org/10.5169/seals-879663</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Clerodendrum nicolsonii (Verbenaceae), a new species from north-eastern India

A. RAJENDRAN  
&  
P. DANIEL

## RÉSUMÉ

RAJENDRAN, A. & P. DANIEL (1993). *Clerodendrum nicolsonii* (Verbenacée), une nouvelle espèce du nord-est de l'Inde. *Candollea* 48: 347-349. En anglais, résumés français et anglais.

*Clerodendrum nicolsonii*, espèce nouvelle du nord-est de l'Inde est décrite et illustrée.

## ABSTRACT

RAJENDRAN, A. & P. DANIEL (1993). *Clerodendrum nicolsonii* (Verbenaceae), a new species from northeastern India. *Candollea* 48: 347-349. In English, French and English abstracts.

A new species, *Clerodendrum nicolsonii* from northeastern India, is described and illustrated.

**KEY-WORDS:** *Clerodendrum nicolsonii* — VERBENACEAE — Deomukh — Assam — India.

While studying the genus *Clerodendrum* L. for a revision of the genus in India as a part of a revision of the Indian Verbenaceae, a few collections from Assam were found to belong to an undescribed species.

## **Clerodendrum nicolsonii** Rajendran & Daniel, spec. nov. (Fig. 1).

*C. wallichii* Merr. affinis foliis decussatis, obovatis vel oblanceolate-ellipticis ad margineum serratis; petiolis usque ad 5 cm longis; bracteis oblanceolatis, ± 3.5 × 1 cm, ad margineum serratis, ad apicem caudate; inflorescentis paniculatis, erectis; calycibus pubescentibus, noninflatis; corollis laeviter 5-lobatis, lobis obovatis; calycibus fructiferis viridibus, fructibus in calyces inclientibus differt.

**Typus:** India, Assam, North Lakhimpur, Deomukh Reserve Forest, 16.5.1966, D. M. Verma 46361 (holo: ASSAM); paratypi: India, Assam, North Lakhimpur, 23.5.1966, D. M. Verma 46654 (ASSAM); Assam, Deomukh, 13.11.1978, G. D. Pal 70238 (ARUN).

Differs from *C. wallichii* in leaves being obovate or oblanceolate-elliptic, petioles up to 5 cm long, inflorescence erect panicles, composed of 6-8 branched cymes, 8-15 cm long, pubescent, bracts oblanceolate, serrate, caudate, ± 3.5 cm long, calyx not inflated, teeth lanceolate, pubescent outside, corolla indistinctly 5-lobed, lobes obovate, entire, tube narrow, cylindric, ± 2.5 cm long, drupes globose, less than 1 cm across, fruiting-calyx green, pubescent and enclosing drupe.




Fig. 1. — *Clerodendrum nicolsonii* Rajendran & Daniel, spec. nov.  
A, habit; B, flower; C, fruit.

Shrub, 3-3.5 m high, pubescent when young, nearly glabrous when mature, acutely 4-angular, slightly sulcate between angles; nodes annulate with a band of hirsute pubescence. Leaves obovate or oblanceolate-elliptic, obtuse or acute to cuneate at base, faintly and irregularly serrate at margins, abruptly acuminate at apex, 4-20 × 1.5-7 cm, chartaceous, dark green above, paler beneath, glabrous; lateral nerves 6-8 pairs, ascending, arcuate at margins, distinct beneath; petioles slender, canaliculate, 0.8-5 cm long. Panicles terminal, erect, composed of 6-8 decussate-opposite branched cymes, 8-15 × 6-13 cm; peduncles acutely 4-angular, sulcate between angles, 2.5-3.5 cm long, pubescent; bracts oblanceolate, serrate at margins, caudate at apex, ± 3.5 × 1.5 cm; bractlets subulate, ± 6 mm long. Flowers numerous; pedicels slender, 0.5-1 mm long, pubescent. Calyx ± 5 × 1 mm, 5-toothed; teeth lanceolate with a prominent midrib, pubescent outside. Corolla hypocrateriform, white, 5-lobed; lobes subequal, obovate, obtuse, ± 5 × 3 mm, glabrous; tube narrow, cylindric, ± 2.5 × 0.1 cm, glabrous outside. Stamens 4, didynamous, attached below the mouth of corolla tube; filaments filiform, 4-5 cm long, exserted; anthers greyish black. Ovary globose, ± 1 mm; style slender, equaling filaments; stigma 2-lobed; lobes equal, subulate. Drupes globose, 5-6 mm, black, seated on persistent accrescent calyx; fruiting-calyx distinctly 4-lobed; lobes lanceolate.

*Flowering and fruiting.* — May-November.

*Distribution.* — Known to occur only in Deomukh Reserve Forest in Assam, India.

*Note.* — This new species is named after Dr. Dan H. Nicolson, Smithsonian Institution, Washington, DC., whose generosity with literature and expertise on nomenclature have been of immense help in our revision of the Indian Verbenaceae.

#### ACKNOWLEDGEMENTS

We are thankful to Dr. N. P. Balakrishnan, Joint Director, Botanical Survey of India, Coimbatore, for encouragement and facilities, and to Dr. V. J. Nair, Indian Liaison Officer, Kew, for the Latin diagnosis. A. R. is thankful to the officers-in-charge of ARUN, ASSAM, CAL, NEHU and SHC for permission to study the specimens there.

