

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 48 (1993)

Heft: 2

Artikel: What is the Gran Chaco vegetation in South America? : II. A redefinition ; contribution to the study of the flora and vegetation of the Chaco, VII

Autor: Prado, Darién E.

DOI: <https://doi.org/10.5169/seals-879682>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

What is the Gran Chaco vegetation in South America? II. A redefinition

Contribution to the study of the flora and vegetation of the Chaco.

VII

DARIÉN E. PRADO

ABSTRACT

PRADO, D. E. (1993). What is the Gran Chaco vegetation in South America? II. A redefinition. Contribution to the study of the flora and vegetation of the Chaco. VII. *Candollea* 48: 615-629. In English, English and Spanish abstracts.

The twenty-three more relevant woody communities of the Gran Chaco are studied following classical phytosociological and numerical analysis techniques (PCA and CLINK), in an attempt to redefine the present concept of the Chaco province in South America. The congruence between the different results is striking. On this basis, the communities Gallery Forest, "Selva de Ribera", "Tipa-Pacará" Forest, "Palo blanco" Forest and the Calcareous Forest of Mato Grosso do Sul cannot be regarded as chaquenian, while the Austro-Brazilian Transitional Forest and three different communities from Mato Grosso are considered transitional. A new map of the Chaco *sensu stricto* is presented, with a wide belt of transitional vegetation on the east of the region, whereas gallery forests and hills vegetation are completely excluded.

RESUMEN

PRADO, D. E. (1993). ¿Qué es la vegetación del Gran Chaco de Sudamérica? II. Redefinición. Contribución al estudio de la flora y de la vegetación del Chaco. VII. *Candollea* 48: 615-629. En español, resúmenes en inglés y en español.

Las veintitrés comunidades leñosas más relevantes del Gran Chaco fueron estudiadas siguiendo técnicas de análisis fitosociológico clásico y numérico (PCA y CLINK), tratando de redefinir el concepto actual de la provincia del Chaco en Sud América. La congruencia entre los distintos resultados es sorprendente. Sobre esta base, las comunidades Bosque en Galería, Selva de Ribera, Bosque de Tipa y Pacará, Bosque de Palo Blanco y los Bosques Calcáreos de Mato Grosso do Sul no pueden ser aceptadas como chaqueñas, mientras que el bosque Transicional Austro-Brasileño y tres comunidades diferentes de Mato Grosso son consideradas transicionales. Se presenta un nuevo mapa del Chaco *sensu stricto*, que muestra una amplia faja de vegetación transicional en el este de la región, al tiempo que los bosques en galería y la vegetación de los cerros son totalmente excluidos.

KEY-WORDS: Chaco — Classical phytosociology — Floristics groups — Numerical analysis — Vegetation — Woody communities.

Introduction

In a previous contribution a review of the available knowledge on the Gran Chaco vegetation of Bolivia, Paraguay and Argentina was presented (PRADO, 1993). In studying this problem the need of a thorough analysis of the present day notion of the Chaco arose, since in the literature available the concept of the Chaco as a phytogeographical province has simply been equated with the Chaco as a geographical region (see review in PRADO, 1993), an established view that has prevailed unchallenged so far. The comparative study showed that the geographical region known as

the "Gran Chaco" in fact includes rather different kinds of vegetation, and there is a number of floristic elements of widely different lineages in the woody communities still regarded as chaquenian. Such studies then provide the basis for an attempt to redefine the "Chaco proper" by more objective techniques.

The twenty-three woody vegetation units encompassed in the analysis have been regarded as pertaining to the Gran Chaco by different authors. These include the Gallery Forests in the islands of the river Paraná and tributaries, considered chaquenian by RAGONESE (1941) and RAGONESE & CASTIGLIONI (1970), and often confused with the next one, the "Selva de Ribera" (SCHULZ, 1967; PRADO & al., 1989), as in REBORATTI & NEIFF (1986). The Subandean Piedmont Forests of NW Argentina and SW Bolivia (PRADO, 1991), bordering the western limit of the Chaco and included in it by HUECK (1959, 1972) and UNZUETA (1975), consists of two different units: the "Tipa-Pacar " and the "Palo blanco" forests. The Austro-Brazilian Transitional Forest (PRADO, 1991) and the nearby "quebrachales" of *Schinopsis balansae* have, more often than not, been confused as a single vegetation type (e.g. HUECK, 1972; CABRERA, 1976). Also arguable is the phytogeographical position of the five communities described by PRADO & al. (1992) for a sector of alleged Chaco vegetation in Mato Grosso do Sul (HUECK, 1955). There is little argument, in contrast, about the chaquenian character of the "quebrachales" of three "quebrachos" dominated by *Schinopsis quebracho-colorado*, of white "quebracho" (*Aspidosperma quebracho-blanco*), and of two "quebrachos" in the drier western Chaco; the same is true for the "Palosantales" of *Bulnesia sarmientoi*, the white "quebracho" dominated Arid Chaco Woodland of SW Chaco (SARMIENTO, 1972), together with the main four azonal woody communities ("Algarrobales", "Cardonales", two variants of "Palmares", and "Vinalares"; PRADO, 1993). Also enclosed in the analysis are the Pampean and Subandean varieties of the Sierra Chaco Forests.

Methods

In order to assess to what an extent the woody chaquenian communities and neighbouring seasonal formations are interrelated, reliable floristic lists were selected from the available literature and worked out in phytosociological tables on a basis of presence-absence of species. These data were then analyzed following three different techniques:

a) Classic phytosociological analysis of the Z rich-Montpellier school, in the more modern version modified by MUELLER-DOMBOIS & ELLENBERG (1974). This consists essentially in listing all the species (rows) occurring in certain areas or localities (columns), and then mechanically search for species with common patterns of distribution in the columns. Subsequent reshuffling of columns and rows generally results in the grouping of some species which seem to be exclusive to a determined group of localities, exclusive to a single area, or common to most or all of them. These species groups are taken to indicate the presence of common environmental factors by which they are restricted in their phytogeographical distribution. Therefore, such groups have an ecological indicator value, and they are referred to as "floristic groups (FG)" (MUELLER-DOMBOIS & ELLENBERG, 1974). Because of the subjective nature of this analyses it is necessary to compare the results with those of more objective statistical studies, as the following.

b) Multivariate numerical methods:

b.1. CLINK (Complete Linkage) algorithm from WISHART's (1987) package, employing S RENSEN's (1948) similarity index (also known as Czekanowski-Dice coefficient). This agglomerative technique of classification fuses the individuals (localities/areas) into increasingly larger discrete groups, based on their similarity matrix, and a dendrogram is provided to show their relationships. The fusion is interrupted according to a subjective criterion, generally when recognizable clusters of ecologically-floristically related individuals are formed, or when no individual is left isolated (MATEUCCI & COLMA, 1982).

b.2. PCA (Principal Components Analysis) from the JMP INTM statistical software. This ordination technique, contrary to the classificatory ones, does not establish discrete classes but

displays the individuals under study in a hyper-dimensional space along axes of continuous variation. The axes (“principal components”) are numbered according to the decreasing percentage of accumulated variation they concentrate. Thus, Axis I always comprises the highest value of variation absorbed by any possible axis, which can be interpreted as representing one particular environmental or ecological factor. The individuals are then displayed in successive two-dimensional plots (“scattergrams”), of which only the Axis I-Axis II coordinates are shown and discussed here.




Fig. 1. — Dendrogram resulting from Clink classification analysis of the main woody communities of the Chaco and neighbouring formations. Letters at the bottom indicate the communities studied (see Table 1 for key), and figures in the horizontal lines the similarity index value at each level of fusion.

Results

All the vegetation types described by PRADO (1993) and usually regarded as part of the Gran Chaco region have been analyzed in Table 1. A number of FG, namely II, V, VI and XII amongst others, indicate very close links between those woody communities of wetter and less frost-affected areas, such as A (Gallery Forest), B (“Selva de Ribera”), C (“Tipa-Pacará” Forest), D (“Palo blanco” Forest) and the Austro-Brazilian Transitional Forest (E), alongside the communities described by PRADO & al. (1992) for Mato Grosso do Sul (F, G, H, I and J). Apart from FG XXI, which comprises species with a wide ecological plasticity, there are scarce connections between communities A to J and what will be here regarded as Chaco s.s. (communities K to W). The FG XXII comprises the species that relate the Austro-Brazilian Transitional Forest (E) to the Chaco proper, which are on the one hand of scarce importance here and sometimes accidental, and on the other




Fig. 2. — Scattergram from PCA analysis of the main woody communities of the Chaco and neighbouring formations.

are outweighed by the much more relevant links to communities A to D. The Chaco forest at Porto Murtinho (J) is the only one between the communities of Mato Grosso do Sul that can be regarded as truly chaquenan, as shown by Table 1 and coinciding with PRADO & al. (1992). The FG XXIII, XXIV, XXV, XXVI, XXVII, XXVIII, XXXIII and XXXIV illustrate the strong internal homogeneity of the Chaco s.s. woody communities.

Fig. 1 shows the dendrogram resulting from the CLINK classification analysis applied to the same set of communities shown in Table 1. It is noteworthy that the order of the communities rendered by CLINK is basically very similar to that proposed in the phytosociological table, save for the position of the Chaco at Porto Murtinho (J). Group α comprises communities A to E, while β those of Mato Grosso do Sul (F to I). Meanwhile, groups γ and δ consist of all the communities of the Chaco s.s., separated into two groups probably because of an east-west floristic gradient within the province.

The results of the PCA analysis shown in Fig. 2 present three basic clusterings of communities: the wetter ones (A to F) are together in cluster α' , the Mato Grosso do Sul communities are grouped in another one (β'), while the bulk of the Chaco s.s. in cluster γ' shows stronger links within itself than with the rest of the neighbouring formations. Once again the Chaco of Porto Murtinho (J) is in a transitional position and could pertain to either of two clusters, while the Austro-Brazilian Transitional Forest (E) unmistakably joins the group of wetter communities.

Discussion

The results of three different types of analysis applied to the Gran Chaco vegetation have been presented: the classical phytosociological approach and, as a more objective yardstick to evaluate this, two numerical analyses were performed, one to classify (CLINK) and one to ordinate (PCA) the samples. The congruence between the different results is striking. Out of the 23 vegetation units that had been regarded as chaquenan by different authors (see PRADO, 1993 and Table 1), five

Table 1. — Classical phytosociological analysis of the main woody communities of the Chaco and neighbouring formations. Source of the floristic lists is indicated.

Column: **A**, Gallery forest, Paraná River (FRANCESCHI & LEWIS, 1979). **B**, "Selva de Ribera" (PRADO & al., 1989; MORELLO & ADÁMOLI, 1974). **C**, "Tipa-Pacará" forest (MEYER, 1963; DIGILIO & LEGNAME, 1966; BROWN & al., 1985). **D**, "Palo blanco" forest (MEYER, 1944; CORO, 1956; ADÁMOLI & al., 1972; CABRERA, 1976). **E**, Austro-Brazilian Transitional Forest (LEWIS & PIRE, 1981, and author's field experience; also MORELLO & al., 1971, MORELLO & ADÁMOLI, 1974, and CABRERA, 1976). **F**, Calcareous woodlands, Mato Grosso do Sul (PRADO & al., 1992). **G**, *Schinopsis balansae* parkland, M. Gr. do Sul (PRADO & al., 1992). **H**, *Aspidosperma-Mimosa* scrubland, M. Gr. do Sul (PRADO & al., 1992). **I**: *Diplokeleba-Tabebuia-Capparis* scrubland, M. Gr. do Sul (PRADO & al., 1992). **J**, Chaco forest, Pto Murtinho, M. Grosso do Sul (PRADO & al., 1992). **K**, "Quebrachal" of *Schinopsis balansae* (LEWIS & PIRE, 1981, and author's field experience). **L**, "Quebrachal" of 3 "quebrachos" (RAGONESE, 1941; LEWIS & PIRE, 1981, and author's field experience). **M**, "Quebrachal" of 2 "quebrachos" (MORELLO & ADÁMOLI, 1968; ADÁMOLI & al., 1972, and author's field experience). **N**, "Quebrachal" of white "quebracho" (MORELLO & al., 1971; LEWIS & PIRE, 1981). **O**, "Palosantal" of *Bulnesia sarmientoi* (MORELLO & al., 1971, and ADÁMOLI & al., 1972). **P**, "Algarrobal" of *Prosopis* spp. (MORELLO & al., 1971). **Q**, "Vinalar" of *Prosopis ruscifolia* (MORELLO & al., 1971). **R**, "Palmar" of *Copernicia australis*, Western Chaco (CABRERA, 1976). **S**, "Palmar" of *Copernicia australis*, Eastern Chaco (CABRERA, 1976; LEWIS & PIRE, 1981, and author's field experience). **T**, Arid Chaco woodland (SAYAGO, 1969). **U**, "Cardonal" of *Stetsonia coryne* (SAYAGO, 1969). **V**, North Sierra Chaco (MARLANGE, 1972). **W**, South Sierra Chaco (SAYAGO, 1969).

			A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W
			Gall	Sal	T	P	P	Aug	C	S	AS	DI	Tr	Ch											
1	Laur	Nectandra																							
2	Legm	Albizia																							
3	Myrt	Hexachlamis																							
4	Capp	Crateva																							
5	Apoc	Peschiera																							
6	Legm	Inga																							
7	Legm	Acacia																							
8	Ulm	Celtis																							
9	Euph	Croton																							
10	Flac	Banara																							
11	Arec	Arecastrum																							
12	Legp	Geoffroea																							
13	Anno	Rollinia																							
14	Poly	Ruprechtia																							
15	Legm	Enterolobium																							
16	Myrt	Eugenia																							
17	Ruta	Fagara																							
18	Comb	Terminalia																							
19	Myrs	Rapanea																							
20	Sapo	Pouteria																							
21	Legc	Holocalyx																							
22	Mora	Ficus																							
23	Sapi	Sapindus																							
24	Mora	Sorocea																							
25	Meli	Trichilia																							
26	Nyct	Pisonia																							
27	Cbl	Schaefferia																							
28	Pter	Acrostichum																							
29	Cecr	Cecropia																							
30	Ruta	Pilocarpus																							
31	Ster	Guazuma																							
32	Tili	Luehea																							
33	Legc	Peltoporum																							
34	Ulm	Phyllostylon																							
35	Ulm	Celtis																							
36	Bign	Tabebuia																							
37	Legm	Anadenanthera																							
38	Ahac	Astronium																							
39	Ruta	Fagara																							
40	Capp	Capparis																							
41	Sapi	Diplokeleba																							
42	Sora	Patagonula																							
43	Myrt	Myrcianthes																							
44	Sapi	Allophylus																							
45	Nyct	Pisonia																							
46	Legc	Gleditsia																							
47	Sapo	Chrysophyllum																							
48	Mora	Maclura																							
49	Legm	Pithecellobium																							
50	Arac	Pentapanax																							
51	Cari	Carica																							
52	Phyt	Phytolacca																							
53	Sola	Brunfelsia																							
54	Anac	Astronium																							
55	Legc	Pterogyne																							
56	Sapi	Cupania																							
57	Sapo	Chrysophyllum																							
58	Ulm	Trema																							
59	Bign	Jacaranda																							
60	Myrt	Myrcianthes																							
61	Bign	Tecoma																							
62	Piper	Piper																							
63	Tili	Hellicarpus																							

I
II
III
IV
V
VI
VII
VIII
IX

			A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W			
			Gall	Seli	T	Pad	Aub	Calc	Sbil	Asm	DiTn	ChP	Ch	O	X	2	2	Wn	P	as	Aln	Vin	Pmw	Pme	ACW	Card	HSC	SSC
54	LEGP	Tipuana	tipu					1	1																			
55	LEGP	Myroxylon	balsamum					1	1																			
56	LEGM	Parapirotadenia	excelsa					1	1																			
57	URTI	Urera	baccifera					1	1																			
58	BOMB	Pseudobombax	argentinum					1	1																			
59	MELI	Cedrela	odorata					1	1																			
70	LEGP	Lonchocarpus	lilloi					1	1																			
71	LEGM	Acacia	albicorticata					1	1																			
72	SAPI	Athyana	weimannifolia					1	1																			
73	LEGC	Senna	spectabilis v. sp					1	1																			
74	BORA	Cordia	trichotoma/alillo					1	1																			
75	AREC	Acrocomia	chonta					1	1																			
76	MELI	Trichilia	clauseni					1	1																			
77	ELAE	Mutingia	calabura					1	1																			
78	RUBI	Pogonopus	tubulosus					1	1																			
79	COMP	Vernonia	fulva					1	1																			
80	COMP	Cnicothamnus	lorentzii					1	1																			
81	BUDD	Buddleja	albomaculata					1	1																			
82	RUTA	Fagara	hemialis					1	1																			
83	BOMB	Celiba	speciosa					1	1																			
84	RUBI	Calycoconvallium	multiflorum					1	1																			
85	LEGP	Amburana	cearensis					1	1																			
86	EUPH	Cnidioscolus	vitifolius v. cnic					1	1																			
87	ANAC	Schinopsis	brasiliensis					1	1																			
88	STER	Helicteres	ihotskyana					1	1																			
89	EUPH	Sapum	sp					1	1																			
90	RUBI	Randia	armata					1	1																			
91	STER	Sterculia	striata					1	1																			
92	LEGP	Machaerium	nirtum					1	1																			
93	EUPH	Croton	sp l					1	1																			
94	RHAM	Ziziphus	oblongifolius					1	1																			
95	POLY	Coccoloba	sp					1	1																			
96	ERYT	Erythroxylum	sp					1	1																			
97	CACT	Cereus	uruguayanus					1	1																			
98	BORA	Cordia	glabrata					1	1																			
99	AREC	Acrocomia	totali					1	1																			
100	CELA	Maytenus	ilicifolia					1	1																			
101	RHAM	Rhamnidium	elaeocarpon					1	1																			
102	RUBI	Chomelia	obtusifolia					1	1																			
103	BOMB	Pseudobombax	sp					1	1																			
104	BIGN	Tabebuia	ochracea					1	1																			
105	RUBI	Tocovena	formosa					1	1																			
106	COMB	Terminalia	argentea					1	1																			
107	Unct	Pisonia ?	sp					1	1																			
108	LEGM	Mimosa	hexandra					1	1																			
109	LEGM	Mimosa	sp					1	1																			
110	LEGM	Prosopis	rubriflora					1	1																			
111	EUPH	Jatropha	ribifolia					1	1																			
112	LEGM	Prosopis	sp					1	1																			
113	NYCT	Bougainvillea	sp					1	1																			
114	LEGM	Goldmani	paraguensis					1	1																			
115	EUPH	Aporosaella	chacoensis					1	1																			
116	RHAM	Scutia	buxifolia					1	1																			
117	ACHA	Achatocarpus	praecox					1	1																			
118	SAPO	Sideroxylon	obtusifolium					1	1																			
119	SANT	Acanthosvris	falcata					1	1																			
120	MYRT	Myrcianthes	displatensis					1	1																			
121	BOMB	Celiba	chodatii					1	1																			
122	EUPH	Sapum	haematospermum					1	1																			
123	LEGP	Geoffroea	decorticans					1	1																			
124	APOC	Aspidosperma	quebracho-blanco					1	1																			
125	RHAM	Ziziphus	mistol					1	1																			
126	LEGM	Prosopis	nigra					1	1																			
127	ANAC	Schinus	fasciculatus					1	1																			
128	LEGM	Acacia	praecox					1	1																			
129	ANAC	Schinopsis	balansae					1	1																			
130	LEGC	Caesalpinia	paraguariensis					1	1																			
131	LEGC	Cercidium	praecox					1	1																			
132	BIGN	Tabebuia	nodosa					1	1																			
133	CAPP	Capparis	tweediana					1	1																			
134	CAPP	Capparis	speciosa					1	1																			
135	SANT	Jodina	rhombofolia					1	1																			
136	LEGM	Acacia	aroma					1	1																			
137	CAPP	Capparis	retusa					1	1																			
138	VERB	Aloisia	gratissima					1	1																			
139	SIMA	Castela	coccinea					1	1																			
140	CELA	Maytenus	vitis-idaea					1	1																			
141	ULMA	Celtis	spinosa					1	1																			
142	LEGM	Acacia	caven					1	1																			
143	LEGM	Prosopis	alba					1	1																			
144	ULMA	Celtis	pallida					1	1																			
145	SOLA	Grabowskia	duplicata					1	1																			
146	LEGC	Senna	apovilla					1	1																			
147	ZYGO	Porlieria	microphylla					1	1																			
148	SOLA	Cestrum	parqui					1	1																			
149	LEGM	Prosopis	affinis					1	1																			
150	AREC	Trithrinax	campestris					1	1																			
151	CACT	Opuntia	sp					1	1																			
152	CHEN	Holmbergia	spedei					1	1		</																	

are definitely non-chaquenian (Gallery Forest, "Selva de Ribera", "Tipa-Pacar  Forest, "Palo blanco" Forest, and Calcareous Forest), four are here considered transitional with neighbouring formations (the Austro-Brazilian Transitional Forest and three of the communities of Mato Grosso do Sul), and 14 correspond to what is here defined as Chaco *sensu stricto*: the four "Quebrachales", the "Palosantal", "Algarrobal", "Vinalar", "Cardonal", the two "Palmares", both Sierra Chaco variants, the Arid Chaco Woodland and the Chaco Forest at Porto Murtinho. A brief discussion on the excluded and transitional communities follows.

a) Excluded communities

Between the communities listed in Table 1, none of the Gallery Forest (A), "Selva de Ribera" (B), "Tipa-Pacar  Forest (C), "Palo blanco" Forest (D) and the Calcareous Forest of Mato Grosso do Sul (F) can be regarded as chaquenian in character. Climate, soils and water regime are entirely different from that which prevails in the Chaco s.s., and as a result their flora is essentially of Amazonian lineage (*sensu* CABRERA & WILLINK, 1980). All of these vegetation types are well defined, each characterized by a floristic group comprising the species which are exclusive to them within the scope of this study. Moreover, the numerical analyses (Figs. 1 & 2) have shown that they are more closely inter-related, and linked to what has been called "Austro-Brazilian Transitional Forest" (PRADO, 1991), than to the Chaco s.s. communities. Therefore, none of these five vegetation types should be included in the Chaco phytogeographical province, as the more modern treatments have indicated (e.g. CABRERA, 1976; PRADO, 1991).

b) Transitional communities

The four transitional units consist of three from the Mato Grosso do Sul area studied by PRADO & al. (1992): the *Schinopsis balansae* parkland (G), the *Aspidosperma-Mimosa* scrubland (H), and the *Diplokeleba-Tabebuia-Capparis* scrubland (I), and finally the "Austro-Brazilian Transitional Forest" (E). The first three communities contain a very low percentage of true Chaco species (PRADO & al., 1992), along with a number of cerrado and semi-deciduous forest elements, and none of the three can be fully ascribed to any of the major phytogeographical units which converge in the Pantanal, i.e. Cerrados, Amazonian, Paranense and Chaco provinces (AD MOLI, 1982). Furthermore, the numerical analyses (Fig. 2) show these communities in a clear transitional position between the Chaco s.s. and the wetter vegetation units.

The Austro-Brazilian Transitional Forest (E) is undoubtedly the climax community in the east of the Gran Chaco area (LEWIS & PIRE, 1981, *sub* "bosque chaque o"), i.e. the sector referred to as "Eastern Chaco" in all phytogeographical classifications (MORELLO & AD MOLI, 1968; RAGONESE & CASTIGLIONI, 1970; CABRERA, 1971 & 1976). Four true Chaco communities also manage to extend into this eastern area. These are the *Schinopsis balansae* "quebrachal" (K), *Prosopis* spp. "algarrobales" (P), *Copernicia australis* "palmares" (S), and *Stetsonia coryne* "cardonales" (see PRADO, 1993). They are evidently edaphic-dependent, and their presence in an area of over 900 mm yearly rainfall is allowed only by the physiological drought caused by the high salt content of the soil and consequent higher osmotic potential. Consequently, and contrary to the position sustained by RAGONESE & CASTIGLIONI (1970) and CABRERA (1976), the "quebrachal" of *Schinopsis balansae* should not be regarded as a climax community, since it rather thrives on non-climatogenic saline soils, with high clay level, periodical waterlogging, elevated Na⁺⁺ concentration and very strongly alkaline subsoils (ESPINO & al., 1983). Thus, true Chaco vegetation is found in a strip of some 100 km wide, parallel to the Paraguay and Paran  rivers, in an area where climatic, edaphic and geomorphological conditions are different from those in the center and west of the Chaco s.s. Presumably, the salinity factor is responsible for this outlying representation, since Chaco elements can survive in edaphic conditions which are hostile for the subtropical humid forests species of the Austro-Brazilian Transitional Forest. This hypothesis is exactly the opposite to HUECK's (1972), who postulated that chaquenian species would occur on higher, drier ground within the floodable, water-modelled Eastern Chaco. On the contrary, such drier areas are covered by a different kind of forest dominated by species such as *Patagonula*

americana, *Phytolacca dioica*, *Ruprechtia laxiflora*, *Gleditsia amorphoides*, *Tabebuia impetiginosa*, *Pisonia zapallo*, and *Enterolobium contortisiliquum*. All of these species occur elsewhere in subtropical humid forests in S Brazil, E Paraguay and N Argentina. Indeed, the species listed above are all widespread in South American forests, and extend to NE Brazil or humid Atlantic forests (KLEIN, 1967), or even to Central America and Mexico in similar ecosystems (e.g. *Tabebuia impetiginosa*, in GENTRY, 1979).

Most species of the Austro-Brazilian Transitional Forest are elements which are common to the five vegetation types here excluded altogether from the Chaco s.s., such as the species listed above together with *Diplokeleba floribunda*, *Astronium balansae*, *Chrysophyllum gonocarpum*, *Pithecellobium scalare* and *Brunfelsia australis* (see Table 1, FGs II, V and VI). Seven of the most typical Chaco species¹ may also occur in this transitional unit, but they are never dominant trees and seem to be at the extreme of their ecological range in this area. Hardly any of these chaquenian intruders can be found further east in wetter areas either in Argentina, Paraguay or Brazil, and it is likely that their existence in this unit could be due to anthropogenic alteration of the environment (overgrazing, selective felling), or even brought in by cattle. Therefore it is proposed that this vegetation type, which has received several different names by different authors, and has generally been included in the Chaco province (CABRERA, 1971 & 1976), is better called Austro-Brazilian Transitional Forest and should be excluded altogether from the Chaco s.s.

This forest is basically a very impoverished version of the Brazilian Subtropical Forests (sensu ANDRADE-LIMA, 1966, and KLEIN, 1972), which is here at the extreme western limit of its distribution. Furthermore, and despite the presence of some chaquenian elements in it, the Austro-Brazilian Transitional Forest should be regarded as a component of the Paranense province of the Amazonian Dominion (sensu CABRERA & WILLINK, 1980), since the dominant trees are species which are mainly distributed in other districts of this province where, however, they may be of lesser importance. Thus, of the Paranense province floristic stock, these species are probably those that can tolerate a few frosts each year, and can still compete successfully against the chaquenian species pushing east in areas with less alkaline and less waterlogged soils.

This interpretation is reinforced by the fact that most of the dominant species of the Austro-Brazilian Transitional Forest reappear further west, jumping over 700 km of dry Chaco plains, in the Yungas province (SMITH, 1962; RAGONESE & CASTIGLIONI, 1970; CABRERA, 1976) in the "Tipa-Pacará" (C) and "Palo blanco" (D) Forests. There is a very close relationship between those three vegetation units, as shown by several floristic groups of Table 1 and strongly supported by the numerical analysis. However, although units C & D also include some widespread chaquenian species, it is of interest that few authors tried to include these two forest types in the Chaco (e.g. HUECK, 1972), whilst virtually all phytogeographers have merged without hesitation the Austro-Brazilian Transitional Forest with the Chaco, with perhaps the sole exception of CASTELLANOS & PÉREZ-MOREAU (1944) in their map of vegetation of Argentina.

c) *The Eastern Chaco problem*

The fact that the so-called "Eastern Chaco" is in effect a meeting point of quite diverse floristic elements, i.e. true xerophytic chaquenian communities and humid subtropical and tropical species, has been perceived by some previous authors. Thus, CABRERA (1970) was aware that the boundary between the Chaquenian and Amazonian Dominia, which meet in the Eastern Chaco, is confused, and he remarked that the whole of the river Paraguay basin is an immense transition area with a very complicated intermingled pattern. A striking example of such intermingling is that the current concept of "Eastern Chaco" even includes "paratodales" of *Tabebuia caraiba* in floodable localities associated with the river Pilcomayo valley in SE Paraguayan Chaco region and NE Formosa in Argentina (CHODAT & VISCHER, 1977; FIEBRIG, 1933; RAGONESE & CASTIGLIONI, 1970; MORELLO & ADÁMOLI, 1974, p. 42). *T. caraiba* can hardly be accepted

¹These are: *Geoffroea decorticans*, *Aspidosperma quebracho-blanco*, *Ziziphus mistol*, *Prosopis nigra*, *Schinus fasciculatus*, *Acacia praecox* and *Caesalpinia paraguariensis* (see Table 1, FG XXII). Sometimes even *Schinopsis balansae* may appear in this forest, but its occurrence is very occasional and marginal to the unit (LEWIS, J. P., in litt.).

as Chaco species (see distribution map in PRADO & GIBBS, in press) since it also occurs in NE Brazil in the "sertão" and "agreste" (ANDRADE-LIMA, 1960 & 1989), is also an abundant species in the Cerrados (HERINGER & al., 1977; FURLEY & RATTER, 1988; RATTER & al., 1988), and the "paratodales" are one of the main features of the Pantanal landscape (RATTER, 1984; ALLEM & VALLS, 1987)!

Although edaphic factors can explain the eastward expansion of some Chaco taxa, the reason why southern or central Brazilian hygrophilous elements can expand westwards to interdigitate with chaquenian communities must be climatic. In Chaco vegetation maps the line drawn to separate the so-called "Eastern Chaco" from the rest of the province (see subregions map in MORELLO & ADÁMOLI, 1968; also HUECK & SEIBERT, 1981) seems to coincide roughly with a rainfall isoline somewhere between 950 to 1000 mm per year (PRADO, 1991), and also with the alleged limit between humid and dry climates, the MI (Moisture Index) = 1 line (BOX, 1986). In RAGONESE & CASTIGLIONI's map (1970), where their concept of "Eastern Chaco" covers the eastern half of the Gran Chaco region, this line coincides roughly with the western boundary of the "Pilaguense" and "Bosque Chaqueño" districts.

The climate classification systems of Koeppen and the two systems proposed by Thornthwaite have been compared for Argentina by BURGOS & VIDAL (1951), who concluded that Thornthwaite's Second System is the most fitted to the distribution of the natural vegetation. For the Argentine Chaco, this correlation is very good: the transitional belt proposed here (Fig. 4) matches very well with an equivalent narrow band along the river Paraná with C₂B₄'ra' climate (Fig. 3), and so does the zero isoline for the MI according to Thornthwaite's Second System (BURGOS & VIDAL, 1951, Fig. 7D). The similarities extend even to most of the chaquenian districts proposed by RAGONESE & CASTIGLIONI (1970), such as the "Pilaguense", "Matacense", and "Santiagoño" districts, to which the corresponding climate types are C₁A'da', DA'da' and DB₄'da' respectively, and the districts "Campestre", Mixed Forests & Savannas and Chaquenian Forest as a group with the corresponding climate type C₁B₃'da'. It is noteworthy that a very similar kind of climate to that of the eastern Transitional Belt appears to the west of the Chaco in areas occupied by both the "Tipa-Pacará" and "Palo blanco" Forests!

Given the weight of floristic and correlated climatic and edaphic evidence, the question could be posed now why the "Eastern Chaco" has traditionally been considered as truly chaquenian, as in MORELLO & ADÁMOLI (1968), RAGONESE & CASTIGLIONI (1970), CABRERA (1976), HUECK & SEIBERT (1981), RAMELLA & SPICHIGER (1989), SPICHIGER & RAMELLA (1989), and ZELLWEGER & al. (1990). The reasons are more historical than botanical. The rainforests or thorny dry woodlands formed a barrier to an eastward expansion for centuries with the result that most of the exsiccata collected on the eastern side of the Paraná and Paraguay rivers have simply the locality "Chaco", whether they are xerophytic or humid forest species, just because the collector ferried the river or sailed along the tributaries. The NE Argentine Chaco region and the Paraguayan Chaco region were cautiously explored by naturalists in a narrow fringe parallel to the big rivers or their tributaries, mainly the Pilcomayo (see the picturesque account by KERR, 1968), at the beginning of this century. HOCHREUTINER (1923) remarked that the Paraguayan Chaco was known only along the banks of the Paraguay and Pilcomayo rivers, and FIEBRIG (1933) was honest enough to admit that up to that time scientific exploration extended only for 100 to 150 km west of the river Paraguay (though the present author still believes that is an overestimate)¹. Geographically speaking the term Chaco will continue as the denomination of the whole of the region, but from a phytogeographical point of view the name must have a more restricted usage.

¹The first botanical collector to cross the Paraguayan Chaco by land was Teodoro Rojas, who followed the Paraguayan lines during the "Chaco War" and reached the Parapetí river in Bolivia in 1935. More evidence comes from the zoological field; the endemic peccary genus *Catagonus*, considered for long as extinct and known only from fossils, was rediscovered as late as in 1972 in the heart of the Paraguayan Chaco (WETZEL & al., 1975).




Fig. 3. — Climate types for the Argentine Chaco, according to the Second System of Thornthwaite. Modified from GALMARINI & R. DEL CAMPO, 1964.




Fig. 4. — A new proposal for the geographical limits of the Chaco sensu stricto. The transitional belt is indicated by oblique hatching.

d) *The Chaco sensu stricto*

By excluding these extraneous communities it is now possible to re-establish the geographical limits of the Chaco as a phytogeographical province. On the basis of a strict floristic list of woody and succulent species (PRADO, 1991), phytosociological study of the more relevant plant communities in the region both in classical and numerical analyses, and putting the vegetation of this area against the background of similar formations in South America (PRADO, 1991), a map of what is here regarded as Chaco *sensu stricto* is presented (Fig. 4). It must be noted that the Sierra Chaco, although taken as a separate entity throughout the analysis, is accepted as part of the Chaco s.s.; however, the *Polylepis australis* woodlands are not to be regarded as chaquenian, as in CABRERA (1976). In the construction of this new map some previous vegetation maps have been employed; i.e. CORO (1956) and ELLENBERG (1981) for the Bolivian Chaco, VERVOORST (in HAWKES & HJERTING, 1969) for NW Argentina up to La Rioja province, RAGONESE & CASTIGLIONI (1970) for SW Chaco in the provinces of La Rioja, San Juan, San Luis and Córdoba in part, SAYAGO (1969) and LUTI & al. (1979) for the Córdoba province, and D'ANGELO & al. (1987) together with LEWIS (1981) for the southern and eastern limit in the Santa Fe province. Also taken into consideration were the maps of the neighbouring phytogeographic provinces Monte (MORELLO, 1958) and Espinal (LEWIS & COLLANTES, 1973).

The extremely patterned vegetation in the east of the Chaco s.s. cannot be mapped with any precision at this stage with the knowledge available to date. A wide belt of transitional vegetation (Fig. 4) has been left open to further studies which could determine the exact localities where true Chaco vegetation can be found. However, to delimit this belt, information was taken from maps in CASTELLANOS & PÉREZ-MOREAU (1944), MORELLO & ADÁMOLI (1967), RAGONESE & CASTIGLIONI (1970), LEWIS (1981), ESSER (1982), and PRADO & al. (1989 & 1992), together with the present author's field experience and the plant distribution maps presented elsewhere (PRADO, 1991; PRADO & GIBBS, in press). Other differences with previous concepts of the Chaco shown in this new map consist in the exclusion of the vegetation of some of the "cerros" of Paraguay, such as the Cerros León, Cabrera and Chovoreca, clearly linked to the Subandean Piedmont Forests (RAMELLA & SPICHIGER, 1989) and also to the arboreal forms of Caatingas of NE Brazil (PRADO, 1991). Also excluded are the gallery forests on the rivers crossing the Chaco, particularly in their lower courses close to the Paraguay and Paraná rivers, and the vegetation of the Río Timane in the Paraguayan Chaco. Of the alleged Chaco vegetation in SW Brazil only a reduced sector remains as such (Fig. 4), together with some limited neighbouring transitional areas. It is noteworthy as well that the expansion of the Chaco s.s. beyond the line Santa Cruz de la Sierra-Chiquitos is here dramatically limited with respect to previous maps.

ACKNOWLEDGEMENTS

To Marta Bianchi and Peter Gibbs for their encouragement, and Paulo Oliveira for his vital help with computing. To J. P. Lewis and everybody at Plant Ecology, Cs. Agrarias, Rosario. This work was partially funded by the University of St. Andrews and the ORS Awards (U. K.) and CONICET and the National University of Rosario, Argentina.

REFERENCES

- ADÁMOLI, J. (1982). O pantanal e suas relações fitogeográficas com os Cerrados. Discussão sobre o conceito de "Complexo do Pantanal". In: Soc. Bot. Brasil (ed.), *An. XXXII Congr. Nac. Bot., Teresina*, 1981: 109-119. Universidade Federal do Piauí.
- ADÁMOLI, J., R. NEUMANN, A. D. R. DE COLINA & J. MORELLO (1972). El Chaco aluvional salteño (Convenio INTA-Prov. de Salta). *Rev. Inv. Agropec., INTA, Serie 3, Clima y Suelo* 9: 165-237.
- ALLEM, A. A. & J. F. M. VALLS (1987). *Recursos forrageiros nativos do Pantanal matogrossense*. Documentos, 8. EMBRAPA-CENARGEN, Brasília. x + 339 pp.
- ANDRADE-LIMA, D. DE (1960). Estudos fitogeográficos de Pernambuco. *Arq. Inst. de Pesq. Agron. Recife* 5: 305-341.
- ANDRADE-LIMA, D. DE (1966). Vegetação. In: IBGE (ed.), *Atlas Nacional do Brasil*. vol. II: 11. Conselho Nacional de Geografia, Rio de Janeiro.
- ANDRADE-LIMA, D. DE (1989). *Plantas das Caatingas*. Academia Brasileira de Ciências, Rio de Janeiro.

- BOX, E. O. (1986). Some climatic relationships of the vegetation of Argentina, in global perspective. *Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 91: 181-216.
- BROWN, A. D., S. C. CHALUKIAN & L. M. MALMIERCA (1985). Estudio florístico-estructural de un sector de selva semidecidual del noroeste argentino. I. Composición florística, densidad y diversidad. *Darwiniana* 26: 27-41.
- BURGOS, J. J. & A. L. VIDAL (1951). The climates of the Argentine republic according to the new Thornthwaite classification. *Ann. Ass. Amer. Geogr.* 41: 237-263.
- CABRERA, A. L. (1970). La vegetación del Paraguay en el cuadro fitogeográfico de América del Sur. *Bol. Soc. Arg. Bot.* 11 (Suppl.): 121-131.
- CABRERA, A. L. (1971). Fitogeografía de la República Argentina. *Bol. Soc. Arg. Bot.* 14: 1-42.
- CABRERA, A. L. (1976). *Regiones Fitogeográficas Argentinas*. 2nd ed. Enc. Arg. Agricultura y Jardinería. ACME S.A.C.I., Buenos Aires. 85 pp.
- CABRERA, A. L. & A. WILLINK (1980). *Biogeografía de América Latina*. 2nd ed. Serie de Biología. Secr. Gral OEA, Washington, DC. 122 pp.
- CASTELLANOS, A. & R. A. PÉREZ-MOREAU (1944). Los tipos de vegetación de la República Argentina. *Monogr. Inst. Est. Geogr., Tucumán* 4: 1-154, 30 pl., 1 map.
- CHODAT, R. & W. VISCHER (1977, orig. 1916). *La végétation du Paraguay*. J. Cramer, Vaduz. 558 pp.
- CORO, M. (1956). Distrito forestal de Yacuiña. Descripción de su área correspondiente a clima, suelo y vegetación. *Bol. Forestal, La Paz* 1: 53-72.
- D'ANGELO, C. H., D. E. PRADO, S. L. STOFELLA & J. P. LEWIS (1987). The Subchaquenian vegetation of the province of Santa Fe (Argentina). *Phytocoenologia* 15(3): 329-352.
- DIGILIO, A. P. L. & P. R. LEGNAME (1966). Los árboles indígenas de la provincia de Tucumán. *Op. Lilloana* 15: xxvii, 1-42, 107 descr.
- ELLENBERG, H. (1981). *Desarrollar sin destruir: respuestas de un ecólogo a 15 preguntas de agrónomos y planificadores bolivianos*. First ed. Instituto de Ecología, UMSA, La Paz. 55 pp.
- ESPINO, L. M., M. A. SEVESO & M. A. SABATIER (1983). *Mapa de Suelos de la Prov. de Santa Fe*. MAG Sta Fe-INTA EERA Rafaela. 220 pp.
- ESSER, G. (1982). *Vegetationsgliederung und Kakteenvegetation von Paraguay*. Akad. der Wissenschaften u. der Lit., Mainz. Mathematisch-Naturwissenschaftliche Klasse, Wiesbaden. 113 pp., 76 pl., 1 map.
- FIEBRIG, C. (1933). Ensayo fitogeográfico sobre el Chaco Boreal. *Rev. Jard. Bot. y Mus. Hist. Nat., Paraguay* 3: 1-87.
- FRANCESCHI, E. A. & J. P. LEWIS (1979). Notas sobre la vegetación del valle santafesino del Río Paraná (Rep. Argentina). *Ecosur* 6: 55-82.
- FURLEY, P. A. & J. A. RATTER (1988). Soil resources and plant communities of the central Brazilian cerrado and their development. *J. Biogeogr.* 15: 97-108.
- GALMARINI, A. & J. R. DEL CAMPO (1964). *Rasgos fundamentales que caracterizan el clima de la región chaqueña*. Cons. Nac. Desarrollo, Publ. N° 9, Buenos Aires. 178 pp.
- GENTRY, A. H. (1979). Distribution patterns of neotropical Bignoniaceae: some phytogeographic implications. In: LARSEN, K. & L. B. HOLM-NIELSEN (ed.), *Tropical Botany*: 339-354. Academic Press, London.
- HAWKES, J. G. & J. P. HJERTING (1969). *The potatoes of Argentina, Brazil, Uruguay and Paraguay. A biosystematic study*. Ann. Bot. Mem. N° 3. Clarendon Press & Oxford Univ. Press, Oxford. xxiii + 525 pp., 150 pl.
- HERINGER, E. P., G. M. BARROSO, J. A. RIZZO & C. T. RIZZINI (1977). A flora do Cerrado. In: FERRI, M. G., *IV Simposio sobre o cerrado*: 211-234. Editora USP, São Paulo.
- HOCHREUTNER, B. (1923). La végétation du Paraguay. *Bull. Inst. Nat. Genève* 45: 1-49.
- HUECK, K. (1955). Bosques chaquenhos e extração de tanino no Brasil. *Rev. Bras. Geog.* 17: 343-346.
- HUECK, K. (1959). Bosques secos de la zona tropical y subtropical de la América del Sur. *Bol. Inst. For. Lat. Amer. Invest. y Capac.* 4: 1-49.
- HUECK, K. (1972). *As florestas da América do Sul, Ecologia, composição e importância econômica*. Transl. H. Reichardt. Univ. de Brasília & Ed. Polígono, São Paulo.
- HUECK, K. & P. SEIBERT (1981). *Vegetationskarte von Südamerika*. Second ed., G. Fischer Verlag, Stuttgart. x + 90 pp.
- KERR, J. G. (1968). *A naturalist in the Gran Chaco*. Greenwood Publ., New York.
- KLEIN, R. M. (1967). Aspectos predominantes da vegetação sul-brasileira. In: Soc. Bot. Brasil (ed.), *An. XV Congr. Nac. Bot.*, Porto Alegre, 1964: 255-276.
- KLEIN, R. M. (1972). Árvores nativas da floresta subtropical do Uruguai. *Sellowia* 24.: 9-62.
- LEWIS, J. P. (1981). La vegetación de la provincia de Santa Fe. *GAEA* 9: 121-148.
- LEWIS, J. P. & M. COLLANTES (1973). El Espinal Periestépico. *Cien. Invest.* 29: 360-377.
- LEWIS, J. P. & E. F. PIRE (1981). *Reseña sobre la vegetación del Chaco santafesino*. Serie Fitogeográfica N° 18. INTA, Buenos Aires. 42 pp.
- LUTI, R., M. SOLÍS, F. GALERA, M. FERREIRA, M. HERRERA & J. C. BARRERA (1979). Vegetación. In: VÁZQUEZ, J., R. MIATELLO & M. ROQUE (eds.), *Geografía Física de la Provincia de Córdoba*: 297-368. Buenos Aires.
- MARLANGE, M. (1972). *Contribution à l'étude phyto-écologique du Chaco argentin*. Ph. D. thesis, Univ. des sciences et techniques du Languedoc, Acad. de Montpellier.

- MATEUCCI, S. D. & A. COLMA (1982). *Metodología para el estudio de la vegetación*. Serie de Biología. Secretaría General de la Organización de los Estados Americanos, Washington, DC.
- MEYER, T. (1944). Un viaje botánico al departamento de Orán (Provincia de Salta). *Rev. Geog. Americana* 21: 255-264.
- MEYER, T. (1963). Estudios sobre la selva de Tucumán. La selva de Mirtáceas de "Las Pavas". *Op. Lilloana* 10: 1-144, 55 pl.
- MORELLO, J. (1958). La provincia fitogeográfica del Monte. *Op. Lilloana* 2: 5-155, 58 pl.
- MORELLO, J. & J. ADÁMOLI (1967). *Vegetación y ambiente del nordeste del Chaco argentino. Guía de viaje, tramo Resistencia-Puerto Pilcomayo, IX Jorn. Arg. Bot.* INTA, Centro Reg. Chaqueño, E.E.A. Cnia. Benítez, mimeogr., 75 pp.
- MORELLO, J. & J. ADÁMOLI (1968). *Las grandes unidades de vegetación y ambiente del Chaco argentino. Primera parte: objetivos y metodología*. Serie Fitogeográfica N° 10. I.N.T.A., Buenos Aires. 125 pp.
- MORELLO, J. & J. ADÁMOLI (1974). *Las grandes unidades de vegetación y ambiente del Chaco argentino. Segunda parte: Vegetación y ambiente de la provincia del Chaco*. Serie Fitogeográfica N° 13. I.N.T.A., Buenos Aires. 130 pp.
- MORELLO, J., N. E. CRUDELLI & M. SARACENO (1971). *Los vinalares de Formosa (Rep. Argentina). La colonizadora leñosa Prosopis ruscifolia Gris*. Serie Fitogeográfica N° 11. I.N.T.A., Buenos Aires. 116 pp.
- MUELLER-DOMBOIS, D. & H. ELLENBERG (1974). *Aims and methods of vegetation ecology*. J. Wiley & Sons, New York.
- PRADO, D. E. (1991). *A critical evaluation of the floristic links between Chaco and Caatingas vegetation in South America*. Ph.D. thesis, University of St. Andrews.
- PRADO, D. E. (1993). What is the Gran Chaco vegetation in South America? I. A review. Contribution to the study of flora and vegetation of the Chaco. V. *Candollea* 48: 145-172.
- PRADO, D. E., E. A. FRANCESCHI & M. B. BIANCHI (1989). El bosque del escarpe occidental del Río Paraná (Argentina). Composición florística y estructura. *Acta Bot. Bras.* 3(2, suppl.): 99-108.
- PRADO, D. E. & P. E. GIBBS (in press). Patterns of species distributions in the dry seasonal forests of South America. *Ann. Missouri Bot. Gard.* 80(4).
- PRADO, D. E., P. E. GIBBS, A. POTT & V. J. POTT (1992). The Chaco-Pantanal transition in Southern Mato Grosso, Brazil. Pp 451-470. In: FURLEY, P. A., PROCTOR, J. & RATTER, J. A. (eds.), *Nature & Dynamics of Forest-Savanna Boundaries*. Chapman & Hall, London.
- RAGONESE, A. E. (1941). La vegetación de la provincia de Santa Fe (R.A.). *Darwiniana* 5: 369-416.
- RAGONESE, A. E. & J. A. CASTIGLIONI (1970). La vegetación del Parque Chaqueño. *Bol. Soc. Arg. Bot.* 11(Suppl.): 133-160.
- RAMELLA, L. & R. SPICHIGER (1989). Interpretación preliminar del medio físico y de la vegetación del Chaco Boreal. Contribución al estudio de la flora y de la vegetación del Chaco. I. *Candollea* 44: 639-680.
- RATTER, J. A. (1984). *Report of a visit to the Unidade de Execução de Pesquisa de Ambito Estadual (UEPAE) de EMBRAPA (Pantanal Station, Corumbá, MS)*. Royal Botanic Garden, Edinburgh, mimeogr.
- RATTER, J. A., A. POTT, V. J. POTT, C. N. DA CUNHA & M. HARIDASAN (1988). Observations on woody vegetation types in the Pantanal and at Corumbá, Brazil. *Notes RBG Edinb.* 45: 503-525.
- REBORATTI, H. J. & J. J. NEIFF (1986). Los bosques fluviales del Chaco oriental: primera aproximación. *Amb. Subtropical* 1: 36-46.
- SARMIENTO, G. (1972). Ecological and floristic convergences between seasonal plant formations of tropical and subtropical South America. *J. Ecol.* 60: 367-410.
- SAYAGO, M. (1969). Estudio fitogeográfico del norte de Córdoba. *Bol. Acad. Nac. Cienc., Córdoba* 46: 123-427.
- SCHULZ, A. G. (1967). *Reserva Biológica de Colonia Benítez. Catálogo de la flora de la selva*. INTA, Centro Reg. Chaqueño, E.E.A. Cnia. Benítez, mimeogr., 7 pp.
- SMITH, L. B. (1962). Origins of the flora of southern Brazil. *Contr. US Nat. Herb.* 35: 215-250.
- SØRENSEN, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species content. *K. Dansk. Vidensk. Selsk. Biol. Skrift* 5: 1-34.
- SPICHIGER, R. & L. RAMELLA (1989). The forests of the Paraguayan Chaco. In: HOLM-NIELSEN, L. B. & al. (eds.), *Tropical forests, botanical dynamics, speciation and diversity*: 259-270. Academic Press, London.
- UNZUETA, O. (1975). *Mapa ecológico de Bolivia. Memoria explicativa*. Min. Asuntos Camp. y Agrop., Rep. de Bolivia, La Paz. vi + 312 pp.
- WETZEL, R. M., R. E. DUBOS, R. L. MARTIN & P. MYERS (1975). Catagonus, an "extinct" peccary, alive in Paraguay. *Science* 189: 379-381.
- WISHART, D. (1987). *Clustan User Manual. Cluster analysis software*. St. Andrews: Computing Laboratory, University of St. Andrews. Fourth version.
- ZELLWEGER, C., R. PALESE, P. PERRET, L. RAMELLA & R. SPICHIGER (1990). Concept and use of an integrated database system for the Chaco. Application to a preliminary checklist. Contribution to the study of the flora and vegetation of the Chaco. II. *Candollea* 45: 681-690.

