

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	49 (1994)
<b>Heft:</b>	1
<b>Artikel:</b>	Calli, cell and plantlet suspension cultures of bryophytes
<b>Autor:</b>	Felix, Hansruedi
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-879533">https://doi.org/10.5169/seals-879533</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Calli, cell and plantlet suspension cultures of bryophytes

HANSRUEDI FELIX

## ABSTRACT

FELIX, H. (1994). Calli, cell and plantlet suspension cultures of bryophytes. *Candollea* 49: 141-158. In English, English and French abstract.

Calli, cell cultures or cultures of whole plantlets have been established of a range of bryophytes. The cultures have favourable biochemical and physiological characteristics, making them a system of high potential for morphological and physiological studies as well as for the in vitro production of valuable bryophyte-derived metabolites. To date relatively few papers have been published on cultures of bryophytes, the use of these cultures for production of metabolites is even more rare. This review summarizes the publications on calli, cell cultures or cultures of whole plantlets of bryophytes.

## RÉSUMÉ

FELIX, H. (1994). Calli, cultures de cellules et de plantes entières des bryophytes. *Candollea* 49: 141-158. En anglais, résumés anglais et français.

Des calli, des cultures de cellules ou des plantes entières ont été établis pour de nombreuses bryophytes. Les cultures ont des caractères biologiques et physiologiques favorables pour des études morphologiques et physiologiques et pour la production de substances de haute valeur. Jusqu'à maintenant il n'y a pas beaucoup de publications sur la culture des bryophytes. L'utilisation des cultures pour la production des métabolites est encore plus rare. Cet exposé rassemble les publications sur les calli et les cultures de cellules et de plantes entières.

**KEY WORDS:** Bryophytes — Mosses — Cell culture — Secondary metabolites — Protonemata culture — Plantlet culture — Photoautotropic growth.

## 1. Introduction

Plant cell culture has been proposed as an alternative supply of plant mass and phytochemicals to the normal plantation system. For bryophytes there are no extended plantation systems, thus the need for bryophyte cell cultures is even more obvious. Such systems can provide a product of constant quality and quantity, free from constraints of season, climate or disease considerations. No process, however, will be adapted unless it can compete on cost, or confer some real advantages over existing sources. Bryophyte cell cultures can act as a supply of completely novel substances opening new market areas.

In this review it will be shown, that progress in bryophyte cell culture did not go as fast as with cell culture techniques of higher plants, although even there only a few commercial applications are known. There is, however, an interesting fact with bryophyte cell cultures: Many research groups cultivate whole plantlets or protonemata in culture instead of single cells or cell aggregates. This is not possible with higher plants.

Bryophyte calli have been discovered a long time ago (VON WETTSTEIN, 1953; MEYER, 1953). Bryophyte cell cultures were mainly used to answer morphological and physiological questions. A research group in Saarbrücken (Germany; ZINSMEISTER & al., 1991) uses bryophyte culture techniques to get enough plant material to isolate interesting secondary metabolites, which can be screened in medical and agrochemical tests.

Several authors summarized the use of calli and suspension cultures. HUNECK (1986) stated that relatively few papers have been published on the biochemistry and chemistry of cell cultures of bryophytes. CHOPRA & KUMRA (1988) discussed the study of callus formation and its differentiation as an additional tool to investigate life cycles.

The present review lists the bryophytes cultivated as calli, cell suspension or plantlet suspension; it does not contain extended information on axenic cultures of whole bryophytes on solid media, which can be found elsewhere (LAL, 1984; SARGENT, 1988; McQUEEN, 1991). Additionally information is summarized about the investigations carried out with these cultures. The aim of this review is to encourage further research in this branch of biology and bio-organic chemistry.

## 2. Establishment of cell cultures

As shown in Table 1 (row A) bryophyte cell cultures originate from different sources: a) from spores, b) from gametophytes, c) from gametophytic buds and gemmae, d) from sporogons, e) from protoplasts. There is indication that the origin of the callus plays a role: Different types of callus tissues appear to originate from different tissues of thallus (TRIPATHY, 1978a, 1978b). They differ in their external and internal morphology, pattern of growth, potentiality for differentiation and physiological requirements.

In a first step bryophyte parts need to be sterilized. The majority of species are surface sterilized using the washing machine technique (BASILE, 1972; BASILE & BASILE, 1988) on either shoot tips or unopened capsules. Clorox (9%) for 1 minute works best for most species, but the time and concentration have to be modified for delicate and tough specimens. Spore suspensions from surface sterilized capsules are spread on agar-plates, and spores from untreated capsules are spread by the top technique described by DOYLE (1967). A few species, e.g. *Treubia* and *Cryptothallus*, have proven to be totally recalcitrant to surface sterilization, probably due to obligate, symbiotic fungi and fungal and bacterial (especially blue-green algae) contaminants, which are able to destroy bryophyte cell cultures.

LAL (1984) and SARGENT (1988) list a whole number of species cultured under sterile conditions together with recommended growth conditions. SIMON (1988) emphasizes particularly the sterile cultivation of *Sphagnum* species, which can be a source for calli or suspension cultures. KATOH (1988) describes in a general way possibilities to get calli and suspension cultures. Table 1 summarizes literature on culture of bryophytes in general (row a) and literature on effects of different factors on growth of calli or suspension cultures (rows c, d, e, f, g, h). Plant hormones and other compounds like chelated iron, vitamins, light, temperature, humidity, sugars, pH, bacteria all influence growth. Plant hormones (BOPP, 1983; COVE & ASHTON, 1984) and sugars (KATOH, 1988; JONES, 1978) seem to play a major role. According to the literature (Table 1, row c) hormones strongly influence bryophytes as well as higher plants. UV light can substitute for growth regulators (KUMRA & CHOPRA, 1982). The effect of sugars (Table 1, row f) was tentatively explained by JONES (1978) who found soluble carbohydrates in inorganic culture solutions containing aseptically cultivated *Sphagnum* shoots. He concluded that to grow normally in liquid culture, *Sphagnum* requires sugars to compensate for those carbohydrates lost by leakage. Activated charcoal can inhibit calussing of protonema (RAHBAR, 1982), indicating adsorption of an essential compound.

Not all bryophytes are cultured in the form of undifferentiated tissue (calli, suspension), many authors use axenic suspension cultures of whole plantlets, protonemata or chloronemata (Table 1, row q). These cultures have special requirements. Repeatedly researchers mentioned that inorganic media are insufficient to allow normal growth and development of *Sphagnum* in axenic cultures (SIMOLA, 1969; HINTIKKA, 1972; JONES, 1978; BAKER & BOATMAN, 1985; KAJITA

& al., 1987). In non-agitated batch cultures the protonema development, bud formation, and growth of shoots are poor without any addition of an organic carbon source. SIMOLA (1969, 1975, 1979) studied the effect of several sugars, amino acids and dipeptides on the growth of *Sphagnum* in axenic cultures. Ca. 200-230 mg dry weight were produced in 100 days when one 10 mm shoot tip was used as inoculum per flask containing nutrient medium with 1% sucrose as a carbon source (SIMOLA, 1977; RUDOLPH & al., 1988).

Several authors developed fermenters for the cultivation of bryophytes (Table 1, row B; BOYD & al., 1988; RUDOLPH & al., 1988; JACUBOWSKI & RUDOLPH, 1989; OHTA & TAKATA, 1990; ZINSMEISTER & al., 1991). RUDOLPH & al. (1988) describe a continuous feed fermenter for whole plantlets. Multicellular organisms form large aggregates, which are difficult to deal with in fermenters, for clumping soon takes place, together with layering. It is difficult to achieve adequate agitation, especially turbulence, without damaging the branched leafy moss shoots. Therefore common types of stirrers are not appropriate. An airlift fermenter (rising current of air) is specially suited to agitate delicate moss shoots. The airlift system described by BOYD & al. (1988) provides a supply of protonemal tissue for analysis, and allows time courses to be followed, for example after the addition of a plant hormone. *Physcomitrella patens* will develop from the protonemal stage in this system to give small plantlets, consisting of a protonemal ball with gametophores, but such development is delayed compared with growth on solid media (20-24 days instead of 10-14 days). If gametophytic tissue is required, the fermenter will need some alterations, as clogging of the outflow will be a problem. Alternatively, if longer production runs of protonemal tissue are required, it may be possible to manipulate the culture medium in order to extend the protonemal phase. To keep the culture axenic proved to be most difficult, even though there was no organic carbon source in the medium. Only by a rigorous test of all possible contamination sources was this problem eliminated. For continuous culture, a representative outflow is essential, in order that the specific growth rate, calculated from the dilution rate and the steady state dry weight, is accurate.

### 3. Effect of physical parameters on growth

As can be expected the environmental conditions play a major role in the growth of bryophytes in cultures. The influence of light including UV light (Table 1, row d) seems to be especially important. Bryophytes grow better in culture, if light is not completely absent. Photoautotrophic growth of bryophytes in culture is easier achieved than with higher plants (see chapter on photoautotrophic growth), most probably this is due to a higher chlorophyll content. Other physical parameters like temperature, humidity and pH (Table 1, rows e and g) were also investigated.

### 4. Photoautotrophic growth

The majority of cell cultures of higher plants exhibit a low rate of photosynthetic growth, probably because the chlorophyll content of the cells is low. Callus cells from bryophytes contain a high concentration of chlorophyll (ALLSOPP, 1957; WARD, 1960a & 1960b), but few attempts have been made to establish photoautotrophic suspension cultures of bryophyte cells (Table 1, row k). The cells of liverworts grow rapidly in the light in liquid culture media containing glucose, they contain a high amount of chlorophyll. In contrast to most cultured cells from seed plants, liverwort cells apparently do not grow in the dark even when glucose is present. A cell line isolated from *Marchantia polymorpha* was capable of photoautotrophic growth under air enriched with 1% CO<sub>2</sub> (KATOH, 1983b), *Sphagnum imbricatum* could also be grown under elevated CO<sub>2</sub> concentration (KAJITA & al., 1987). When chlorophyllous cells of *Marchantia paleacea* that had been grown heterotrophically in the dark were transferred to phototrophic conditions, these cells started to grow photoautotrophically without a lag phase (NGUMI & al., 1990), maybe the chlorophyll content is less controlled by light.

Table 1. — Bryophytes listed alphabetically together with a brief description of investigations carried out.

Table 1 (cont.)

Table 1 (cont.)

Bryophytes	A	B	Investigations carried out												References							
			a	b	c	d	e	f	g	h	i	k	l	m	n	o	p	q	r	s	t	u
<i>Marchantia polymorpha</i> .....	4	C	X	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	ONO, 1973
<i>M. polymorpha</i> .....	4	C	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ONO, 1974, 1976a & 1976b
<i>M. polymorpha</i> .....	4	S	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	OHTA & al., 1977
<i>M. polymorpha</i> .....	4	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ONO & al., 1979
<i>M. polymorpha</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	KATOH, 1983a & 1983b; KATOH & al., 1979 & 1980
<i>M. polymorpha</i> .....	4	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	OHYAMA & al., 1982 & 1983; TANAKA & al., 1984
<i>M. polymorpha</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	KONNO & al., 1983; TAKIO, 1987; TAKIO & HINO, 1988
<i>M. polymorpha</i> .....	4	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	SUGAWARA, 1983
<i>M. polymorpha</i> .....	2	S	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ABE & OHTA, 1983 & 1984; OHTA & al., 1983 & 1984
<i>M. polymorpha</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ONO & OKAMOTO, 1984
<i>M. polymorpha</i> .....	4	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	IMOTO & OHTA, 1985
<i>M. polymorpha</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	KONNO & al., 1987; SATO & KATO, 1988
<i>M. polymorpha</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	OHTA & TAKATA, 1990
<i>M. polymorpha</i> .....	4	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	HAMADA & al., 1991
<i>M. polymorpha</i> .....	1	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ALLSOPP & ILAHI, 1969; ILAHI & ALLSOPP, 1969 & 1970
<i>Moerckia floriotiana</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ALLSOPP & ILAHI, 1970a & 1970b
<i>Notocladia confusa</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ILAHI, 1972
<i>N. confusa</i> .....	1	C	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ONO & al., 1988
<i>Peltia endiviifolia</i> .....	2	C,S	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ILAHI & ALLSOPP, 1969 & 1970
<i>P. epiphylla</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	BECKER, 1990
<i>P. epiphylla</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ONO & al., 1988
<i>P. neesiana</i> .....	2	C,S	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ILAHI, 1972
<i>Petalophyllum rafisii</i> .....	2	C	X	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	BEUTELMANN & al., 1978
<i>Physcomitrella patens</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	GRIMSLEY & WITHERS, 1983
<i>P. patens</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	RESKI & ABEL, 1985
<i>P. patens</i> .....	2	S	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ABEL & al., 1989
<i>P. patens</i> .....	2	S,f	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	BOYD & al., 1988
<i>Physcomitrium coorgense</i> .....	3	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	ILA, 1961
<i>P. pyriforme</i> .....	5	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	BAUER, 1957 & 1961
<i>P. pyriforme</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	MENON & LAL, 1972
<i>P. pyriforme</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	MENON & LAL, 1974
<i>P. pyriforme</i> × <i>F. hygrometrica</i> .....	2	C	.	X	.	.	.	.	.	.	.	X	.	.	.	.	.	.	.	.	.	MENON & HARTMANN, 1988a & 1988b
																					BAUER, 1961 & 1963	

Table 1 (cont.)

Table 1 (cont.)

Bryophytes	A	B	Investigations carried out												References								
			a	b	c	d	e	f	g	h	i	k	l	m	n	o	p	q	r	s	t	u	
<i>Scapania nemorea</i> ....	2	S	.	.	x	.	.	.	.	.	.	.	.	x	x	.	.	.	.	.	.	.	BECKER, 1990
<i>S. nemorea</i> ....	4	C	.	x	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	BASILE, 1965
<i>Sphaerocarpos donnellii</i> ....	3	S	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	MACHLIS & DOYLE, 1962; MACHLIS, 1962
<i>S. donnellii</i> ....	3	S	.	.	.	x	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	MILLER & MACHLIS, 1968
<i>S. donnellii</i> ....	3	S	.	.	.	x	.	.	x	.	.	.	.	x	.	.	.	.	.	.	.	GRUSAK & al., 1980	
<i>S. texanus</i> ....	3	S	x	.	x	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	MONTAGUE & TAYLOR, 1971	
<i>S. texanus</i> ....	3	S	x	.	x	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	STAHLY & al., 1977	
<i>Sphagnum cuspidatum</i> ....	2	S	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	BAKER & BOATMAN, 1985
<i>S. cuspidatum</i> ....	2	S	.	x	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	RUDOLPH & al., 1988
<i>S. fallax</i> ....	2	S	.	x	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	HINTIKKA, 1972
<i>S. fallax</i> ....	2	f	.	.	.	.	.	.	.	.	x	.	.	.	x	.	.	.	.	.	.	.	JACUBOWSKI & RUDOLPH, 1989
<i>S. fallax</i> ....	2	S	.	.	x	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	RUDOLPH & al., 1988
<i>S. fimbriatum</i> ....	2	S	.	x	.	.	.	.	.	.	.	x	.	.	.	x	.	.	.	.	.	.	SIMOLA, 1977
<i>S. imbricatum</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	KAJITA & al., 1987
<i>S. imbricatum</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	TAKAMI & al., 1990
<i>S. imbricatum</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	RUDOLPH & al., 1988
<i>S. magellanicum</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	.	x	.	.	.	.	.	.	JACUBOWSKI & RUDOLPH, 1989
<i>S. magellanicum</i> ....	2	f	.	.	.	.	.	.	.	.	x	.	.	.	.	x	.	.	.	.	.	.	RUDOLPH & al., 1988
<i>S. majus</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	.	x	.	.	.	.	.	.	SPOERLE & al., 1990
<i>Symphyogyna brongniartii</i> ....	2	S	.	x	.	.	.	.	.	.	x	.	.	.	x	.	.	.	.	.	.	.	SPIESS, 1976
<i>Thuidium delicatulum</i> ....	2	c	.	x	.	.	.	.	.	.	x	.	.	.	x	.	.	.	.	.	.	.	BECKER, 1990
<i>Trichocolea tomentella</i> ....	1	S	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	

Areas of research:

a establishment of cultures

b differentiation, morphogenetic studies

c effects of plant hormones and other compounds

d effects of light (UV light included) on growth

e effects of temperature or humidity on growth

f effects of sugars on growth

g effect of pH on growth

h effects of *Agrobacterium tumefaciens* and other bacteria

i enzymology

j photoautotrophic growth

k chloroplast DNA

l excretion of compounds

m production of primary and secondary metabolites

n culture of protoplasts

o photosynthetic activity

p whole plants in culture (incl. protonemata, chloronemata)

q mutants

r preservation of cultures

s oxygen metabolism

t cell wall biosynthesis

u

A = Origin of cultures:

1 callus of undefined origin

2 from spores

3 from whole gametophytes, surface sterilized

4 gametophytic buds, gemmae

5 sporogons

6 from protoplasts

B = Type of culture:

c callus

s suspension culture

f fermenter

Bryophyte	Type of culture	Doubling time (days)	References
<i>Barbula unguiculata</i> .....	photoautotrophic growth, suspension culture	4,5	TAKAMI & al., 1988b
<i>Physcomitrella patens</i> .....	protonemata in airlift fermenter	1,1	BOYD & al., 1988
<i>P. patens</i> .....	protonemata in spherical aeration vessel	4,4	WANG & al., 1981a
<i>Marchantia polymorpha</i> ...	suspension culture, photoautotrophic growth	1,1	KATOH, 1983b
<i>M. polymorpha</i> .....	jar-fermenter, photoautotrophic growth	2	OHTA & TAKATA, 1990
<i>Pellia endiviifolia</i> .....	suspension culture	7,4	ONO & al., 1988
<i>Ricciocarpos natans</i> .....	axenic culture of whole bryophytes	4,1	WURZEL & BECKER, 1990
<i>Sphagnum</i> sp. .....	cell suspension culture in the light	3,3	KAJITA & al., 1987
<i>Sphagnum fallax</i> .....	fermenter, whole plants	5	RUDOLPH & al., 1988
<i>Marchantia paleacea</i> .....	cell suspension culture, photoautotrophic growth	4,7	NGUMI & al., 1990

Table 2. — Growth rates of bryophyte cultures.

A high cell density makes it quite difficult to irradiate cells in large-scale culture such as a jar-fermenter. If illumination is provided from outside of the vessel, only the cells near the surface of side walls will receive sufficient intensity of light. In order to cultivate liverwort cells in a jar-fermenter under illumination, a new culture method was developed in which light is introduced directly into the culture vessel by using optical fibres (OHTA & TAKATA, 1990).

### 5. Growth rates in suspension cultures and fermenters

In general microbial cells have rapid growth rates with doubling times measured in hours, whereas in plant cell suspension cultures this is measured in days. One of the fastest growing cultures has been tobacco with doubling times of 18 hours, but a more typical time would be 2-3 days (NOGUCHI & al., 1977). Comparing these values with doubling times of bryophyte cultures (Table 2), the growth rates of bryophytes are very similar. It is, however, remarkable, that without special optimization some bryophytes grow as fast as the particular tobacco strain. Over all, similar problems will be observed in a bioreactor with higher plants and bryophytes, a run of a plant suspension can be from one to four weeks which means a much greater difficulty in maintaining sterility than with bacteria, because slow growth of plant cells means that almost any contamination will outgrow the culture.

### 6. Storage of cell cultures

Continuous vegetative propagation in sterile culture is used as a storage method for bryophytes, since many do not produce spores due to self-sterility or the presence of a developmental block in the gametophyte. Usually it is possible to cope with subculturing demands. However, the number of stock strains can increase beyond this point and, more importantly, there is evidence of strains undergoing changes with time in culture. The risk of mutations accumulating in culture and risks of loss of irreplaceable strains by microbial contaminations, which can be very difficult to eradicate, have led GRIMSLEY & WITHERS (1983) to develop a stable storage by cryopreservation in liquid nitrogen.

## 7. Bryophytes used

In Table 1 bryophytes are listed alphabetically together with a brief description of the investigations carried out and the kind of culture used (callus, suspension culture, culture of whole plants, fermenters). It is evident, as in many areas of biology, that there are favorite organisms, e.g. *Marchantia polymorpha*, *Physcomitrium pyriforme*, *Sphagnum* spp. Compared to the countless number of investigations carried out using cell cultures of higher plants, the use of bryophyte cultures is more limited. Not even all bryophyte families were considered. The reason for this neglect may be, that there are no crops among bryophytes, bryophytes are also not used as a model for higher plants. The interest may increase a little bit, if there is a need for sufficient plant material to isolate secondary metabolites (ZINSMEISTER & al., 1991).

## 8. Investigations carried out

Cell cultures of hepaticas and mosses are used for the same purposes as cultures of higher plants, namely to get reproducible and comparable results. Essentially the same investigations are carried out: studies of morphogenesis, differentiation, effects of different factors on growth, physiology, biochemical studies, enzymes, photosynthesis, oxygen metabolism, cell wall biosynthesis, genetics, production of primary and secondary metabolites.

### 8.1. Differentiation, morphogenetic studies

Bryophytes have proven to be a good material to study morphogenesis (Table 1, row b). Calli can readily be induced from different parts of the gametophyte and from the sporophyte, and it can be made to differentiate into gametophytes or sporophytes, respectively, by modifying the cultural conditions. This is possible by variation of single elements or by changing the relative concentration of elements in the different solutions (Table 1, rows c and f). Therefore it must be shown for each bryophyte material which of the nutritional solutions may provide the optimal results in vitro. The use of protoplasts as starting material for morphological studies has also been described (Table 1, row o). Agrobacteria, the causal agents of the crown gall, cane gall, and hairy root diseases in higher plants, as well as other bacteria, accelerate normal development and induce abnormal changes in bryophytes also. Pathogenic strains are typically most effective in bringing about these developmental changes, which require physical contact between the bacteria and the moss. SPIESS & al. (1981) conclude that many bacteria closely associated with mosses in nature are capable of influencing moss development in the protonemal stage and thus could play a role in the normal initiation of gametophore development. While some of these strains may prove to be agrobacteria, the tests employed were insufficient to establish such an identification. Besides studying the effect of bacteria the influence of fungi on bryophyte development also deserves further attention (FELIX, 1988).

### 8.2. Biochemical studies

Compared to the biochemical knowledge about higher plants, much less is known about bryophytes. Bryophyte cell cultures are an ideal source for tissue, because reproducible and comparable results can be obtained. A few enzymological investigations were carried out (Table 1, row i), whereby enzymes were extracted from cell cultures. Chloroplast DNA was examined in bryophytes (Table 1, row l), photosynthetic activity (Table 1, row p), oxygen metabolism (Table 1, row t) and cell wall biosynthesis (Table 1, row u) were followed. A detailed description of the reports is not given here, because the experiments must be seen in other contexts than just the use of cell cultures.

### 8.3. Genetic studies

COVE & ASHTON (1984) and KNIGHT & al. (1988) describe in detail the use of mutants to study the role of hormones in moss development and other processes, in these reviews it was not indicated, whether also cell cultures were mutagenized. BEUTELMANN & al. (1978) treated spores with N-methyl-N'-nitro-N-nitrosoguanidine and established cell cultures from them. TAKIO (1990) selected mutants directly from cell cultures after incubation of cells in chlorate.

### 8.4. Production of primary and secondary metabolites

Between 25% and 30% of prescribed pharmaceuticals are extracted from plant sources. Plants are also used as sources of food colourings and flavourings, and also in the production of agrochemicals. These products supply a market worth billions of dollars per annum. For compounds of plant origin the source is usually assumed to be a higher plant. The chemical analysis of bryophytes was neglected for a long time, the main reason for this fact may be the difficulty to collect sufficient amounts of bryophytes to isolate the pure compound. The easier it was to characterize minute amounts of isolated compounds the more secondary metabolites could be investigated. Nevertheless only about 5% of all known bryophyte species were investigated for their chemical constituents.

An interesting feature of bryophytes is that they are relatively free from attack by parasitic microorganisms (FELIX, 1988), e.g. herbarium specimen need no special treatment like those of higher plants. The cause of this may be anatomical, immunological or may be due to the presence of antimicrobial activity or other factors. In recent publications on disease resistance of plants (KUC, 1990) and on the role of secondary metabolites in chemical defence mechanisms (HARBORNE, 1990) bryophytes were not mentioned at all, obviously this phenomenon was not studied in bryophytes. In this context it would also be interesting to study phytoalexins as part of induced defence reactions in bryophytes as it is done in higher plants (BARZ & al., 1990). It is not known whether phytoalexins exist in bryophytes at all, some hints can be found in several publications (ZINSMEISTER & MUES, 1990). Besides antibiotic activity of bryophytes (BANERJEE & SEN, 1979; ASAOKAWA, 1990) other biological effects of compounds from bryophytes were observed: allergenic effects, cytotoxic and cytostatic effects, cardiotonic activity, carcinogenic activity, antiinflammatory activity, vasopressin-antagonistic activity, release of hyperoxide, piscicidal, molluscicidal and feed repellent effects, growth regulation in higher plants, enzyme inhibition (HUNECK, 1983; ZINSMEISTER & MUES, 1990; ZINSMEISTER & al., 1991; ASAOKAWA, 1990). A relatively large screening program for antitumor agents was carried out (SPJUT & al., 1986). In this report the problem of collecting enough material was mentioned: samples weighing 0.5 — 2 kg (dried) were collected for each of 208 species and varieties. In vitro cultures offer the possibility to obtain sufficient material (BECKER, 1990; KATO & TAKEDA, 1990; OHTA & TAKATA, 1990; SAUERWEIN & BECKER, 1990). This technique is used for intensive screening programs at the University of Saarbrücken, Germany (BECKER, 1990; ZINSMEISTER & al., 1991).

Mainly liverwort cultures were screened for secondary metabolites. SAUERWEIN & BECKER (1990) and WURZEL & BECKER (1990) found that *Fossombronia pusilla* and *Ricciocarpus natans*, respectively, produced the same pattern of secondary metabolites as field collected material (perrottianal, santonin, 7 terpenes with antimicrobial activity). The content of the terpenoid decreases on nitrogen and phosphorous-deficient media. MORAIS & BECKER (1990) found that differentiated cultures on agar produced sesquiterpenes in tenfold increased amount compared to undifferentiated suspension cultures. TAKEDA & KATO (1981, 1983a, 1983b) found that *Calypogeia granulata* had a higher yield in essential oils in suspension cultured cells than for freely growing, intact plants. The hypothesis of bryophytes producing secondary metabolites independent of their developmental stage is controversial (H. BECKER, personal communication). ABE & OHTA (1983, 1984) and OHTA & al. (1983, 1984) investigated prelunularic acid and lunularic acid in various liverworts. Lunularic acid may be a secondary metabolite or an endogenous bryophyte growth regulator, it may control drought resistance. The accumulation of lunularic acid is inversely correlated with the amount of phosphate in the medium. The age-dependent accumulation of

lunularic acid can be explained as the result of depletion of phosphate in the medium. Some compounds are excreted into the medium (Table 1, row m). Excretion does not seem to be dependent on the size of the molecules.

HAMADA & al. (1991) describe a green cell suspension culture of the liverwort *Marchantia polymorpha* converting testosterone to 6 $\beta$ -hydroxytestosterone and epitestosterone to androst-4-ene-3,17-dione.

Commercial interest in secondary metabolites of bryophytes from cell or whole plant culture origin is still rare. Lack of material for extraction and following difficulties with structure analysis may have been the reason. The chemical company BASF together with ZINSMEISTER & al. (1991) developed an extended screening program. MITSUBISHI-GAS-CHEM. (1988) may also have a commercial interest in secondary metabolites.

#### REFERENCES

- ABE, S. & Y. OHTA (1983). Lunularic acid in cell suspension cultures of *Marchantia polymorpha*. *Phytochem.* 22: 1917-1920.
- ABE, S. & Y. OHTA (1984). The concentrations of lunularic acid and prelunularic acid in liverworts. *Phytochem.* 23: 1379-1381.
- ABEL, W.O., W. KNEBEL, H.-U. KOOP, J. R. MARIENFELD, H. QUADER, R. RESKI, E. SCHNEPF & B. SPOERLEIN (1989). A cytokinin-sensitive mutant of the moss *Physcomitrella patens*, defective in chloroplast division. *Protoplasma* 152: 1-13.
- ALLSOPP, A. (1957). Controlled differentiation in cultures of two liverworts. *Nature* 179: 681-682.
- ALLSOPP, A. & I. ILAHI (1969). Studies in the Metzgeriales (Hepaticae). I. The effects of certain sugars on the growth and morphology of some thalloid species. *Phytomorph.* 19: 242-253.
- Allsopp, A. & I. ILAHI (1970a). Studies in the Metzgeriales (Hepaticae). III. The effects of sugars and auxins on *Noteroclada confluens*. *Phytomorph.* 20: 9-16.
- ALLSOPP, A. & I. ILAHI (1970b). Studies in the Metzgeriales (Hepaticae). VII. Regeneration in *Noteroclada confluens* and *Blasia pusilla*. *Phytomorph.* 20: 173-182.
- ASAKAWA, Y. (1990). Terpenoids and aromatic compounds with pharmacological activity from bryophytes. In: ZINSMEISTER, H. D. & R. MUES (eds.), *Proc. Phytochem. Soc. Europe 29. Bryophytes: their chemistry and chemical taxonomy*, pp. 369-410. Clarendon Press, Oxford.
- BAKER, R. G. & D. J. BOATMAN (1985). The effect of carbon dioxide on the growth and vegetative reproduction of *Sphagnum cuspidatum* in aqueous solutions. *J. Bryol.* 13: 399-406.
- BANERJEE, R. D. & S. P. SEN (1979). Antibiotic activity of bryophytes. *Bryologist* 82: 141-153.
- BARZ, W., W. BLESS, G. BÖRGER-PAPENDORF, W. GUNIA, U. MACKENBROCK, CH. OTTO & E. SÜPER (1990). Phytoalexins as part of induced defence reactions in plants: their elicitation, function and metabolism. In: *Bioactive compounds from plants. Ciba Foundation Symposium 154*: 140-153. John Wiley & Sons Inc., Chichester.
- BASILE, D. V. (1965). Growth, development and gemma formation in the liverwort *Scapania nemorosa* as influenced by L-arginine, L-histidine and L-glutamic acid. *Amer. J. Bot.* 52: 443-454.
- BASILE, D. V. (1972). A method for surface-sterilizing small plant parts. *Bull. Torrey Bot. Club* 99: 313-316.
- BASILE, D. V. (1979). Hydroxyproline-induced changes in form, apical development, and cell wall protein in the liverwort *Plagiochila arctica*. *Amer. J. Bot.* 66: 776-783.
- BASILE, D. V. & M. R. BASILE (1988). Procedures used for the axenic culture and experimental treatment of bryophytes. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 1-16. Hattori Bot. Lab., Nichinan.
- BAUER, L. (1957). Regenerationsversuche am Sporogon von *Physcomitrium pyriforme* (Brid.). *Ber. Deutsch. Bot. Ges.* 70: 424-432.
- BAUER, L. (1961). Über die Kalluswuchsform des Laubmoosporophyten. *Naturwissenschaften* 48: 507-508.
- BAUER, L. (1963). On the physiology of sporogonium differentiation in mosses. *J. Linn. Soc. (Bot.)* 58: 343-351.
- BAUER, L. (1966). Isolierung und Testung einer kinetinartigen Substanz aus Kalluszellen von Laubmoosporophyten. *Z. Pflanzenphysiol.* 54: 241-253.
- BECKER, H. (1990). Secondary metabolites from in vitro cultures of liverworts. In: ZINSMEISTER, H. D. & R. MUES (eds.), *Proc. Phytochem. Soc. Europe 29. Bryophytes: their chemistry and chemical taxonomy*, pp. 339-348. Clarendon Press, Oxford.
- BEUTELMANN, P. (1973). Untersuchungen zur Biosynthese eines Cytokinins in Calluszellen von Laubmoosporophyten. *Planta* 112: 181-190.
- BEUTELMANN, P. & L. BAUER (1977). Purification and identification of a cytokinin from moss callus cells. *Planta* 133: 215-217.

- BEUTELMANN, P., D. J. COVE & E. HARTMANN (1978). Cytokinin producing protonema mutants of the moss *Physcomitrella patens*. *Pl. Physiol.* 61: 48.
- BOPP, M. (1983). Developmental physiology of bryophytes. *New Manual of Bryology (Nichinan)* 1: 276-324.
- BOYD, P. J., J. HALL & D. J. COVE (1988). An airlift fermenter for the culture of the moss *Physcomitrella patens*. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 41-45. Hattori Bot. Lab., Nichinan.
- BÜNNING, E. & D. VON WETTSTEIN (1953). Polarität und Differenzierung an Mooskeimen. *Naturwissenschaften* 40: 147-148.
- CHOPRA, R. N. & S. SOOD (1973). In vitro studies in Marchantiales. I. Effects of some carbohydrates, agar, pH, light and growth regulators on the growth and sexuality in *Riccia crystallina*. *Phytomorph.* 23: 230-244.
- CHOPRA, R. N. & S. C. BHATLA (1981). Effect of physical factors on gametangial induction, fertilization and sporophyte development in the moss *Bryum argenteum* grown in vitro. *New Phytol.* 89: 439-447.
- CHOPRA, R. N. & S. DHINGRA-BABBAR (1986). Studies on callus induction, its growth and differentiation in *Marchantia palmata* Nees. I. Effect of some amino acids, complex organic substances and activated charcoal. *J. Hattori Bot. Lab.* 60: 193-202.
- CHOPRA, R. N. & A. KAPUR (1989). Studies on callus induction in the liverwort *Mannia dichotoma*. *Phytomorph.* 39: 357-362.
- CHOPRA, R. N. & S. KUMRA (1986). Studies on callus induction, its growth and differentiation in the liverwort *Asterella wallichiana* (Lehm. & Lindenb.) Grolle. *Cryptogamie, Bryol. Lichénol.* 7: 249-255.
- CHOPRA, R. N. & S. KUMRA (1988). Callus formation and its differentiation. In: *Biology of Bryophytes*, pp. 119-130, John Wiley & Sons, Inc., New Delhi (India).
- COVE, D. J. & N. W. ASHTON (1984). The hormonal regulation of gametophyte development in bryophytes. In: DYER, A. F. & J. G. DUCKETT (eds.), *The Experimental Botany of Bryophytes*, pp. 177-201, Academic Press Inc. Orlando, Florida, USA.
- DHINGRA-BABBAR, S. & R. N. CHOPRA (1985). Induction of callus in the liverwort *Marchantia palmata* Nees. *J. Bryol.* 13: 585-590.
- DHINGRA-BABBAR, S. & R. N. CHOPRA (1986a). Studies on callus induction, its growth and differentiation in *Marchantia palmata* Nees. I. Effect of some amino acids, complex organic substances and activated charcoal. *J. Hattori Bot. Lab.* 60: 193-202.
- DHINGRA-BABBAR, S. & R. N. CHOPRA (1986b). Studies on callus induction, its growth and differentiation in *Marchantia palmata* Nees. II. Effect of some growth regulators. *Beitr. Biol. Pflanzen* 61: 467-476.
- DOYLE, W. T. (1967). Mosses and liverworts. In: WILT, F. H. & N. K. WESSELS (eds.), *Methods in developmental biology*, pp. 329-339, T.Y. Crowell, New York.
- EULER, R., P. BEUTELMANN & E. HARTMANN (1988). The metabolism of polyunsaturated C<sub>20</sub>-fatty acids in suspension cultures of a moss protonema. *Biological Chemistry Hoppe-Seyler* 369: 18.
- FELIX, H. (1988). Fungi on bryophytes, a review. *Bot. Helv.* 98: 239-269.
- GAY, L. (1976). The development of leafy gametophytes from isolated protoplasts of *Polytrichum juniperinum* Willd. *Z. Pflanzenphysiol.* 79: 33-39.
- GRIMSLEY, N. H. & L. A. WITHERS (1983). Cryopreservation of cultures of the moss *Physcomitrella patens*. *Cryo-Letters* 4: 251-258.
- GRIMSLEY, N. H., N. W. ASHTON & D. J. COVE (1977). The production of somatic hybrids by protoplast fusion in the moss, *Physcomitrella patens*. *Mol. Gen. Genet.* 154: 97-100.
- GRUSAK, M. A., R. J. THOMAS & B. H. MARSH (1980). Wall regeneration by plasmolysed cells from tissue suspension cultures of *Sphaerocarpos donnellii*. *J. Cell Sci.* 43: 167-175.
- HAMADA, H., H. KONISHI, H. J. WILLIAMS & A. I. SCOTT (1991). Biotransformation of testosterone isomers by a green cell suspension culture of *Marchantia polymorpha*. *Phytochem.* 30: 2269-2270.
- HANDA, A. K. & M. N. JOHRI (1976). Cell differentiation by 3',5'-cyclic AMP in a lower plant. *Nature* 259: 480-482.
- HANDA, A. K. & M. N. JOHRI (1977). Cyclic adenosine 3':5'-monophosphate in moss protonema; a comparison of its levels by protein kinase and Gilman assays. *Plant Physiol.* 59: 490-496.
- HANDA, A. K. & M. N. JOHRI (1979). Involvement of cyclic adenosine-3',5'-monophosphate in chloronema differentiation in protonema cultures of *Funaria hygrometrica*. *Planta* 144: 317-324.
- HANSEN, C. E. & P. ROSSI, P. (1991a). Effects of culture conditions on accumulation of arachidonic and eicosapentaenoic acids in cultured cells of *Rhytidadelphus squarrosus* and *Eurhynchium striatum*. *Phytochem.* 30: 1837-1841.
- HANSEN, C. E. & P. ROSSI (1991a). Incorporation and metabolism of radiolabelled linoleic acid in cultures cells of the moss *Rhytidadelphus squarrosus*. *Phytochem.* 30: 2899-2903.
- HARBORNE, J. B. (1990). Role of secondary metabolites in chemical defence mechanisms in plants. In: *Bioactive compounds from plants. Ciba Foundation Symposium* 154: 126-139. John Wiley & Sons Inc., Chichester.
- HARTMANN, E. (1971). Über den Nachweis eines Neurohormones beim Laubmooscallus und seine Beeinflussung durch das Phytochrom. *Planta* 101: 159-165.
- HARTMANN, E. (1972). Über den Allantoinstoffwechsel des Laubmooskallus und seine Beeinflussung durch Licht. *Biochem. Physiol. Pflanzen* 163: 421-431.

- HARTMANN, E. (1973). Über die Wirkung des Phytochroms beim Laubmooskallus. I. Photomorphogenese und Stoffwechsel. *Beitr. Biol. Pflanzen* 49: 1-34.
- HARTMANN, E. (1974). Über die Wirkung des Phytochroms beim Laubmooskallus. II. Sauerstoffmetabolismus und Acetylcholinwirkung. *Z. Pflanzenphysiol.* 71: 349-365.
- HARTMANN, E. & H. KILBINGER (1974). Gas-liquid-chromatographic determination of light-dependent acetylcholine concentrations in moss callus. *Biochem. J.* 137: 249-252.
- HINTIKKA, V. (1972). Variation in gametophyte morphology of Sphagnum fallax in aseptic culture. *Ann. Bot. Fenn.* 9: 91-96.
- HUNECK, S. (1983). Chemistry and biochemistry of bryophytes. *New Manual of Bryology (Nichinan)* 1: 1-116.
- HUNECK, S. (1986). The application of cell cultures in bryology. *Bryol. Times* 38: 1-3.
- ILAHI, I. (1972). Callus formation in some members of the order Metzgeriales (Hepaticae). *Biologia* 18: 125-130.
- ILAHI, I. & A. ALLSOPP (1969). Studies in the Metzgeriales (Hepaticae). II. The effects of certain auxins on the growth and morphology of some thalloid species. *Phytomorph.* 19: 381-389.
- ILAHI, I. & A. ALLSOPP (1970). Studies in the Metzgeriales (Hepaticae). VI. Regeneration and callus formation in some thalloid species. *Phytomorph.* 20: 126-136.
- IMOTO, S. A. & Y. OHTA (1985). Intracellular localization of lunularic acid and prelunularic acid in suspension cultured cells of Marchantia polymorpha. *Plant Physiol.* 79: 751-755.
- JACUBOWSKI, S. & H. RUDOLPH (1989). Glutamate dehydrogenase in Sphagnum species. *J. Hattori Bot. Lab.* 67: 395-398.
- JOHRI, M. M. (1974). Differentiation of caulinema cells by auxins in suspension cultures of Funaria hygrometrica. In: *Plant Growth Substances 1973 Part IV*, pp. 925-933. Tokyo, Hirokawa Publ. Co.
- JOHRI, M. M. (1975). The protonema of Funaria hygrometrica as a system for studying cell differentiation. In: MOHAN RAM, H.Y., J. J. SHAH & C. K. SHAH (eds.), *Form, structure and function in plants*, pp. 116-124, Meerut, India: Sarita Prakashan.
- JOHRI, M. M. & B. M. A. HENRIQUES (1990). Nitrate reductase from the moss Funaria. In: ZINSMEISTER, H. D. & R. MUSS (eds.), *Proc. Phytochem. Soc. Europe 29. Bryophytes: their chemistry and chemical taxonomy*, pp. 265-273. Clarendon Press, Oxford.
- JONES, D. G. (1978). Aspects of growth and development in Sphagnum cuspidatum. *Bull. Br. Bryol. Soc.* 3: 8-10.
- KAJITA, M., S. TAKIO, S. TAKAMI & S. HINO (1987). Establishment and growth characterization of suspension culture of cells from the moss, Sphagnum imbricatum. *Physiol. Plant.* 70: 21-26.
- KAPUR, A. & R. N. CHOPRA (1989). Studies on callus growth and differentiation in the liverwort Mannia dichotoma. *J. Hatt. Bot. Lab.* 67: 323-330.
- KAPUR, A. & R. N. CHOPRA (1991). The effect of some complex organic substances on callus growth in the liverwort Mannia androgyna. *Cryptogamie, Bryol. Lichénol.* 12: 155-164.
- KATOH, K. (1983a). Photosynthesis and photoassimilation of glucose during photomixotrophic growth of Marchantia polymorpha cells in suspension culture. *Physiol. Plant.* 57: 67-74.
- KATOH, K. (1983b). Kinetics of photoautotrophic growth of Marchantia polymorpha cells in suspension culture. *Physiol. Plant.* 59: 242-248.
- KATOH, K. (1988). Isolation and maintenance of callus and cell suspension cultures of bryophytes. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 99-105. Hattori Bot. Lab., Nichinan.
- KATOH, K., Y. OHTA, Y. HIROSE & T. IWAMURA (1979). Photoautotrophic growth of Marchantia polymorpha L. cells in suspension culture. *Planta* 144: 509-510.
- KATOH, K., M. ISHIKAWA, K. MIYAKE, Y. OHTA, Y. HIROSE & T. IWAMURA (1980). Nutrient utilization and requirement under photoheterotrophic growth of Marchantia polymorpha: Improvement of the culture medium. *Physiol. Plant.* 49: 241-247.
- KATOH, K. & R. TAKEDA (1990). Growth and production of secondary metabolites by cultured bryophyte cells. In: ZINSMEISTER, H. D. & R. MUSS (eds.), *Proc. Phytochem. Soc. Europe 29. Bryophytes: their chemistry and chemical taxonomy*, pp. 349-358. Clarendon Press, Oxford.
- KAUL, K. N., G. C. MITRA & B. K. TRIPATHI (1961). Morphogenetic responses of the thallus of Marchantia to several growth substances. *Curr. Sci.* 30: 131-133.
- KAUL, K. N., G. C. MITRA, & B. K. TRIPATHI (1962). Responses of Marchantia in aseptic culture to well-known auxins and antiauxins. *Ann. Bot.* 26: 447-467.
- KNIGHT, C. D., D. J. COVE, P. J. BOYD & N. W. ASHTON (1988). The isolation of biochemical and developmental mutants in Pyscomitrella patens. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 47-58. Hattori Bot. Lab., Nichinan.
- KONNO, H., Y. YAMASAKI & K. KATOH (1983). Exopolygalacturonase from suspension cultures of Marchantia polymorpha. *Plant Physiol.* 73: 216-222.
- KONNO, H., Y. YAMASAKI & K. KATOH (1987). Fractionation and partial characterization of pectic polysaccharides in cell walls from liverwort (Marchantia polymorpha) cell cultures. *J. Exp. Bot.* 38: 711-722.
- KUC, J. (1990). Compounds from plants that regulate or participate in disease resistance. In: *Bioactive compounds from plants. Ciba Foundation Symposium 154*: 213-225. John Wiley & Sons Inc., Chichester.
- KUMRA, S. (1984). Callus induction in the liverwort Asterella wallichiana (L. et L.) Grolle. *J. Plant Physiol.* 115: 263-269.

- KUMRA, P. K. & R. N. CHOPRA (1982). Effect of some growth substances, vitamins and ultraviolet radiation on callus induction in the moss *Bryum coronatum* Schwaegr. *Z. Pflanzenphysiol.* 108: 143-150.
- KUMRA, S. & R. N. CHOPRA (1987). Callus initiation, its growth and differentiation in the liverwort *Asterella wallichiana* (Lehm. et Lindenb.) Grolle I. — Effect of auxins and cytokinins. *J. Hattori Bot. Lab.* 63: 237-245.
- LAL, M. (1961). In vitro production of apogamous sporogonia in *Physcomitrium coorgense* Broth. *Phytomorph.* 11: 263-269.
- LAL, M. (1984). The culture of bryophytes including apogamy, apospory, parthenogenesis and protoplasts. In: DYER, A. F. & J. G. DUCKETT (eds.), *The Experimental Botany of Bryophytes*, pp. 97-115, Academic Press Inc. Orlando, Florida, USA.
- LAW, D. M., D. V. BASILE & M. R. BASILE (1985). Determination of endogenous indole-3-acetic acid in *Plagiochila arctica* (Hepaticae). *Plant Physiol.* 77: 926-929.
- LISA, L. (1971). Nitrogen nutrition of liverwort *Riccia fluitans* in axenic culture. *Acta Universitatis Carolinae — Biologica*: 185-190.
- MACHLIS, L. (1962). The effects of mineral salts, glucose, and light on the growth of the liverwort, *Sphaerocarpos donnellii*. *Physiol. Plant.* 15: 354-362.
- MACHLIS, L. & W. T. DOYLE (1962). Submerged growth of pure cultures of the liverwort, *Sphaerocarpos donnellii*. *Physiol. Plantarum* 15: 351-353.
- MCQUEEN, C. B. (1991). Laboratory and greenhouse cultures and the experimental taxonomy of bryophytes. *Adv. Bryol.* 4: 103-120.
- MAEDA, M. (1979). Isolation and culture of protoplasts from moss protonemata. *Bot. Mag. Tokyo* 92: 105-110.
- MEHRA, K. (1984). Effect of ammonium and nitrate nitrogen on callus induction in *Hyophila involuta* (Musci). *Beitr. Biol. Pflanzen* 59: 333-339.
- MEHRA, P. N. & D. PENTAL (1976). Induction of apospory, callus, and correlated morphogenetic studies in *Athalamia pusilla* Kash. *J. Hattori Bot. Lab.* 40: 151-183.
- MENON, M. K. C. & M. LAL (1972). Influence of sucrose on the differentiation of cells with zygote-like potentialities in a moss. *Naturwissenschaften* 59: 514.
- MENON, M. K. C. & M. LAL (1974). Morphogenetic role of kinetin and abscisic acid in the moss *Pyscomitrium*. *Planta* 115: 319-328.
- MENON, M. K. C. & E. HARTMANN (1988a). Secretion of protein-bound hydroxyproline from moss callus cells. *J. Plant Physiol.* 132: 569-574.
- MENON, M. K. C. & E. HARTMANN (1988b). Extracellular secretion of polysaccharides from moss callus cells. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 107-111. Hattori Bot. Lab., Nichinan.
- MEYER, D. E. (1953). Tumorartige Wucherungen bei einem Lebermoos. *Naturwissenschaften* 40: 297-298.
- MILLER, D. H. & L. MACHLIS (1968). Effects of light on the growth and development of the liverwort, *Sphaerocarpos donnellii* Aust. *Plant Physiol.* 43: 714-722.
- IMITSUBISHI-GAS-CHEM. (1988). Cultivation of bryophyta cell culture — callus culture in pyrroloquinoline quinone culture medium. Patent Number JP 63233783 (Kokai), WPI accession No.: 88-318071 (8845)
- MONTAGUE, M. J. & J. TAYLOR (1971). The effects of glucose, indole-3-acetic acid, and 2,4-dichlorophenoxyacetic acid on the growth of *Sphaerocarpos texanus*. *Bryologist* 74: 18-22.
- MORAIS, R. M. S. C. & H. BECKER (1990). Growth and secondary product formation of in vitro cultures from the liverwort *Reboulia hemisphaerica*. *Z. Naturforsch.* 46c: 28-32.
- NAEF, J. (1975). Dispositif de culture de protonemas de mousse en milieu liquide. *Saussurea* 6: 307-312.
- NAEF, J. & M.-TH. MISSET (1973). Culture de protonemas de *Polytrichum* en milieu liquide. *Ber. Schweiz. Bot. Ges.* 83: 205-212.
- NGUMI, V. W., S. TAKIO & S. TAKAMI (1990). Photoautotrophic growth in cell suspension culture from the liverwort, *Marchantia paleacea* var. diptera. *J. Plant Physiol.* 137: 25-28.
- NOGUCHI, M., T. MATSUMOTO, Y. HIRATA, K. YAMAMOTO, A. KATSUYAMA, A. KATO, S. AZECHI & K. KATO (1977). Improvement of growth rates of plant cell cultures. In: BARZ, W., E. REINHARD & M. H. ZENK (eds.), *Plant Tissue Culture and Its Biotechnological Applications*, pp. 85-94, Springer Verlag, Berlin.
- OHTA, Y., K. KATO & K. MIYAKE (1977). Establishment and growth characteristics of a cell suspension culture of *Marchantia polymorpha* L. with high chlorophyll content. *Planta* 136: 229-232.
- OHTA, Y., M. ISHIKAWA, S. ABE, K. KATO & Y. HIROSE (1981). Growth behavior of a liverwort, *Jungermannia subulata* Evans, in a cell suspension culture. The role of organic acids required for cell growth. *Plant Cell Physiol.* 22: 1533-1540.
- OHTA, Y. & Y. HIROSE (1982). Induction and characteristics of cultured cells from some liverworts of *Jungermanniales*. *J. Hattori Bot. Lab.* 53: 239-244.
- OHTA, Y. & T. TAKATA (1990). Secondary metabolites production in cultured cells of liverworts. In: ZINSMEISTER, H. D. & R. MUES (eds.), *Proc. Phytochem. Soc. Europe* 29. *Bryophytes: their chemistry and chemical taxonomy*, pp. 359-366. Clarendon Press, Oxford.
- OHTA, Y., S. ABE, H. KOMURA & M. KOBAYASHI (1983). Prelunularic acid, a probable immediate precursor of lunularic acid. First example of a "prearomatic" intermediate in the phenylpropanoid-polymalonate pathway. *J. Amer. Chem. Soc.* 105: 4480-4481.
- OHTA, Y., S. ABE, H. KOMURA & M. KOBAYASHI (1984). Prelunularic acid in liverworts. *Phytochem.* 23: 1607-1609.

- OHYAMA, K., L. R. WETTER, Y. YAMANO, H. FUKUZAWA & T. KOMANO (1982). A simple method for isolation of chloroplast DNA from *Marchantia polymorpha* L. cell suspension cultures. *Agric. Biol. Chem.* 46: 237-242.
- OHYAMA, K., Y. YAMANO, H. FUKUZAWA, T. KOMANO, H. YAMAGISHI, S. FUJIMOTO & M. SUGIURA (1983). Physical mappings of chloroplast DNA from liverwort *Marchantia polymorpha* L. cell suspension cultures. *Mol. Gen. Genet.* 189: 1-9.
- ONO, K. (1973). Callus formation in liverwort, *Marchantia polymorpha*. *Jap. J. Genetics* 48: 69-70.
- ONO, K. (1974). Cytological studies on callus of *Marchantia polymorpha*. *Miscellanea Bryologica et Lichenologica* 6: 143.
- ONO, K. (1976a). Cytological observations on the calluses and restored thalluses in *Marchantia polymorpha*. *Jap. J. Genetics* 51: 11-18.
- ONO, K. (1976b). Chromosome variation and redifferentiation of thalluses in *Marchantia* callus subcultured for 4 years. *Miscellanea Bryologica et Lichenologica* 7: 107-108.
- ONO, K., K. OHYAMA & O. L. GAMBORG (1979). Regeneration of the liverwort *Marchantia polymorpha* L. from protoplasts isolated from cell suspension culture. *Plant Sci. Lett.* 14: 225-229.
- ONO, K. & K. OKAMOTO (1984). Isolation and culture of protoplasts from the liverwort cell suspension cultures and the moss protonemata, *J. Hattori Bot. Lab.* 56: 201-207.
- ONO, K., Y. MURASAKI & K. KAWAUCHI (1987). Establishment and characteristics of a cell suspension culture from a moss, *Atrichum undulatum*. *Bot. Mag. Tokyo* 100: 217-221.
- ONO, K., Y. MURASAKI & M. TAKAMIYA (1988). Induction and morphogenesis of cultured cells of bryophytes. *J. Hattori Bot. Lab.* 65: 391-401.
- PADIDAM, M. & M. M. JOHRI (1991). Regulation of nitrate reductase by molybdenum in chloronema cell cultures of the moss *Funaria hygrometrica*. *Plant Sci.* 76: 19-28.
- PADIDAM, M., K. VENKATESWARLU & M. M. JOHRI (1991). Ammonium represses NADPH-nitrate reductase in the moss *Funaria hygrometrica*. *Plant Sci.* 75: 185-194.
- RAHBAR, K. (1982). Influence of organic compounds and activated charcoal on callusing of protonema in the moss *Hyophila involuta*. *Z. Pflanzenphysiol.* 106: 469-473.
- RAHBAR, K. & R. N. CHOPRA (1980). Preliminary studies on callus induction in the moss: *Hyophila involuta*. *Z. Pflanzenphysiol.* 99: 199-205.
- RAHBAR, K. & R. N. CHOPRA (1982). Effect of liquid medium, activated charcoal and pH on the onset of reproductive phase in the moss *Bartramidula bartramoides*. *Z. Pflanzenphysiol.* 106: 185-189.
- RASHID, A. (1971). Callusing and regeneration of cell-aggregates and free-cells of a hepatic: *Asterella angusta* Aust. *Experiencia* 27: 1498-1499.
- RASHID, A. & R. N. CHOPRA (1969). The apogamous sporophytes of *Funaria hygrometrica* and their cultural behaviour. *Phytomorph.* 19: 170-178.
- RESKI, R. & W. O. ABEL (1985). Induction of budding on chloronemata and caulinemata of the moss, *Physcomitrella patens*, using isopentenyl-adenine. *Planta* 165: 354-358.
- RIPETSKIJ, R. T. (1978). Regeneration of isolated protoplasts of the moss *Pottia intermedia*. *Ukr. Bot. Zh.* 35: 467-469.
- RIPETSKIJ, R. T. (1979). On the nature of stable apogamy in diplophase of mosses. *Ukr. Bot. Zh.* 36: 1-5.
- RUDOLPH, H., M. KIRCHHOFF & S. GLIESMANN (1988). Sphagnum culture techniques. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 25-34. Hattori Bot. Lab., Nichinan.
- SARGENT, M. L. (1988). A guide to the axenic culturing of a spectrum of bryophytes. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 17-24. Hattori Bot. Lab., Nichinan.
- SATO, N. & K. KATO, K. (1988). Analysis and biosynthesis of diacylglycerol-N,N,N-trimethylhomoserine in the cells of *Marchantia* in suspension culture. *Plant Sci.* 55: 21-25.
- SAUERWEIN, M. & H. BECKER (1990). Growth, terpenoid production and antibacterial activity of an in vitro culture of the liverwort *Fossumbronia pusilla*. *Planta Med.* 56: 364-367.
- SAXENA, P. K. & A. RASHID (1982). Induction of callus from protonemal cells of the moss *Anoectangium thomsonii* Mitt. *Beitr. Biol. Pflanzen* 57: 301-308.
- SCHNEIDER, M. J., P. D. VOTH & R. F. TROXLER (1967). Methods of propagating bryophyte plants, tissues and propagules. *Bot. Gaz. (Chicago)* 128: 169-174.
- SHARMA, S., R. K. JAYASWAL & M. M. JOHRI (1979). Cell-density-dependent changes in the metabolism of chloronema cell cultures. *Plant Physiol.* 64: 154-158.
- SIMOLA, L. K. (1969). The effect of various mono- and disaccharides on the growth of *Sphagnum nemoreum* thalli in sterile cultures. *Physiol. Plant.* 22: 1079-1084.
- SIMOLA, L. K. (1975). The effect of several protein amino acids and some inorganic nitrogen sources on the growth of *Sphagnum nemoreum*. *Physiol. Plant.* 35: 194-199.
- SIMOLA, L. K. (1977). The tolerance of *Sphagnum fimbriatum* towards lead and cadmium. *Ann. Bot. Fenn.* 14: 1-5.
- SIMOLA, L. K. (1979). Dipeptide utilization by *Sphagnum fimbriatum*. *J. Hattori Bot. Lab.* 46: 49-54.
- SIMON, E. (1988). Axenic cultures of *Sphagnum* on solid substrates. In: GLIME, J. M. (ed.), *Methods in Bryology, Proc. Bryol. Meth. Workshop, Mainz*, pp. 1-16. Hattori Bot. Lab., Nichinan.

- SOOD, S. (1974). In vitro studies in Marchantiales — II. Effect of mineral nutrients, chelates and organic nitrogenous sources on the growth and sexuality in *Riccia crystallina*. *Phytomorph.* 24: 186-197.
- SPIESS, L. D. (1975). Comparative activity of isomers of zeatin and ribosyl-zeatin on *Funaria hygrometrica*. *Plant Physiol.* 55, 583-585.
- SPIESS, L. D. (1976). Developmental effects of zeatin, ribosyl-zeatin, and *Agrobacterium tumefaciens* B6 on certain mosses. *Plant Physiol.* 58: 107-109.
- SPIESS, L. D. (1979). Antagonism of cytokinin induced callus in *Pylaisiella selwynii* by nucleosides and cyclic nucleotides. *Bryologist* 82: 47-53.
- SPIESS, L. D., B. B. LIPPINCOTT & J. A. LIPPINCOTT (1973). Effect of hormones and vitamine B<sub>12</sub> on gametophore development in the moss *Pylaisiella selwynii*. *Amer. J. Bot.* 60: 708-716.
- SPIESS, L. D., J. C. TURNER, P. G. MAHLBERG, B. B. LIPPINCOTT & J. A. LIPPINCOTT (1977a). Adherence of agrobacteria to moss protonema and gametophores viewed by scanning electron microscopy. *Amer. J. Bot.* 64: 1200-1208.
- SPIESS, L. D., B. B. LIPPINCOTT & J. A. LIPPINCOTT (1977b). Comparative response of *Pylaisiella selwynii* to Agrobacterium and Rhizobium species. *Bot. Gaz.* 138: 35-40.
- SPIESS, L. D., B. B. LIPPINCOTT & J. A. LIPPINCOTT (1981). Bacteria isolated from moss and their effect on moss development. *Bot. Gaz.* 142: 512-518.
- SPJUT, R. W., M. SUFFNESS, G. M. CRAGG & D. H. NORRIS (1986). Mosses, liverworts, and hornworts screened for antitumor agents. *Economic Botany* 40: 310-338.
- SPOERLE, J., H. BECKER, N. S. ALLEN & M. P. GUPTA (1990). Occurrence of (-)-geosmin and other terpenoids in an axenic culture of the liverwort *Symphygyna bronniartii*. *Z. Naturforsch.* 46c: 183-188.
- STAHLY, N. C., J. N. TAMBS & J. TAYLOR (1977). An assessment of fructose as a carbohydrate source of carbon for the in vitro growth of *Sphaerocarpos texanus*. *J. Hattori Bot. Lab.* 43: 69-71.
- SUGAWARA, Y., K. MORI, H. MATSUSHIMA & M. TAKEUCHI (1983). Enhancement of cell division in *Marchantia* protoplast culture by activated charcoal. *Z. Pflanzenphysiol.* 109: 275-278.
- SUSSMAN, M. R. & H. KENDE (1977). The synthesis and biological properties of 8-azido-N<sup>6</sup>-benzyladenine, a potential photoaffinity reagent for cytokinin. *Planta* 137: 91-96.
- SZWEJKOWSKA, A., I. GUZOWSKA & J. GALLAS (1968). Studies on the activity of kinetin in cultures of *Funaria hygrometrica*. II. Effect of kinetin on isolated protonema cells. *Acta Soc. Bot. Pol.* 37: 201-206.
- TAKAMI, S., M. YASUNAGA, S. TAKIO, J. KIMURA & S. HINO (1988a). Establishment of suspension cultures of cells from the hornwort, *Anthoceros punctatus* L. *J. Hattori Bot. Lab.* 64: 429-435.
- TAKAMI, S., S. NISHIKAWA, S. TAKIO & S. HINO (1988b). Photoautotrophic growth in suspension culture of cells from the moss, *Barbula unguiculata*. *Physiol. Plant.* 73: 227-231.
- TAKAMI, S., T. NISHIZAKI, M. HARADA, V. W. NGUMI & S. TAKIO (1990). Glutamate dehydrogenase and glutamine synthetase in *Sphagnum* species. *J. Hattori Bot. Lab.* 68: 441-447.
- TAKEDA, R. & K. KATO (1981). Growth and sesquiterpenoid production by *Calypogeia granulata* Inoue cells in suspension culture. *Planta* 151: 525-530.
- TAKEDA, R. & K. KATO (1983a). Sesquiterpenoids in cultured cells of liverwort, *Calypogeia granulata* Inoue. *Bull. Chem. Soc. Japan* 56: 1265-1266.
- TAKEDA, R. & K. KATO (1983b). 3,10-Dihydro-1,4-dimethylazulene, a labile biosynthetic intermediate isolated from cultured cells of liverwort *Calypogeia granulata* Inoue. *J. Am. Chem. Soc.* 105: 4056-4058.
- TAKIO, S. (1987). Coenzyme requirements of nitrate reductase in extracts from suspension cultured cells of four bryophyte species. *J. Hattori Bot. Lab.* 62: 269-280.
- TAKIO, S. (1989). Effect of cytokinin on morphological changes of suspension cultured cells of the moss, *Barbula unguiculata*. *Plant Cell Reports* 7: 603-606.
- TAKIO, S. (1990). Nitrate reductase inactivator in a chlorate resistant mutant of the liverwort, *Marchantia paleacea* var. diptera. *J. Plant Physiol.* 136: 30-34.
- TAKIO, S., M. KAJITA, S. TAKAMI & S. HINO (1986). Establishment and growth characterization of suspension cultures of cells from *Barbula unguiculata*. *J. Hattori Bot. Lab.* 60: 407-417.
- TAKIO, S., S. TAKAMI & S. HINO (1988). Photosynthetic ability of dark-grown *Marchantia paleacea* cell suspension culture. *J. Plant Physiol.* 132: 195-198.
- TAKIO, S. & S. HINO (1988). Nitrate reductase from suspension cultured cells of *Marchantia polymorpha* L. *J. Plant Physiol.* 132: 470-473.
- TAKIO, S. & T. NAGATA (1990). Changes in chloroplast DNA levels during the culture of photosynthetically active cells of a moss, *Barbula unguiculata*. *J. Plant Physiol.* 137: 147-150.
- TAKIO, S., C. AKITA, V. M. NGUMI & S. TAKAMI (1990). Photosynthetic ability in dark-grown *Reboulia hemisphaerica* and *Barbula unguiculata* cells in suspension culture. *Plant Cell Reports* 8: 575-578.
- TANAKA, A., Y. YAMANO, H. FUKUZAWA, K. OHYAMA & T. KOMANO (1984). In vitro DNA synthesis by chloroplasts isolated from *Marchantia polymorpha* L. cell suspension cultures. *Agric. Biol. Chem.* 48: 1239-1244.
- TRIPATHI, B. K. (1978a). Culture in vitro de tissus et de protonémas provenant de thalles de *Marchantia nepalensis* L. et L. *C. R. Acad. Sc. Paris Série D* 286: 851-854.

- TRIPATHI, B. K. (1978b). Etudes sur l'histogenèse de trois types de cals provenant de thalles de *Marchantia nepalensis* L. et L. *Rev. Gén. Bot.* 85: 91-106.
- ULYCHNA, K. O. (1971). Characteristics of morphogenesis in regenerates of moss sporogonia. *Ontogenesis* 2: 165-169.
- VASHISTHA, B. D. & R. N. CHOPRA (1989). Effect of some physical factors on vegetative growth and archegonial formation in the female clone of *Riccia frostii* Aust. *Phytomorph.* 39: 141-147.
- WANG, T. L., D. J. COVE, P. BEUTELMANN & E. HARTMANN (1980). Isopentenyladenine from mutants of the moss, *Physcomitrella patens*. *Phytochem.* 19: 1103-1105.
- WANG, T. L., R. HORGAN & D. COVE (1981a). Cytokinins from the moss *Physcomitrella patens*. *Plant Physiol.* 68: 735-738.
- WANG, T. L., P. BEUTELMANN & D. COVE (1981b). Cytokinin biosynthesis in mutants of the moss *Physcomitrella patens*. *Plant Physiol.* 68: 739-744.
- WARD, M. (1960a). Some techniques in the culture of mosses. *Bryologist* 63: 213-217.
- WARD, M. (1960b). Callus tissues from the mosses *Polytrichum* and *Atrichum*. *Science* 132: 1401-1402.
- WARD, M. & S. E. FREDERICK (1967). Propagation of aberrant gametophytes from aggregates and from single cell of moss callus. *Phytomorph.* 17: 371-374.
- WETTSTEIN, D. VON (1953). Beeinflussung der Polarität und undifferenzierte Gewebebildung aus Moossporen. *Z. Botanik* 41: 199-226.
- WURZEL, G. & H. BECKER (1990). Growth and terpenoid production of an axenic culture from the liverwort *Ricciocarpus natans*. *Z. Naturforsch.* 45c: 13-18.
- WURZEL, G., H. BECKER, TH. EICHER & K. TIEFENSEE (1990). Molluscicidal properties of constituents from the liverwort *Ricciocarpos natans* and of synthetic lunularic acid derivatives. *Planta Medica* 56: 444-445.
- ZINSMEISTER, H. D. & R. MUES (eds.) (1990). *Proc. Phytochem. Soc. Europe* 29. *Bryophytes: their chemistry and chemical taxonomy*, 470 pp. Clarendon Press, Oxford.
- ZINSMEISTER, H. D., H. BECKER & T. EICHER (1991). Moose, eine Quelle biologisch aktiver Naturstoffe? *Angew. Chemie* 103: 134-151