

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	49 (1994)
Heft:	1
Artikel:	Floristic notes from aquatic stands of Central Greece (Stereia Ellas)
Autor:	Koumpli-Sovantzi, Lemonia / Vallianatou, Irini
DOI:	https://doi.org/10.5169/seals-879538

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Floristic notes from aquatic stands of Central Greece (Stereia Ellas)

LEMONIA KOUMPLI-SOVANTZI
&
IRINI VALLIANATOU

ABSTRACT

KOUMPLI-SOVANTZI, L. & I. VALLIANATOU (1994). Floristic notes from aquatic stands of Central Greece (Stereia Ellas). *Candollea* 49: 195-207. In English, English and French abstracts.

This paper lists a total of 164 taxa of vascular plants (161 flowering plants, 3 ferns) found in and around some aquatic ecosystems of Central Greece, some of which are investigated for the first time.

RÉSUMÉ

KOUMPLI-SOVANTZI, L. & I. VALLIANATOU (1994). Notes floristiques des stations humides de la Grèce centrale. *Candollea* 49: 195-207. En anglais, résumé anglais et français.

164 taxa de plantes vasculaires (161 plantes à fleur, 3 fougères) de stations humides ont été récoltés en Grèce centrale. Quelques-unes d'entre elles sont étudiées pour la première fois.

KEY-WOROS: Floristics — Aquatic habitats — Central Greece.

Introduction

In C. Greece a limited number of aquatic ecosystems (lakes, rivers, rivulets, streams, canals, springs, lagoons, pools, marshes, saltworks) is found. The present floristic study deals with the plants living in the water or on the shores and banks of some of them.

Some parts of C. Greece are well or reasonably well explored areas and some others ill-explored (GREUTER & al., 1976). Its aquatic ecosystems must be considered as not completely investigated, despite the fact that recently some of them have been the object of several publications.

Going from the west to the east, the vegetation of the Mesolongi lagoon has been treated by WOLFF (1968), the aquatic flora of the province of Aitolias & Akarnanias by KOUMPLI-SOVANTZI (1983, 1989), the vegetation of the lake Lisiimakhia by KOUMPLI-SOVANTZI & VALLIANATOU (1985), the vegetation of the delta of the Akheloos river by SEVERIN & al. (1989-1990), the flora of Sperkhios valley by STEPHANIDES (1948), the evolution of the halophytic vegetation of the two largest plains of Attiki (Athens and Marathon) by ECONOMIDOU (1975), whilst the Marathon plain (marsh of Skhinia) has been also dealt with the following publications: HARITONIDOU (1977, 1983), PHITOS (1988) and BROFAS & KARETSOS (1992).

The purpose of this study is to contribute to the floristic knowledge of the aquatic habitats of Central Greece, completing the floristic lists of the already investigated ecosystems or giving for the first time lists of the rest.

Between 1983 and 1992 the authors paid sixteen short visits to the investigated ecosystems; during these visits plant specimens were collected and some measurements of the salinity, conductivity and pH were taken. The authors have in mind to continue this study, visiting other ecosystems of the area in the near future.

List of localities

The aquatic ecosystems under study are found at the Provinces of Aitolias & Akarnanias, Fokidhos, Fthiotidhos, Voiotias and Attikis. The investigated localities are indicated with dots and numbers on the map (Fig. 1) and are shortly described below.




Fig. 1. — Map of Sterea Ellas. Dots with numbers indicate the investigated sites.

I. *Nomos Aitolias & Akarnanias*

1. Mesolongi: saltworks (canals, ditches and pans separated by banks).

II. *Nomos Fokidhos*

2. Kira: rivulet (width = 4 m, depth = 0.5 m; 13.10.1990: salinity = 6‰, conductivity = 8000 µmhos, pH = 7.9; 23.8.1992: salinity 8‰, conductivity = 15 000 µmhos, pH = 7.0); wetland.

III. *Nomos Fthiotidhos*

3. Sperkhios river: bank of the river, near the bridge on the National road Athens — Lamia.

IV. *Nomos Voiotias*

4. Iliki-Paralimni: canal which connects lake Iliki and lake Paralimni (29.8.1992: salinity = 0.5‰, pH = 8.4, conductivity = 400 µmhos).
5. Lake Paralimni (29.8.1992: salinity = 0.8‰, pH 8.6, conductivity = 1150 µmhos).

V. *Nomos Attikis*

6. Oropos: vernal coastal pool becoming quite dry in summer (29.5.1992: salinity = 12‰, conductivity = 17 500 µmhos, pH = 7.2).
7. Loutsa: streamlet of water flowing towards the Vraonas bay.
8. Marathon: technical lake (29.9.1992: salinity = 0.5‰, conductivity = 780 µmhos, pH = 7.9); rivulet (29.9.1992: salinity = 0.2‰, conductivity = 600 µmhos, pH = 7.7).
9. Skhinias: canal (width = 4 m, depth = 1.5 m, 29.9.1992: salinity = 2‰, conductivity = 2600 µmhos, pH = 7.4); marsh; pools.

Species list

The following catalogue of taxa is based on the authors' collections (Herb. L. Koumpli-Sovantzi and I. Vallianatou, duplicates in ATHU). The collectors' name are abbreviated by the capital letters *K. S.* and *V.* respectively and followed by the collecting numbers appear on the list after the localities and the corresponding collecting dates. When no collecting number is cited the report is based on field notes by the authors.

The determinations were mainly made according to CASPER & KRAUSCH (1980-1981), COOK & al. (1974), DAVIS (1965-1988), FASSETT (1957) and TUTIN & al. (1964-1980). Nomenclature follows that of TUTIN & al. (1964-1980) except for the part revised by GREUTER & al. (1984, 1986, 1989) where the latter is followed.

P TERIDOPHYTA*Equisetaceae*

Equisetum arvense L. — Sperkhios river (9.5.1983: *K. S. 1851*).

Equisetum ramosissimum Desf. — Marathon (29.9.1992: *K. S. 4581a*).

Equisetum telmateia Ehrh. — Marathon (29.9.1992: *K. S. 4581*).

A NGIOSPERMATE — DICOTYLEDONES*Salicaceae*

Salix alba L. — Sperkhios river (9.5.1983: *K. S. 1853*); Marathon (29.9.1992: *K. S. 4584*).

Populus alba L. — Sperkhios river (9.5.1983: *K. S. 1859*); Marathon (29.9.1992: *K. S. 4589*).

Polygonaceae

Polygonum aviculare L. — Mesolongi (3.6.1991: *K. S. 4413*).

Persicaria lapathifolia (L.) S. F. Gray — Iliki-Paralimni (29.8.1992: *K. S. 4541*); Marathon (29.9.1992: *K. S. 4591*).

Rumex conglomeratus Murray — Mesolongi (3.6.1991: *K. S. 4411*).

Rumex cf. obtusifolius L. subsp. *transiens* (Simkovics) Rech. fil. — Mesolongi (3.6.1991: K. S. 4412).

Rumex palustris Sm. — Iliki-Paralimni (29.8.1992: K. S. 4540a).

Chenopodiaceae

Beta maritima L. — Mesolongi (3.6.1991: K. S. 4422).

Chenopodium ambrosioides L. — Sperkhios river (9.5.1983: K. S. 1855).

Chenopodium botrys L. — Mesolongi (3.6.1991: K. S. 4423).

Atriplex portulacoides L. — Mesolongi (3.6.1991: K. S. 4416); Oropos (27.4.1992: V. 5904, 29.8.1992: K. S. 4561).

Atriplex prostrata DC. — Mesolongi (3.6.1991: K. S. 4415, 20.10.1991: K. S. 4450).

Halocnemum strobilaceum (Pallas) MB. — Mesolongi (20.10.1991: K. S. 4448).

Arthrocnnum macrostachyum (Moric.) Moris — Mesolongi (3.6.1991: K. S. 4424); Skhinias (10.11.1991: K. S. 4476); Oropos (29.8.1992: K. S. 4557).

Sarcocornia perennis (Miller) A. J. Scott — Mesolongi (20.10.1991: K. S. 4452); Skhinias (10.11.1991: K. S. 4475, 29.8.1992: K. S. 4626); Oropos (27.4.1992: V. 5916, 29.9.1992: K. S. 4558).

Salicornia europaea L. — Mesolongi (3.6.1991: K. S. 4425).

Suaeda maritima (L.) Dumort. — Mesolongi (20.10.1991: K. S. 4453); Oropos (29.8.1992: K. S. 4560).

Suaeda vera J. F. Gmelin — Skhinias (29.9.1992: K. S. 4627).

Salsola soda L. — Mesolongi (3.6.1991: K. S. 4426, 20.10.1991: K. S. 4451); Oropos (29.8.1992: K. S. 4559).

Amaranthaceae

Amaranthus albus L. — Sperkhios river (9.5.1983: K. S. 1846).

Amaranthus retroflexus L. — Sperkhios river (9.5.1983: K. S. 1856).

Caryophyllaceae

Herniaria hirsuta L. — Skhinias (26.5.1988: K. S. 3513).

Spergularia salina J. Presl & C. Presl — Skhinias (26.5.1988: K. S. 3509); Oropos (27.4.1992: V. 5908); Mesolongi (3.6.1991: K. S. 4417).

Silene colorata Poiret — Mesolongi (3.6.1991: K. S. 4419).

Silene gallica L. — Mesolongi (3.6.1991: K. S. 4420).

Petrorhagia velutina (Guss.) P. W. Ball & Heywood — Mesolongi (3.6.1991: K. S. 4421).

Velezia rigida L. — Mesolongi (3.6.1991: K. S. 4418).

Ceratophyllaceae

Ceratophyllum demersum L. — Paralimni (15.7.1983: K. S. 1895).

Ranunculaceae

Nigella damascena L. — Mesolongi (3.6.1991: K. S. 4414).

Ranunculus sceleratus L. — Iliki-Paralimni (29.8.1992: K. S. 4542).

Papaveraceae

Fumaria capreolata L. — Oropos (27.4.1992: V. 5900).

Cruciferae

Hymenolobus procumbens (L.) Nutt. — Oropos (27.4.1992: V. 5915).

Sinapis alba L. subsp. *alba* — Oropos (27.4.1992: V. 5905).

Platanaceae

Platanus orientalis L. — Sperkhios river (9.5.1983: K. S. 1845); Marathon (29.9.1992: K. S. 4592).

Rosaceae

Rubus sp. — Sperkhios river (9.5.1983: K. S. 1847).

Potentilla reptans L. — Marathon (29.9.1992: K. S. 4576).

Leguminosae

Ononis reclinata L. — Mesolongi (3.6.1991: K. S. 4408).

Melilotus indicus (L.) All. — Oropos (27.4.1992: V. 5906).

Melilotus neapolitanus Ten. — Mesolongi (3.6.1991: K. S. 4407).

Melilotus segetalis (Brot.) Ser. — Mesolongi (3.6.1991: K. S. 4406).

Medicago lupulina L. — Marathon (29.9.1992: K. S. 4577).

Trifolium angustifolium L. — Mesolongi (3.6.1991: K. S. 4402).

Trifolium campestre Schreber — Mesolongi (3.6.1991: K. S. 4403).

Trifolium infamia-ponertii Greuter — Skhinias (26.5.1988: K. S. 3517).

Trifolium nigrescens Viv. — Skhinias (26.5.1988: K. S. 3522); Mesolongi (3.6.1991: K. S. 4405).

Trifolium pallidum Waldst. & Kit. — Mesolongi (3.5.1991: K. S. 4404).

Trifolium resupinatum L. — Skhinias (26.5.1988: K. S. 3514).

Scorpiurus muricatus L. — Mesolongi (3.6.1991: K. S. 4401).

Euphorbiaceae

Euphorbia oblongata Griseb. — Marathon (29.9.1992: K. S. 4582).

Euphorbia prostrata Aiton — Sperkhios river (9.5.1983: K. S. 1843).

Euphorbia hirsuta Vahl — Mesolongi (3.6.1991: K. S. 4400).

Guttiferae

Hypericum perforatum L. — Marathon (29.9.1992: K. S. 4586).

Tamaricaceae

Tamarix hampeana Boiss. & Heldr. — Skhinias (26.5.1988: K. S. 3521).

Tamarix smyrnensis Bunge — Marathon (29.9.1992: K. S. 4598).

Frankeniaceae

Frankenia pulverulenta L. — Mesolongi (3.6.1991: K. S. 4410).

Lythraceae

Lythrum junceum Banks & Solander — Mesolongi (20.10.1991: K. S. 4449).

Onagraceae

Epilobium hirsutum L. — Iliki-Paralimni (29.8.1992: K. S. 4543); Marathon (29.9.1992: K. S. 4601).

Haloragaceae

Myriophyllum spicatum L. — Paralimni (15.7.1983: K. S. 1894, 29.8.1992: K. S. 4551); Marathon (29.9.1992: K. S. 4602).

Umbelliferae

Foeniculum vulgare Miller subsp. *piperitum* (Ucria) Coutinho — Skhinias (29.9.1992: K. S. 4608).

Bupleurum fruticosum L. — Marathon (29.9.1992: K. S. 4596).

Bupleurum gracile D'Urv. — Mesolongi (3.6.1991: K. S. 4399).

Apium nodiflorum (L.) Lag. — Skhinias (10.11.1991: K. S. 4483, 29.9.1992: K. S. 4604); Marathon (29.9.1992: K. S. 4584).

Torilis nodosa (L.) Gaertner — Mesolongi (3.6.1991: K. S. 4398).

Primulaceae

Anagallis arvensis L. — Mesolongi (3.6.1991: K. S. 4397); Oropos (27.4.1992: V. 5903).

Samolus valerandi L. — Marathon (29.9.1992: K. S. 4583); Skhinias (29.9.1992).

Plumbaginaceae

Limonium narbonense Miller — Mesolongi (20.10.1991: K. S. 4457); Skhinias (29.9.1992: K. S. 4616).

Limonium virgatum (Willd.) Fourr. — Skhinias (29.9.1992: K. S. 4624).

Gentianaceae

Centaurium spicatum (L.) Fritsch — Skhinias (10.11.1991: K. S. 4470).

Centaurium tenuiflorum (Hoffmanns. & Link) Fritsch subsp. *acutiflorum* (Schott) Zeltner — Mesolongi (3.6.1991: K. S. 4396); Marathon (29.9.1992: K. S. 4575).

Apocynaceae

Nerium oleander L. — Sperkhios river (9.5.1983: K. S. 1848); Marathon (29.9.1992: K. S. 4593).

Convolvulaceae

Cressa cretica L. — Mesolongi (20.10.1991: K. S. 4455).

Calystegia sepium (L.) R. Br. subsp. *sepium* — Skhinias (29.9.1992: K. S. 4611).

Boraginaceae

Anchusa hybrida Ten. — Skhinias (26.5.1988: K. S. 3518).

Verbenaceae

Vitex agnus-castus L. — Mesolongi (3.6.1991); Marathon (29.9.1992: K. S. 4597).

Verbena officinalis L. — Marathon (29.9.1992: K. S. 4588).

Lippia nodiflora (L.) Michx — Iliki-Paralimni (29.8.1992: K. S. 4549).

Callitrichaceae

Callitriche truncata Guss. subsp. *truncata* — Loutsa (5.11.1990: K. S. 4231).

Labiatae

Lycopus europaeus L. — Sperkhios river (9.5.1983: K. S. 1841); Iliki-Paralimni (29.8.1992: K. S. 4548).

Mentha aquatica L. — Marathon (29.9.1992: K. S. 4580).

Mentha pulegium L. — Iliki-Paralimni (29.8.1992: K. S. 4547); Marathon (29.9.1992: K. S. 4579).

Solanaceae

Solanum luteum Miller subsp. *alatum* (Moench) Dostal — Sperkhios river (9.5.1983: K. S. 1842).

Solanum nigrum L. — Sperkhios river (9.5.1983: K. S. 1838).

Scrophulariaceae

Veronica anagalloides Guss. — Iliki-Paralimni (29.8.1992: K. S. 4546).

Plantaginaceae

Plantago coronopus L. — Skhinias (26.5.1988: K. S. 3503a); Oropos (27.4.1992: V. 5913); Mesolongi (3.6.1991: K. S. 4391).

Plantago crassifolia Forskål — Skhinias (26.5.1988: K. S. 3503, 10.11.1991: K. S. 4474, 29.9.1992: K. S. 4615).

Plantago lagopus L. — Skhinias (26.5.1988: K. S. 3516); Mesolongi (3.6.1991: K. S. 4392).

Plantago major L. (s.l.) — Marathon (29.9.1992: K. S. 4590).

Caprifoliaceae

Sambucus nigra L. — Sperkhios river (9.5.1983: K. S. 1860).

Compositae

Aster squamatus (Sprengel) Hieron. — Sperkhios river (9.5.1983: K. S. 1836); Iliki-Paralimni (29.8.1992: K. S. 4550); Marathon (29.9.1992: K. S. 4599); Skhinias (29.9.1992: K. S. 4605).

Aster tripolium L. — Kira (5.5.1991: K. S. 4346); Mesolongi (20.10.1991: K. S. 4458); Skhinias (10.11.1991: K. S. 4468).

Conyza albida Sprengel — Marathon (29.9.1992: K. S. 4600).

Conyza canadensis (L.) Cronq. — Sperkhios river (9.5.1983: K. S. 1839).

- Inula crithmoides* L. — Mesolongi (3.6.1991); Skhinias (29.9.1992: K. S. 4625).
- Dittrichia viscosa* (L.) W. Greuter subsp. *viscosa* — Skhinias (29.9.1992: K. S. 4618).
- Xanthium strumarium* L. subsp. *italicum* (Moretti) D. Löve — Marathon (29.9.1992: K. S. 4587).
- Chamomilla recutita* (L.) Rauschert — Oropos (27.4.1992: V. 5914).
- Galactites tomentosa* Moench — Mesolongi (3.6.1991: K. S. 4395).
- Hedypnois cretica* (L.) Dum.-Courset — Skhinias (26.5.1988: K. S. 3512).
- Picris echioides* L. — Sperkhios river (9.5.1983: K. S. 1837); Mesolongi (3.6.1991: K. S. 4393).
- Tragopogon brevirostris* DC. — Skhinias (10.11.1991: K. S. 4467).
- Crepis foetida* L. subsp. *rhoeadifolia* (Bieb.) Čelak. — Mesolongi (3.6.1991: K. S. 4394).

ANGIOSPERMAE — MONOCOTYLEDONES

Alismataceae

Alisma lanceolatum With. — Skhinias (20.4.1991: K. S. 4233, specimen without flowers).

Juncaginaceae

Triglochin bulbosa L. subsp. *barrelieri* (Loisel.) Rouy — Oropos (27.4.1992: V. 5917, 29.8.1992: K. S. 4554).

Potamogetonaceae

Potamogeton crispus L. — Paralimni (15.7.1983: K. S. 1896).

Potamogeton nodosus Poiret — Skhinias (26.5.1988: K. S. 3530, 10.11.1991: K. S. 4481); Paralimni (15.7.1983: K. S. 1892).

Potamogeton pectinatus L. — Skhinias (26.5.1988: K. S. 3531, 10.11.1991: K. S. 4479).

Ruppiaceae

Ruppia cirrhosa (Petagna) Grande — Mesolongi (20.10.1991: K. S. 4460).

Ruppia maritima L. — Kira (30.7.1990: K. S. 4209, 13.10.1990: K. S. 4228); Oropos (27.4.1992: V. 5898).

Zosteraceae

Zostera noltii Hornem. — Mesolongi (20.10.1991: K. S. 4461).

Zannichelliaceae

Zannichellia palustris L. — Skhinias (26.5.1988: K. S. 3529, 10.11.1991: K. S. 4482).

Cymodocea nodosa (Ucria) Ascherson — Mesolongi, (20.10.1991: K. S. 4462).

Juncaceae

Juncus acutus L. — Skhinias (26.5.1988: K. S. 3525, 10.11.1991: K. S. 4465); Mesolongi (3.6.1991: K. S. 4389).

Juncus articulatus L. — Sperkhios river (9.5.1983: K. S. 1870); Iliki-Paralimni (29.8.1992: K. S. 4533); Skhinias (29.9.1992: K. S. 4612).

Juncus bufonius L. — Skhinias (26.5.1988: K. S. 3510).

Juncus heldreichianus Marsson — Skhinias (10.11.1991: K. S. 4466, 29.9.1992: K. S. 4617).

Juncus littoralis C. A. Meyer — Skhinias (29.9.1992: K. S. 4614):

Juncus maritimus Lam. — Skhinias (10.11.1991: K. S. 4473); Oropos (27.4.1992: V. 5901, 29.8.1992: K. S. 4552).

Juncus subulatus Forskål — Skhinias (26.5.1988: K. S. 3507, 20.4.1991: K. S. 4235, 10.11.1991: K. S. 4478, 29.9.1992: K. S. 4621); Oropos (29.8.1992: K. S. 4553); Mesolongi (3.6.1991: K. S. 4388).

Gramineae

Lolium multiflorum Lam. — Mesolongi (3.6.1991: K. S. 4386).

Lolium rigidum Gaudin subsp. *lepturoides* (Boiss.) Sennen & Mauricio — Skhinias (26.5.1988: K. S. 3505).

Sphenopus divaricatus (Gouan) Reichenb. — Mesolongi (3.6.1991: K. S. 4387); Oropos (27.4.1992: V. 5907).

Puccinellia distans (L.) Parl. — Oropos (27.4.1992: V. 5902).

Puccinellia festuciformis (s.l.) — Mesolongi (3.6.1991: K. S. 4378); Skhinias (10.11.1991: K. S. 4469, 29.9.1992: K. S. 4623).

Cynosurus echinatus L. — Skhinias (26.5.1988: K. S. 3519).

Bromus intermedius Guss. — Mesolongi (3.6.1991: K. S. 4379).

Elymus elongatus (Host) Runemark subsp. *ponticus* (Podp.) Melderis — Mesolongi (20.10.1991: K. S. 4459).

Elymus flaccidifolius (Boiss. & Heldr.) Melderis — Skhinias (29.9.1992: K. S. 4619).

Elymus repens (L.) Gould — Marathon (29.9.1992: K. S. 4595).

Hordeum marinum Hudson — Mesolongi (3.6.1991: K. S. 4380); Oropos (27.4.1992: V. 5909).

Hordeum murinum L. subsp. *leporinum* (Link) Arcangeli — Skhinias (26.5.1988: K. S. 3515); Mesolongi (3.6.1991: K. S. 4381).

Gaudinia fragilis (L.) Beauv. — Mesolongi (3.6.1991: K. S. 4382).

Lophochloa cristata (L.) Hyl. — Skhinias (26.5.1988: K. S. 3511).

Lagurus ovatus L. — Skhinias (26.5.1988: K. S. 3523).

Agrostis stolonifera L. — Marathon (29.9.1992: K. S. 4574).

Polypogon maritimus Willd. subsp. *maritimus* — Mesolongi (3.6.1991: K. S. 4374).

Polypogon monspeliensis (L.) Desf. — Skhinias (26.5.1988: K. S. 3504); Mesolongi (3.6.1991: K. S. 4373); Iliki-Paralimni (29.8.1992: K. S. 4538).

Phleum subulatum (Savi) Ascherson & Graebner subsp. *subulatum* — Mesolongi (3.6.1991: K. S. 4377).

Parapholis filiformis (Roth) C. E. Hubbard — Mesolongi (3.6.1991: K. S. 4383).

Parapholis incurva (L.) C. E. Hubbard — Mesolongi (3.6.1991: K. S. 4384).

Parapholis marginata Runemark — Oropos (27.4.1992: V. 5912).

Phalaris minor Retz. — Mesolongi (3.6.1991: K. S. 4375); Oropos (27.4.1992: V. 9II).

Phalaris paradoxa L. — Mesolongi (3.6.1991: K. S. 4376).

Stipa capensis Thunb. — Mesolongi (3.6.1991: K. S. 4372).

Phragmites australis (Cav.) Trin. ex Steudel — Sperkhios river (9.5.1983: K. S. 1840); Mesolongi (3.6.1991); Iliki-Paralimni (29.8.1992: K. S. 4530); Skhinias (29.9.1992: K. S. 4609); Marathon (29.9.1992: K. S. 4578).

Aeluropus littoralis (Gouan) Parl. — Mesolongi (20.10.1991: K. S. 4456); Oropos (29.8.1992: K. S. 4555); Skhinias (29.9.1992: K. S. 4620).

Crypsis alopecuroides (Piller & Mitterp.) Schrader — Iliki-Paralimni (29.8.1992: K. S. 4539).

Echinochloa crus-galli (L.) Beauv. — Iliki-Paralimni (29.8.1992: K. S. 4535).

Digitaria sanguinalis (L.) Scop. — Iliki-Paralimni (29.8.1992: K. S. 4534).

Paspalum distichum L. — Sperkhios river (9.5.1983: K. S. 1868); Iliki-Paralimni (29.8.1992: K. S. 4537).

Typhaceae

Typha domingensis (Pers.) Steudel — Skhinias (10.11.1991: K. S. 4471, 29.9.1992: K. S. 4606); Iliki-Paralimni (29.8.1992; K. S. 4531).

Typha latifolia L. — Iliki-Paralimni (29.8.1992: K. S. 4532).

Cyperaceae

Scirpus cernuus Vahl — Skhinias (26.5.1988: K. S. 3506).

Scirpus holoschoenus L. — Skhinias (26.5.1988: K. S. 3520, 10.11.1991: K. S. 4472, 29.9.1992: K. S. 4613); Iliki-Paralimni (29.8.1992: K. S. 4536).

Scirpus litoralis Schrader — Skhinias (26.5.1988: K. S. 3508, 20.4.1991: K. S. 4236, 29.9.1992: K. S. 4607).

Scirpus maritimus L. subsp. *maritimus* — Skhinias (20.4.1991: K. S. 4234).

Fimbristylis bisumbellata (Forskål) Bubani — Sperkhios river (9.5.1983: K. S. 1863).

Cyperus diformis L. — Sperkhios river (9.5.1983: K. S. 1877).

Cyperus fuscus L. — Sperkhios river (9.5.1983: K. S. 1866); Iliki-Paralimni (29.8.1992: K. S. 4540).

Cyperus laevigatus L. subsp. *distachyos* (All.) Maire & Weiller — Kira (5.5.1991: K. S. 4347).

Schoenus nigricans L. — Skhinias (10.11.1991: K. S. 4477).

Carex extensa Good. — Kira (5.5.1991: K. S. 4348).

Conclusions

The present study on 9 aquatic ecosystems contributes to the knowledge of their flora and of the flora of Central Greece (Stereia Ellas) in general. The authors give a list of 164 taxa of vascular plants: 3 Pteridophytes and 161 Spermatophytes (101 Dicotyledones and 60 Monocotyledones). They are belonging to 113 genera and 45 families. The most diverse families are Gramineae (31 taxa) and Leguminosae (16 taxa) followed by Compositae (13 taxa), Chenopodiaceae (12 taxa) and Cyperaceae (10 taxa).

The number of families, genera and taxa of each examined ecosystem is reported in Table 1; the areas of Mesolongi saltworks and Skhinia marsh house the longest numbers of entities because of their size and the variety of their habitats. Moreover the authors have examined these two ecosystems more intensively than the others, some of which — especially Sperkhios river — require further investigation.

The areas under study belong to two ecosystem types: fresh water and estuarine ecosystems. From the clear aquatic plants the species *Ceratophyllum demersum*, *Myriophyllum spicatum*, *Potamogeton crispus* and *P. nodosus* inhabit the first ecosystem type, whilst the species *Ruppia cirrhosa*, *R. maritima*, *Zostera noltii*, *Cymodocea nodosa*, *Potamogeton pectinatus* and *Zanichellia palustris* live in the second type. The floristic composition of the areas around the water of the two ecosystem types also appear a remarkable difference.

Ecosystems	Families	Genera	Taxa
1. Mesolongi saltworks (canals, ditches, pans separated by banks)	21	54	67
2. Kira (rivulet)	3	4	4
3. Sperkhios river (bank)	15	20	22
4. Iliki-Paralimni (canal).....	11	19	20
5. Lake Paralimni	3	3	4
6. Oropos (pool)	12	22	23
7. Loutsa (streamlet)	1	1	1
8. Marathon (technical lake)	20	27	29
9. Skhinias (canal, marsh, pools)	19	37	53

Table 1. — Floristic composition of the 9 examined ecosystems.

ACKNOWLEDGEMENTS

We wish to express our thanks to Professor S. Snogerup (Lund) for his valuable help in the identification of some *Elymus*, *Juncus* and *Puccinellia* specimens.

REFERENCES

- BROFAS, G. & G. KARETSOS (1992). Simvoli sti dhierevnisi tis oikologias tou Skhinia-Marathona. *Geotechn. Sci. Issue* 3(1): 32-41.
- CASPER, S. J. & H.-D. KRAUSCH (1980-1981). Pteridophyta und Anthophyta, 1 & 2 Teil. In: ETTL, H., J. GERLOFF & H. HEYNIG (Eds.), *Susswasserflora von Mitteleuropa*, 23 & 24. Stuttgart & New York.
- COOK, C. D. K. & al. (1974). *Water plants of the world*. The Hague.

- DAVIS, P. H. (ed.) (1965-1988). *Flora of Turkey and the East Aegean Islands* 1-10. Edinburgh.
- ECONOMIDOU, E. (1975). La végétation halophytique de l'Attique et sa protection. *Coll. Phytosociol.* 4: 35-41.
- FASSETT, N. C. (1957). *A manual of aquatic plants*, ed. 2. Madison.
- GREUTER, W., H. M. BURDET & G. LONG (eds.) (1984, 1986, 1989). *Med-Checklist*. Vols. 1, 3, 4. Genève & Berlin.
- GREUTER, W., D. PHITOS & H. RUNEMARK (1976). Greece and the Greek Islands. A report on the available floristic information and on current floristic and phytotaxonomic research. *Coll. Int. C.N.R.S.* 235: 67-89.
- HARITONIDOU, P. (1977). I prostasia tou perivallontos tou megalou elous Skhinia Marathon. *I Physis* 10-11: 24-30.
- HARITONIDOU, P. (1983). Environmental protection for the Great Marsh at Skinias. Can its vanishing flora and fauna be saved? *Ann. Musei Goulandris* 6: 45-52.
- KOUMPLI-SOVANTZI, L. (1983). *Studies on the Tracheophytes in the lakes and adjacent hydrobiotopes of Aetoloakarnania, Greece. Taxonomic, floristic, phytogeographical and ecological research*. Diss. Athens.
- KOUMPLI-SOVANTZI, L. (1989). The aquatic flora of Aetoloakarnania (W. Greece). *Willdenowia* 18: 377-385.
- KOUMPLI-SOVANTZI, L. & F. VALLIANATOU (1985). The aquatic vegetation of the lake Lyssimachia (Western Greece). *Thalassographica* 8: 33-41.
- PHITOS, D. (1988). To dhasos Skhinia — Ena khameno orama. *I Physis* 43: 7-12.
- SEVERIN, I., Th. GEORGIADIS & D. CHRISTODOULAKIS (1989-1990). Simvoli sti meleti kai axiologisi ton oikosistimatou tou delta tou Akheloou. *Parartima Protou Tomou Epetir. Tmimatos Violigias A. P. Th.*: 221-230.
- STEPHANIDES, Th. (1948). Synoptic notes on the fresh-water organisms of certain regions of Macedonia. Epirus and Central Greece. *Prakt. Hellen. Hydrobiol. Inst. Acad. Athens* 2: 205-213.
- TUTIN, T. G. & al. (eds.) (1964-1980). *Flora Europaea* 1-5. Cambridge.
- WOLFF, W. J. (1968). The halophilous vegetation of the lagoons of Mesolonghi, Greece. *Vegetatio* 16: 95-134.

