

**Zeitschrift:** Candollea : journal international de botanique systématique = international journal of systematic botany  
**Herausgeber:** Conservatoire et Jardin botaniques de la Ville de Genève  
**Band:** 51 (1996)  
**Heft:** 1

**Artikel:** Two new species of Fuirena (Cyperaceae) from Somalia  
**Autor:** Arnstein Lye, Kaare  
**DOI:** <https://doi.org/10.5169/seals-879397>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 01.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Two new species of *Fuirena* (Cyperaceae) from Somalia

KAARE ARNSTEIN LYE

## RÉSUMÉ

LYE, K. A. (1996). Deux espèces nouvelles de *Fuirena* (Cyperaceae) de Somalie. *Candollea* 51: 1-5. En anglais, résumés anglais et français.

Deux espèces nouvelles, *Fuirena boreocoerulescens* Lye, proche du complexe *F. coerulescens* d'Afrique du Sud, et *F. striatella* Lye, proche de *F. wallichiana* Kunth d'Inde, sont décrites. Ces deux espèces ont des aires de distribution restreintes dans les montagnes du nord de la Somalie, entre 1350 et 1500 m d'altitude.

## ABSTRACT

LYE, K. A. (1996). Two new species of *Fuirena* (Cyperaceae) from Somalia. *Candollea* 51: 1-5. In English, French and English abstracts.

*Fuirena boreocoerulescens* Lye, related to the South African *F. coerulescens*-complex, and *F. striatella* Lye, related to *F. wallichiana* Kunth from India, are described as new. Both species have very restricted distributions in the mountains of Northern Somalia at altitudes ranging from 1350 to 1500 m.

**KEY-WORDS:** CYPERACEAE — *Fuirena* — Somalia — Africa — Taxonomy.

Due to the publication of Cyperaceae for "Flora of Somalia" (ed. M. Thulin), it is necessary to publish two new species of *Fuirena*.

***Fuirena boreocoerulescens* Lye, spec. nova, Fig. 1A-F & 2.**

**Type: Somalia. Sanaag region:** Markat, 48°30'E & 10°59'N, in clay flush, 1460 m, 12 Aug 1957, J. G. Newbould 883 (K, holo).

*Herbae perennes. Rhizoma horizontaliter repens lignosum crassum in intervallis brevibus culmos florentes emittens. Culmus 20-40 cm altus rigidulus trigonus dense pubescens. Spiculae 6-10 mm longae et 4-5 mm latae oblongo-ellipticae multiflorae. Squamae 2.0-2.5 mm longae pubescentes dense imbricatae; apice mucronatae. Setae hypogynae 3. Squamellae hypogynae 3 membranaceae. Nux 1.0-1.2 mm longa.*

Slender perennial with a horizontally creeping rhizome 2-3 mm thick with culms at about 5 mm intervals, rarely crowded. Culms 20-40 cm long and 1-2 mm thick (but sometimes 0.6-0.9 mm thick below the inflorescence); triangular, minutely pubescent with mostly adpressed hairs (most densely hairy below the inflorescence); the hairs about 0.1 mm long only. Leaves 4-6 and spaced all along the culm; the lowermost with a short triangular limb only, other with well developed blades; the largest blade 10-20 cm long and 3-5 mm wide, minutely short hairy on margin at least above, becoming glabrescent with age; sheaths glabrous, the lowermost brownish, the other green; ligule 1-2 mm long, cylindrical, glabrous, pale but with orange or reddish brown minute gland-dots. Inflorescence a terminal cluster of 3-10 crowded spikes subtended by 2-3 leaf-like involucre bracts




Fig. 1. — A-F: *Fuirena boreocoerulescens* Lye, from *Newbould 883*. A: habit  $1/2\times$ ; B: part of culm with base of leaf-blade with ligule  $5\times$ ; C: spikelet  $5\times$ ; D: glumes  $15\times$ ; E: perianth  $15\times$ ; F: nutlet  $15\times$ . G-K: *F. striatella* Lye, from *Newbould 861*. G: habit  $1/2\times$ ; H: part of culm with base of leaf-blade with ligule  $5\times$ ; I: spikelet  $5\times$ ; J: glume  $15\times$ ; K: nutlet with perianth  $15\times$ . — Drawn by Gerd Mari Lye.





Fig. 2. — *Fuirena boreocoerulescens* Lye. Achene, scale 0.1 mm. SEM photograph from the holotype *Newbould 883*.

Fig. 3. — *Fuirena striatella* Lye. Achene, scale 0.1 mm. SEM photograph from the holotype *Newbould 861*.

0.5-3 cm long. Spikes 6-10 mm long and 4-5 mm wide, oval-elliptic with obtuse apex and with 40-60 densely imbricate glumes. Glumes 2.0-2.5 mm long (excluding the 0.7-1 mm long mucro) and 1.2-1.5 mm wide, rectangular, light reddish brown below, greyish above, but often with reddish brown lines, minutely short-hairy both on surface and margin; the midrib 3-nerved, slender, but thickened at apex and excurrent into a thick brown straight hairy mucro. Perianth segments 6; the three outer light reddish brown, hairy, filiform or somewhat flattened in the lower middle part, 1-2 mm long and of very unequal length; the three inner (or at least one) with a well developed obovate stalked light reddish brown blade irregularly thickened and crisped above; the midrib prominent, usually excurrent into an about 0.5 mm long filiform hairy awn; the margin short-hairy or more commonly with hairy lobes or thickenings. Stamens 3; filaments about 2 mm long, flattened, light reddish brown; anthers about 1.5 mm long and 0.3 mm wide, linear, light reddish brown with a short darker connective. Style 1.5-2 mm long with 3 long very hairy branches. Achene 1-1.2 mm long and about 0.7 mm wide, light reddish brown, triangular-oval with cuneate base and tip; the apex glabrous or slightly hairy; the surface cells prominent, isodiametric to rectangular with darker cell walls.

In open grassy area on clay and in clay flush, 1350-1500 m. Endemic to the Sanaag region in northern Somalia, but closely related species occur in South Africa (*Fuirena coerulescens* — complex).

*Other specimens examined.* — **Somalia. Sanaag region:** Escarpment SE of Laasqoray, near Ragad, 48°29'N & 11°00'N, open grassy area on clay on limestone with evergreen bushland with *Buxus* and *Cadia*, 1350 m, 16 Jan 1995, *M. Thulin, A. Dahir & A. Hassan 9085* (K, UPS).

***Fuirena striatella* Lye, spec. nova, Fig. 1 G-K & 3.**

**Type: Somalia. Sanaag region:** Ragad, 48°30'E & 10°57'N, clay flush, 1370 m, 11 Aug 1957, *J. G. Newbould 861* (K, holo).

*Herbae annuae, pubescentes. Culmus 2-10 cm altus trigonus vel teres glabrescens. Spiculae 4-5 mm longae et 2-3 mm latae ovatae multiflorae. Squamae 1.5 mm longae pubescentes dense imbricatae; apice mucronatae (mucro ad 1.0 mm longa). Setae hypogynae 3. Squamellae hypogynae 3 minutae. Nux 0.8-1.0 mm longa.*

A small tussocky annual with few-many crowded culms. Culms 2-10 cm long and about 1 mm thick, but perhaps much longer when not heavily grazed, often almost terete with longitudinal ridges, glabrous or with a few stiff hairs. Leaves 3-5; the lowermost without or with very short blade; sheaths light reddish brown, brownish or green, densely short-hairy with spreading hairs, but uppermost sheaths usually glabrescent; ligule about 1 mm tall, densely short-hairy to glabrous; blades probably to 20 cm or more long and 2-5 mm wide, rather thick, hairy on margin and sometimes on upper surface also. Inflorescence a terminal cluster of 3-10 crowded sessile or subsessile spikelets, sometimes with an additional stalked spikelet-cluster from the uppermost sheath of the stem. Spikes 4-5 mm long and 2-3 mm wide, oval with obtuse apex, with 20-40 densely imbricate glumes. Glumes about 1.5 mm long (excluding the up to 1 mm long mucro), oval-elliptic, reddish brown to greyish, hairy particularly near margin and on the excurrent greenish midrib. Perianth-segments usually six, whitish; the three outer filiform, often curved and only 0.2-0.3 mm long; the three inner about 0.8 mm long, filiform below, but ending in a minute irregular or anchor-like lamina only about 0.2 mm wide. Achene 0.8-1 mm long and about 0.5 mm wide, obovate, obtusely triangular, light reddish brown with longitudinally raised ribs and narrow rectangular surface cells between the ribs.

In clay flush with heavily grazed vegetation, 1350-1400 m. Endemic; only known from the type from the Sanaag region in Northern Somalia. Its nearest relative, *F. wallichiana* Kunth, occurs in India.

ACKNOWLEDGEMENTS

The electroscan-photography was made possible through E. Reed at Laboratory of Analytical Chemistry, Department of Electron Microscopy, The Agricultural University of Norway. The photographs were taken by the author. I would also like to thank Gerd Mari Lye for the drawings and Mette Solsvik for typing and editing the manuscript.

