**Zeitschrift:** Candollea: journal international de botanique systématique =

international journal of systematic botany

**Herausgeber:** Conservatoire et Jardin botaniques de la Ville de Genève

**Band:** 51 (1996)

Heft: 2

**Artikel:** Ephemeral flush vegetation on inselbergs in the Ivory Coast (West

Africa)

Autor: Dörrstock, Stephan / Porembski, Stefan / Barthlott, Wilhelm

**DOI:** https://doi.org/10.5169/seals-879417

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

# Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 18.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Ephemeral flush vegetation on inselbergs in the Ivory Coast (West Africa)

STEPHAN DÖRRSTOCK STEFAN POREMBSKI & WILHELM BARTHLOTT

#### **ABSTRACT**

DÖRRSTOCK, S., S. POREMBSKI & W. BARTHLOTT (1996). Ephemeral flush vegetation on inselbergs in the Ivory Coast (West Africa). *Candollea* 51: 407-419. In English, French and English abstracts.

A detailed description of floristic aspects and of the ecology, biogeography and species diversity of the ephemeral flush vegetation on West African granite inselbergs is given. Due to stressful environmental conditions inselbergs bear a highly distinct vegetation. The ephemeral flush vegetation is a seasonal, low growing herb community which occurs over shallow soils on inclined rocky slopes. Floristically the high number of carnivorous species (e.g. *Utricularia* spp., *Genlisea stapfii*) is remarkable. Species diversity of this habitat increases from the rainforest towards the savanna region of the Ivory Coast. With up to 35 species per relevé the ephemeral flush vegetation is the most speciose community on West African inselbergs.

#### RÉSUMÉ

DÖRRSTOCK, S., S. POREMBSKI & W. BARTHLOTT (1996). Végétation des micro-marécages temporaires des inselbergs de Côte-d'Ivoire (Afrique de l'Ouest). *Candollea* 51: 407-419. En anglais, résumés français et anglais.

Une description floristique, écologique et biogéographique avec considération de la diversité de la végétation des micro-marécages sur les inselbergs granitiques en Afrique de l'Ouest est présentée. La végétation des micro-marécages est un groupement de petites herbes sur sol très mince qui se forme pendant la saison des pluies au pied des pentes rocheuses. L'ensemble floristique de ce groupement se distingue par un grand nombre d'espèces carnivores (e.g. *Utricularia* spp., *Genlisea stapfii*). La richesse floristique de cette association augmente, en Côte-d'Ivoire, de la zone forestière vers la région des savanes. Avec un maximum de 35 espèces par relevé, c'est l'habitat le plus riche en nombre d'espèces sur les inselbergs de l'Afrique de l'Ouest.

KEY-WORDS: Carnivorous species – Climatic and latitudinal gradient – Ephemeral flush vegetation – Inselbergs – Ivory Coast – Seasonality – Species richness.

### Introduction

The term "inselberg", created by BORNHARDT (1900), characterizes granitic and gneissic, mostly dome-shaped rock outcrops which are geomorphologically old (frequently more than 20 million years). A detailed survey of the geomorphology of inselbergs is given by BREMER

CODEN: CNDLAR 51(2) 407 (1996)

CONSERVATOIRE ET JARDIN
BOTANIQUES DE GENÈVE 1996

& JENNINGS (1978). Inselbergs are widespread in many parts of the tropics and subtropics. Due to the lack of substrate and extremes of microclimate they bear a vegetation that is distinctly different from the surroundings.

Most inselbergs harbour a highly seasonal plant community of tiny hydrophytes which occurs on slightly inclined slopes in small seepage zones below patches of vegetation (e.g. *Afrotrilepis pilosa*-mats) where water seeps continuously during the rainy season. Following RICHARDS (1957) the term "ephemeral flush vegetation" will be used for this community. Only marginally mentioned in literature (e.g. ADJANOHOUN, 1964; BONARDI, 1966; GUILLAUMET, 1967; HAMBLER, 1964; MILDBRAED, 1922; POREMBSKI & BARTHLOTT, 1993; RICHARDS, 1957; VILLIERS, 1981) this vegetation type has not been examined in detail so far. The intention of this study is to present a first description of floristical, ecological and biogeographical details of this nearly unknown habitat.

### Study area

The Ivory Coast is situated between 4°25′-10°40′N and 2°30′-8°40′W. Floristically Guineo-Congolian elements are dominating, but in the north of the country Sudanian affinities exist. The climate of the Ivory Coast is influenced by the shift of the ITCZ (intertropical convergence zone). The south of the country is mainly covered by lowland rain forest, secondary grassland and cultivational areas. Precipitation lies between 1600 to 2300 mm per year with a weakly expressed dry season in July/August and a more severe dry season during December/January. The northern part of the country is covered by several types of woodland respectively savanna. Annual rainfall is between 900-1500 mm/year with a dry season from November to March/April. Detailed descriptions of climate, flora and vegetation of the Ivory Coast are given by AKÉ ASSI (1984), ANHUF (1994), ELDIN (1971) and GUILLAUMET & ADJANOHOUN (1971).

Inselbergs in the Ivory Coast consist of granite and gneiss (LENEUF, 1959). They occur all over the country, with a higher number in the savanna region than in the rain forest zone. Inselberg size ranges from several 100 m² and a few metres in height ("shield inselbergs") up to an area of 7 km² and a height of 300 m above the surroundings. The microclimate of inselbergs is characterized by extremes of temperature and insolation. Even on inselbergs situated in rain forest, temperatures on the rock regularly exceed 60°C and relative air humidity falls below 30%. High wind speeds can be recorded on inselbergs which are caused by thermic convection.

# **Inselberg vegetation**

The vegetation of inselbergs can be subdivided into characteristic habitats, e.g. cryptogamic crusts on rock surfaces, rock crevices, shallow depressions, seasonally water-filled rock pools, monocotyledonous mats, ephemeral flush and wet flush vegetation. Cryptogamic crusts on Ivorian inselbergs mainly consist of lichens (in particular *Peltula* spp.). Narrow rock fissures are colonized by short-lived herbs (many Poaceae and Cyperaceae), whilst crevices may be occupied by shrubs (e.g. Hymenodictyon floribundum) and trees (e.g. Ficus spp.). Afrotrilepis pilosa, a poikilohydric stem producing Cyperaceae dominates mat communities of West African inselbergs. Afrotrilepis-mats offer growing sites for a number of geophytes, such as Albuca nigritana, Ceropegia deightonii and Eriospermum abyssinicum. A wet flush community develops on inclined rocky slopes with water steadily running down during the rainy season. Typical are crusts of cyanobacteria which offer establishment sites for tiny vascular plants, like *Utricularia subulata*. For a more detailed description of habitats on inselbergs it is referred to BARTHLOTT & al. (1993). In the following, ephemeral flush vegetation is abbreviated to EFV. General descriptions of the Ivorian inselberg vegetation have been provided by ADJANOHOUN (1964), BONARDI (1966), POREMBSKI & BARTHLOTT (1992, 1993), POREMBSKI & al. (1995) and POREMBSKI & BROWN (1995).

#### Material and methods

Altogether 38 ephemeral flush communities, from inselbergs all over the Ivory Coast (Fig. 1) were investigated during September/October 1993, 7 sites were studied in 1991/1992. The description of the vegetation was carried out according to the Braun-Blanquet relevé method (BRAUN-BLANQUET, 1964) with a modification in cover abundance scale following BARK-MAN & al. (1964) and WILMANNS (1984). The plots were randomly located with a size of 4 m<sup>2</sup>. Relevé size was based on a minimal area analysis.

Life-form spectra were compiled in accordance with the RAUNKIAER (1934) classification, the classification of pollination modes according to syndromes follows FAEGRI & VAN DER PIJL (1979). Dispersal modes were classified according to syndromes after VAN DER PIJL (1982) with certain modifications following BERG (1983) and JENNY (1994).

For water pH-analysis, Merck indicator test-sticks were used. Soil analyses were carried out at the Pedological Institute of the University of Bonn: pH was determined by a glass electrode in a 0.01 M CaCl<sub>2</sub>-solution, the nutrient content was determined by CAL-extract, quantitative analysis of the organic matter was determined by measuring the ash content. The soil type was determined by hand, the soil fraction >2 mm diametre was determined by electronic scales. Water analysis by the Wahnbachtalsperrenverband (using the Inductively Coupled Plasma (ICP)-method).

For germination experiments soil samples were dried for 3 months and subsequently heated (75°C) in a heat box for 3 hours. Afterwards the soil was put out in pots (which were kept in a greenhouse at 25-30°C air temperature) over a layer of granite fine cherts. For control unheated soil samples were hold under the same conditions.

### Results

### Abiotic factors

The ephemeral flush communities studied occurred at the feet of rocky slopes below *Afrotrilepis*-mats and other water-holding vegetation types (Fig. 2) or on sandy soils in transition zones between shield inselbergs and the surrounding vegetation. The degree of slope inclination was mostly between 1-10°, sometimes up to 25°. The sites studied occurred at altitudes between 100 and 620 m asl. and up to 300 m above the surrounding plains. Size of the communities studied ranged between 4 (narrow fringes bordering *Afrotrilepis*-mats) and 200 m<sup>2</sup>.

Soil depth ranged from shallow to very shallow (0.5-12 cm) with lowest values at the community margin towards the rock. Characteristic was a mosaic of different soil depths occurring close together. The soil had a sandy or sandy-loamy texture and a high humus content (5-12% organic substance). The soil fraction >2 mm diametre reached 11-28.5%. The soil can be classified as lithosol or ranker (FAO-UNESCO, 1988). Soil pH varied between 4 to 5.5.

Apart from direct precipitation the sites studied were supplied with seepage water (nutrient content was low, s. Table 1). Depending on the water-holding capacity of the vegetation situated above the ephemeral flush communities seepage water was available for 1-14 days after rainfall. If the supply of seepage water comes to an end, the soil dries out within 2 weeks.

# Physiognomy of the vegetation

The EFV is a low growing (5-70 cm) herb community. Frequently the vegetation cover is interrupted by bare rock. Usually 3-4 intergrading zones can be distinguished according to soil depth (Figs. 3, 4, 5): a transitional zone against the rock, a marginal zone, a central zone and an





Fig. 1. – Map of the Ivory Coast showing natural vegetation zones (modified after MONNIER, 1993). Inselbergs studied indicated by white dots. The absence of ephemeral flush communities is marked by a bar.



 $\label{eq:Fig. 2.} Fig. \ 2. - Ephemeral \ flush \ vegetation \ (EFV) \ at \ the \ feet \ of \ a \ rocky \ slope \ on \ Mt. \ Niangbo. \ White \ bars \ indicate \ the \ border \ of \ this \ community. \ BR = bare \ rock, \ GM = grass \ mats.$ 

Fig. 3. – Central zone of an ephemeral flush community on Mt. Niangbo with *Nemum spadiceum*, *Panicum lindleyanum* and *Utricularia juncea*.

Fig. 4. – Marginal zone of an ephemeral flush community with *Aeollanthus pubescens*, *Burmannia madagascariensis*, *Drosera indica*, *Panicum lindleyanum* and *Utricularia micropetala*.

	Tal	ole 1. – Ar	alyses of	water sam	pled in eph	nemeral flu	ish commu	nities.		
E1							9			13
From relevé nº		1	2	3	4	8	9	10	12	13
Conductance	mS/m	1	3	11.6	1.8	2.1	3.4	2.7	1.8	0.6
NO2	mg/l	<0.01	<0.01	< 0.02	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
NO3	mg/l	1	1	1	1	2	2	1	1	1
NO4	mg/l	0.109	1.067	10.147	1.174	0.042	0.025	0.684	0.458	0.155
P	mg/l	0.02	0.03	0.03	0.02	0.02	<0.02	0.02	<0.02	0.03
0-PO4-P	mg/l	< 0.005	< 0.005	< 0.005	0.011	< 0.005	< 0.005	< 0.005	< 0.005	<0.005
K	mg/l	0.2	0.5	1.9	0.5	0.4	0.6	0.6	0.4	0.1
Mg	mg/l	<0.1	0.1	0.41	<0.1	0.1	0.4	0.1	0.1	<0.1
Ca	mg/l	0.4	1.6	1.7	0.2	0.9	1.5	1.5	0.9	0.2
Na	mg/l	1	2.3	4.8	1.3	3.1	5.5	2.4	1.8	0.5
Mn	mg/l	< 0.005	< 0.005	< 0.005	0.005	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005
Al	mg/l	0.015	0.013	<0.005	0.024	0.018	1.036	0.005	0.04	0.031
Fe	mg/l	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005	0.214	< 0.005	< 0.005	<0.005
Si	mg/l	1.6	3.1	5.4	1.6	1.8	7.7	2.2	1	0.3
SO4	mg/l	0.1	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.1

upper transitional zone (sometimes lacking) towards habitats, like *Afrotrilepis*-mats. The transitional zone against the lichen covered rock is dominated by cyanobacteria. The marginal zone (soil depth 0.5-3 cm) is characterized by the scattered occurrence of tiny annuals (5-20 cm in height) with a high percentage of carnivorous species. Within the central zone (soil depth 3-10 cm) Cyperaceae and Poaceae (20-50 cm in height) dominate. Tall grasses are typical for the upper transitional zone. This zonation may become obscured due to the patchy distribution of different soil depths. Most species recorded were not restricted to a single zone, however, they occurred preferentially in a particular one.

# Floristic composition and phytosociological aspects

One hundred fifteen species out of 29 families were found (s. Table 2). Most species belong to the Cyperaceae (26 species), Poaceae (21 spp.), Lentibulariaceae (10 spp.), Scrophulariaceae (10 spp.) and Eriocaulaceae (6 spp.). Most speciose is the genus *Utricularia* (9 spp.), followed by *Cyperus* (7 spp.) and *Eriocaulon* (6 spp.). Many genera occurred with only one species (e.g. *Xyris*).

Carnivorous species like *Drosera indica*, *Genlisea stapfii* and *Utricularia* spp. were the most conspicuous elements according to their high share in species number compared with other habitats on inselbergs. Carnivorous plants characterize the EFV and discriminate this community from other inselberg habitats. Moreover the high species number of small-sized Poaceae and Cyperaceae is typical for the EFV. *Xyris straminea* occurred with a high degree of constance (95.4%) and forms also an essential component of the EFV. Further species with a high degree of constance were *Drosera indica* (93%) and *Panicum lindleyanum* (90.7%). In particular the latter species is highly characteristic for the EFV since it usually occurs with high cover percentages and has only rarely been found in other communities. Other frequently recorded (> 50%) species were *Utricularia pubescens* (76.6%), *Utricularia subulata* (72.1%), *Aeollanthus pubescens* (69.8%), *Afrotrilepis pilosa* (67.4%), *Bulbostylis congolensis* (67.4%), *Ascolepis protea* (62.8%), *Scleria melanotricha* (62.8%), *Sporobolus pectinellus* (60.5%), *Nemum spadiceum* (55.8%) and *Cyanotis lanata* (51.2%). In comparison, 61% of the species were found in only 10% of the relevés. Therefore, the homotoneity of the relevés is low.

### Affinities to similar habitats

No species found in the EFV is restricted to inselbergs. Forty nine out of the 115 species recorded mainly occur on inselbergs where they are restricted to the EFV (e.g. *Genlisea stapfii*). Floristic links exist with inselberg communities in shallow soil-filled depressions (32 species) with Poaceae dominating and seasonally water-filled rock pools (26 species). Eight species also occur in *Afrotrilepis*-mats, 6 species are colonizers of rock crevices and 5 species grow among rock debris.

Outside inselbergs many species observed within ephemeral flush communities occur on open places over wet sandy soil or on swampy respectively marshy sites as well as on ferricretes (locally known as "bowal"). ADJANOHOUN (1964) described a *Sporobolus pectinellus-Cyanotis lanata* community occurring on ferricretes that has strong physiognomic and floristic affinities with the EFV.

# Phytogeography

More than 50% of the species found are distributed all over tropical Africa or are paleotropical respectively pantropical elements. Thirteen species are paleotropics, 9 species are pantropics, 6 species show a disjunct distribution between tropical Africal and the neotropics. *Utricularia juncea*, which hitherto was only known from the New World, has been found at two sites in the Ivory Coast during this study (DÖRRSTOCK & al., 1995). Fifteen species (e.g. *Eriocaulon plumale* ssp. *jaegeri*, *Genlisea stapfii*, *Lipocarpha filiformis*, *Loudetiopsis capillipes*) are restricted to the Upper Guinea province, 17 species occur in the Upper Guinea-Nigeria/Cameroon-provinces and further 17 species are widespread throughout the Sudano-Zambezian Region. No species is endemic to the Ivory Coast.

#### Phenology

The vegetation period starts 4-8 weeks after the begin of the rainy season with the germination of the first seeds. Experiments showed that for *Drosera indica* the germination rate is increased after heat treatment (3 hours at 75°C) of the seeds. The vegetation cover reaches its maximum at the end of the rainy season. At most sites a cover degree of 70-90% was attained,




Fig. 5. – Schematic cross section of typical ephemeral flush vegetation.

however, at some sites the cover degree reached only 50%. On average the cover degree was higher in the rain forest region (70-100%) than in the savanna zone (50-90%).

The first flowers appeared 3-4 weeks after germination was observed (several Cyperaceae). The main flowering period lies at the end of the rainy season. At this time many fruit bearing individuals can be found too. Certain geophytes (e.g. *Eriospermum abyssinicum*) preferentially flower during the dry season. In the rain forest zone some species (e.g. *Utricularia* spp.) are able to complete their life cycle twice a year due to the prolonged vegetation period.

# Life-forms and vegetative strategies

Most species found were therophytes (79%), followed by hemicryptophytes (9.6%), geophytes (6%), meso-phanerophytes (4.3%) and one chamaephyte (*Afrotrilepis pilosa*). The number of perennials increases in the rain forest region. The high percentage of carnivorous species is indicative of the nutrient deficiency of the sites studied. Remarkable is the occurrence of carnivorous "climbers", such as *Utricularia micropetala*, *U. spiralis* and *U. tortilis*. Using grasses as support these species may expose their flowers more conspicuously.

### Pollination biology

Entomophilous species (57%) dominate over anemophilous taxa. Concerning the number of entomophilous flowers the ephemeral flush community is much richer than other inselberg habitats (own data). However, it is conspicuous that only a very small number of pollinating insects (e.g. *Apis mellifera adansonii*) could be observed. Autogamy seems to be an important factor. Cleistogamous forms of *Utricularia juncea*, *U. pubescens* and *U. subulata* (s. KILLIAN, 1953;

ude (N)	7°45′	8°35	8°4	2′8°	49 ′	8°49′	8°49	8°49	' 9°27	9°36	9°32'	9°15	7°45	6°45′	8°35′ 8	3°35′8	°35′8°3	5′8°3	5′8°35	8°35′	8°35′	8°35′8	35′8	35′ 8°35	8°35	8°25'	8°49′ 8	8°49′8°	49 ' 80	49′8°4	9′8°49	° 8°49	8°49′	8°49′	8°49′	8°36′ 9	°40′	8°29′7	°15′7'	25′ 7°	25′ 5	5°25′
pitude (W)	5°10′	4°40	3°3	5 5°	111	5°11′	5°11	5°11	5°38	6°55	6°29'	6°40	6°40′	7°21′	4°40′4	1°40′4	°40′ 4°4	0 ' '4°4	10 ′ 4°40	4°40′	4°40′	4°40′4	40 ' 49	40 · 4°40	4°40	4°32'	5°11′ E	5°11′5°	11′5°	11′5°1	1′ 5°11	6°11′	5°11′	5°11′	5°11′	3°46′ 6	°30′	6°35′8	°1′ 7	35′ 7°	35 7	7°10′
at above sea level [m]	370	410	31	5	40	530	530	530	470	460	430	410	300	200	340	340 3	180 40	420	420	440	450	500 4	60 4	55 430	400	360	450	450 5	50 62	20 60	570	490	490	500	500	280	525	220 3	70 2	40 2	70	260
ation [°]	10	11	3		6	8	5	8	17	2	12	2	2	8	3	2	3 21	12	2	7	6	3	4	8 3	7	14	6	17	3 3	3 6	2	2	3	9	5	3	3	7	5	3	5	7
ition	SE	SE	sv	/ 5	SE	s	s	s	s	s	SSE	Е	Ε	Е	w	w	N N	N	N	N	N	w	1E	E S	E	s	N	N V	N N	IE N	E	s	sw	s	s	N	Е	s	w s	w s	SW	SE
degree (%)	80	70	75		70	75	65	75	50	60	85	60	90	90	75	50	50 70	85	70	60	50	80	0 8	30 70	50	80			0 5	0 60	60	60	70	70	80	40	70					80
s number		31	34		30	10	10	29	14	32				11																												
												22	19				35 20				32			21 15	18	8			3 2			10	27	8	16	24	17					5
rea [m²]	20	120	40	•	50	25	20	25	15	120	65	30	50	50	8	110	25 50	200	70	60	120	100 1	00 4	40 25	100	20	80	25 2	0 4	0 10	120	60	150	40	30	40	25	40	10	15 1	10	4
straminea era indica	1	1 p	1	2	2m	1		2m	1	2m	2m		2m 2m	1	2m		2m 2n	1 2m	2m	2m	2m		m 1	1 2m	2m 2m	1		2m 2m 2	1 2: m 2:	m 2n		:	2m	*	1	1	1	р	1	1	1	1
ım lindleyanum		2b	3	2	2b	4	4	3	- 1	2b	2b		Zm		3	2a :	2b 3	2m	4	3	3			3 1	2m 2b	4	3	2m 2			3	3	3	3	3	3	1	р 4	3 ;	1 2b 2	1 2b	5
laria pubescens laria subulata	1	1 2m	1		1	1	P	1	p 1	1 2m	1 2m		1.	2a 1	1 2m	1 2	2m 1 2m 1	1	1	1	1	1 2	m 2	m 2m	2m	2m	1	1 2	m 1	1 .	1		1				1		1			2a
inthus pubescens		1			2a	1	p	1							1	1	1 1	2a	i i	1	1	i	1	1 2a	1		1	1	1 1	1 1	1	1	1	1	2a	2a	i					. 28
ilepis pilosa stylis congolensis	1 2a	1 2a			2a 2a	4	D	1	, n	p	p 1		2a 4	2a	1		2a 1 p 1	1	1	1 2a	1 .	1 2a 2	1 2 2b	2b 3 1 2a	1	2b	2a		1 2	a 1	1		1		;		3			2a		
melanotricha	-	1			1	1	3	1		í	2a	- 1			1	1	. 1	1	1	1	i	1	1 2	2a 1	- 1	i	1	1	i .				1				2a	1 :	2a :	2a 2	2a	
opis protea polus pectinellus	1	P 1			1		1	p 1	1	1	1	1 2a	2a	1	2m	P 1	1 p	2m	1 1	1	P	1 n		p 1	1		1	1 2	la 1	1 1		2a	p 1	2a		1	2b					
n spadiceum					1	1		2a			. 1		-	4			. 1	1	2a	2a	1	1	1				3	3	. 1		2a	2b	2a		2b	:		2a :	2b :	2a 2	2a	
tis lanata Irpha filiformis	p 1	r P	p 1		1			1		1 p	P 1	2a	1		2b 2m	1	1 .		р 1	i	i	1		1 . D .			1	1	. 1	1 1	1	1	p 1			2a						
ulon plumale ssp.jaegeri		2a	2m	1 2	2m			2m		1							1 1	2m	1 1	2a	2a				2a	- 1			a .				2a			1		p	1	1		
ulon afzelianum theca loeselioides		1	1		1		P 1	p	1	1 p	1	1		i	2m	1	1 .		1	2a	1	1	. 2	2a .	1		1	1 2	ta 1		1	D	1					1		1		2a
nia exilis	1	p	1		1		-	1	1		1		1	1		p	1 .		1	-	-			. :	1				1 1	i i	1	1			;	:	1	i				
glossum costatum tiopsis capillipes		P	p	2	2a	2a		2a	1		p 2a	1			1	p		2a	1 D	p	1 2b	1		р .	1	. :	2a	2a	1 .	 b 2a	2a		2b	2b	:	2a	1		1	1 2a 2	2b	-
nia schweinfurthii	:	1			1			1		1	- :	:			2m	1 2	2m .		1	1		1	1	1 1	1			:	1 1	1 .	1				-	i	1					÷
m griffonii a floribunda	p	1	1								1	1	1		1		 2m p	2a	1	- 1		2a		2a . . 1	- 1								P	3	3	:	1	. :			2b 2a	-
pogon africanus la scabra	5	3								- 1		2a		į.		2a	. 20	1	2a	9							2a		1 2	b .	2a	2b	2b	2a	1							
nnia madagascariensis		. 1	ρ.		1	-:	- :-	i				ρ.		:		P .	p p	2m	2m	2m	2m				i		1	1		. 2m			i				P .					
aria micropetala ia parviflora		D			2a D	2m		1		n .	2m	÷	*		D	;			;		*	2a					2m	2a	١.	. 1	1		1		1							
aria juncea		2a		2	2a	1		1			- 1					:		2a		2a	2a	1						2a			1		1	-								
iopsis kerstingii oogon curvifolius		1						:						2b	2a	p :	2b .		P	р 1	1	2a 1 2	la 2	. 2b 2a 1	2b	2a										2a						
a stapfii		2m							- 1	1	1			-			. 2m	2m	. :	2m	2m	1 '			i													1				
ka heudelotii stylis alboviridis	p	- 0	p 1	1	P			p		D	P		1								1	p 1											р	р								
a cf.filifolia la arenaria			р					-		P							. P		p	Р	P	р		. р	÷											-						
ulon togoense		1											1		:	:	P :	Р.	P 1		1 1		:	1 .	- :			ρ.			1		Р.			:	:	1				
s capillifolius tis longifolia		p								1											р	1	1	1 .													÷					
tia simplex			p									2a	2a	2			: :														3		2a				2a					
lossum gomezianum tenella	1		1					1	- 1	1	i				:	:	1 1																			1	1					
asp.			p								1 -					p	р .	p							- 1											p						
tium trianthum mimosoides	p					:		Р.								:	p .		÷															D	r	ŗ	:					
nigritiana stylis dichotoma			1							1		1	1																							1						÷
olus infirmus								1				ρ.					: :		. :			1			- 1										i	1 2b		1	:			
thloa indica us squarrosus		1	8							1						D			1	- 1													1			1						
aria spiralis			1											:							1	p		. :			:								1			1				:
a sp. rmum abyssinicum				-	P .					:		р .		1		P	: ::				÷				Via month	de esta de la composição		10000								:	:					
era sp. pis capensis													р			:							-												p	r			;			
pis pusilla			1										:	:			i .		- 3																							
ia leptorhachis izachyrium sp.	1		1									1		:	:	:	: :			. :		:						: 1			8					1						
aris atropurpurea							- 1	- 1		-	- :	1					. :	- :	- 1								1			. :			- 1			:	:					÷
a sp. iella ypsilostyla				- 1	р		1		P							:	: :																		:	:	i	:				
ulon remotum ila lecardii										1	Р			;																								÷				
a pumila			Р.		:	:	1	p p						- !		:	: :		- 1					: :				1						1			:					:
s basserei elina sp.						:				D	D					:											p	p .				-		4								
lobata			r											:				- :		- 1		÷					:	: :							ř		:					
ia togoensis olus festivus	:						- 1		3					:	:	:						;					:	: :							:	:		2b				
a diminuta	- 1										1				V		. :		- 1																	1	:					
orhopalon debilis s cuspidatus					:					- :			1	:	:	1					1							: :								:						
s dubius s haspan			:																										. 1													÷
s pustulatus			1							- 1				:		:	: ::							: :			:	: :					:	÷			:					:
ceae sp. s tenuispica														1	1	:	: :										1	: :														
opogon hagerupii									1	- 1		-	- :			:				- :		÷						: :								:	:					1
itis gangetica itis scotelliana			1			5						1				:		-				:					:															
lon pulchellum													1 .						- 1																			÷				÷
lon sp. drepanothrix *	. :		1		:					1				:			: :			1							ď.	: :						1	:	:	:	:				:
pha kernii aria arenaria																	1 .							. :		* *									-							÷
ria prehensilis			- 1		:					- 1						1	: :											: :					4			:	:			1		:
ria stanfieldii bussei	P				1				, •			-				: 1											:															
tylis abortiva		÷						- :		- :						1	p .					÷														1						
ylis pusilla sp.		-								P						:											1															-
s sp.		÷																	- 1		,			: :										:				p p				:
s tenuiculmis piaceae sp.	P											- 1		: -		:											:	: :					-		1	p						
gathis anobrya								- :				p				7	: :																									
cf. djurensis mia simplex										D			Р .	: -		:						:					:	: :							:							
thus maderaspatensis		-	ź							- 5		- :					. :		- 1							- :										p						÷
la multiflora sca foliosa	P				:	:										:												: :						1		p	:					
is pumilus												- :					р .		- 1					. :	:											-						
ia simplex aria tortilis			P							P		- :	:		:	:	: :										1															4.
ynomene lateritia		÷								-										*										r							1					
aria ochroleuca is sp.	r.					:	:	:	:					:		:		- :			:						1	;														:
fera deightonii phicarpa fistulosa											:	r						-									:															
chne rottboellioides								ŗ		:						:		. :				;																				
chne gracilis																																										
stemon graniticola														:			: :					1 1															ř					

TAYLOR, 1989) were found. Presumably, autogamy occurs in *Aeollanthus pubescens* (RYDING, 1986), Eriocaulaceae (STÜTZEL, 1981) and many Cyperaceae. The latter observation was made in the greenhouse where isolated (i.e. wrapped with plastic bags) specimens of Cyperaceae (*Bulbostylis* sp., *Fimbristylis* sp.) showed a high fruiting rate.

# Dispersal biology

Most species (54.8%) possess dehiscent capsular fruits. The number of diaspores per individual is frequently very high (up to several thousands, e.g. *Utricularia* spp., *Burmannia madagascariensis*). Among the diaspores the following groups can be distinguished: dust seeds (less than 0.5 mm in size), granule seeds (0.6-2 mm in size) and small fruits (e.g. nutlets). Dust seeds were dominating and characteristic for small annuals (e.g. *Drosera indica*, *Utricularia* spp., *Xyris straminea*). Micromorphological testa characters, like raised anticlinal walls (e.g. *Utricularia spiralis*), columnlike projections on anticlinal walls (*Eriocaulon togoense*) and epicuticular waxes (*Drosera indica*) reduce the sink velocity and enhance the water-repellance of the sometimes nauto-hydrochorous seeds (RAUH & al., 1975). Granule seeds and fruits (nutlets, caryopses) were the typical diaspores of many Poaceae and Cyperaceae.

Anemochory (75%) is the most frequent mode of dispersal. In contrast to tiny dust seeds the nearly smooth and relatively heavy granule seeds can probably only be dispersed by strong winds which occur on inselbergs due to upwinds and turbulences. Zoochory is often found as epiornithochory, e.g. diaspores lying in wet ground stick to the feet of visiting birds. Myxospermy occurs in *Rotala* spp. and *Aeollanthus pubescens*. Certain Poaceae and Cyperaceae may be endozoochorous.

Most species (77%) use several dispersal modes ("polychory"): dust seeds can be dispersed by wind, animals or water (28% of all species). Granule seeds can be dispersed by animals and wind (49%). Purely zoochorous were 19% of the species, 4% were purely anemochorous.

### Discussion

#### Abiotic factors

The existence of the EFV depends both on more or less continuous water supply from surrounding vegetation types (size of catchment area and storage capacity influence the area of the ephemeral flush community) during the rainy period and on the occurrence of a prolonged dry season. The shallowness of the soil in connection with its sandy or sandy-loamy texture results in a very low water storage capacity of the EFV-sites. Without water supply the soil dries out quickly. Therefore an essential prerequisite for the existence of this habitat is seepage water running down from other plant communities.

Dry periods of 1-2 weeks (sometimes even longer) during the rainy season occur more frequently in the savanna region than in the rain forest zone of the Ivory Coast. The risk of local extinction due to catastrophic drought is therefore higher for ephemeral flush species growing on savanna zone inselbergs. Surprisingly the EFV is often lacking respectively poor in species on inselbergs situated in the rain forest zone with their relatively high and more regularly distributed annual rainfall. Presumably a few perennial species outcompete a large number of annuals under these more favourable growth conditions. In contrast to this the harsher climatic conditions on savanna zone inselbergs favour the co-existence of less competitive annuals and thus increase the species richness of the EFV.

#### Succession

Although not directly observed, there are hints which indicate successional processes. On inselbergs situated in the rain forest zone ephemeral flush communities mostly occurred as small fringes around *Afrotrilepis*-mats respectively other open plant communities (e.g. shallow depressions, rock pools). It is conceivable that the EFV under these conditions forms a stage in a successional sere which will lead to an *Afrotrilepis*-community. In the savanna region, succession within the EFV towards a community dominated by perennials is regularly thrown back because of disturbing effects (i.e. unpredictable droughts, occasional fire). On average the species richness of ephemeral flush communities is higher on savanna zone inselbergs (up to 35 species per relevé) than on those in the rain forest region (up to 15 species per relevé). In addition in the savanna zone the area of ephemeral flush communities is generally larger in combination with an increased heterogeneity in soil depths within individual stands. We suppose that the higher species richness of ephemeral flush communities on savanna zone inselbergs is due to the combined effect of a certain degree of disturbance and a greater habitat heterogeneity than on rain forest inselbergs. A similar gradient in species richness has also been reported for plant communities in shallow depressions on Ivorian inselbergs (POREMBSKI & al., 1995).

### Phytosociology

Remarkable is the high number of species with low constance, i.e. the unpredictability of the species inventory of ephemeral flush communities which indicates a considerable amount of stochasticity.

For an exact phytosociological classification of the EFV more fieldwork is necessary. Preliminary data indicate that the EFV is floristically relatively uniform and may be unified as single phytosociological unity in West Africa (from Guinea to Nigeria). Close affinities exist to certain other communities, such as those on ferricretes. SCHNELL (1952) described an *Utricularieto-Eriocauletum pumili*-community from ferricretes in the Nimba Mountains (Guinea) which has a similar species inventory than many ephemeral flush communities. Presumably, the EFV belongs to the class *Eriocaulo-Utricularietea* which has been described from wet to shallowly inundated rock depressions on West African inselbergs respectively ferricretes (KNAPP, 1966). Moreover relations seem to exist to the class *Drosero-Xyridetea* described by SCHMITZ (1988) from Rwanda, Burundi and Zaïre. Even in the temperate zone ecologically similar therophyte-rich pioneer communities, like the class *Isoeto-Nanojuncetea* (MOOR, 1936; PIETSCH, 1973) occur. POTT (1992) characterizes them as small, often unstable pioneer communities on open soil and disturbed places.

### Life-forms and life-strategies

Obviously tiny annuals and to a much lesser extent hemicryptophytes respectively geophytes are particularly well adapted to cope with the environmental conditions (e.g. nutrient deficiency, unpredictable droughts during the rainy season) typical for ephemeral flush communities. Most other habitats on Ivorian inselbergs are also characterized by the dominance of therophytes (POREMBSKI & al., 1995; POREMBSKI & BROWN, 1995) what is indicative of extreme environmental conditions.

Therophytic species occurring in ephemeral flush communities are r-strategists which are weak competitors with a short life cycle and a high reproduction rate. This is in strong contrast to *Afrotrilepis*-mats where highly competitive and long-lived K-strategists dominate (POREMBSKI & al., 1995).

# Pollination and dispersal biology

The rarity of pollinator visits recorded in ephemeral flush communities could be an artefact of observation (i.e. time span of observation too short) or an effect of the comparatively small size and patchily distribution of this habitat which results in a decreased attractivity for flower visitors. Perhaps the presence of autogamous and agamospermous species is a direct consequence of a low frequency of pollinator visits.

The nutrient deficiency within ephemeral flush communities favours the production of very small diaspores. These, however, are produced in huge numbers which enhances the probability of dispersal over larger distances. Most ephemeral flush species seem to be relatively effective dispersers what is characteristic for colonizers of patchily distributed growing sites. In particular the production of polychorous dust seeds with combined anemochory, zoochory and nautohydrochory seems to be of importance concerning the effectiveness of dispersal.

#### Germination

Diaspores of many ephemeral flush species seem to have a dormancy, which prevents them from germination during the last weeks of the rainy season. Supposedly the diaspores need high temperatures during the dry season and subsequently a definite duration of permanent moisture (see HINTIKKA, 1990; MEYER & KITCHEN, 1992). This strategy prevents germination after sporadic rainfall in the dry season.

# Phytogeography and floristic affinities

Supposedly there is a relation between the large distributional area of many species found in ephemeral flush communities and their good dispersal ability. Phytogeographical affinities of the EFV are similar to those of other habitats on Ivorian inselbergs. Striking is the absence of endemics which is quite different from the situation in other tropical regions (for a discussion of possible reasons s. POREMBSKI & BROWN, 1995). In West Africa the EFV seems to be floristically uniform from Guinea to Nigeria (RICHARDS, 1957; HAMBLER, 1964; POREMSBKI & al., 1994) with minor differences on the species level. According to JAEGER & WINKOUN (1962) and KEITA & al. (1993), the EFV on ferricretes and rock outcrops on the plateau of Bandiagara (southern Mali) is floristically very similar to the Ivorian EFV. On inselbergs in the rain forest zone of Cameroon many characteristic annuals (e.g. *Drosera indica*, *Utricularia* spp.) are missing (LÉTOUZEY, 1968; VILLIERS, 1981; E. FISCHER, Bonn, pers. comm.), sometimes the whole community is absent (e.g. in Gabon, REITSMA & al., 1992). Similar ephemeral flush communities are also known from Malawi (POREMBSKI, 1996), Zimbabwe (R. SEINE, Bonn, pers. comm.), Madagascar (RAUH, 1973; E. FISCHER, Bonn, pers. comm.), India (BHARUCHA & ANSARI, 1963) and Australia (ORNDUFF, 1987). Own preliminary data from the neotropics indicate that ephemeral flush communities are poorer in species compared to their paleotropical counterparts.

No species recorded during this study is strictly restricted to the EFV. Most species occur in other seasonally available habitats, like marshy or swampy places and even ruderals (e.g. *Fimbristylis dichotoma*) are regularly present.

#### **ACKNOWLEDGEMENTS**

We acknowledge financial support by the Deutsche Forschungsgemeinschaft (Ba 605/4-2) and the Richard-Winter-Stiftung (Bergisch-Gladbach). We thank the Ivorian authorities (Direction Générale des Eaux et Forêts, Abidjan; Institut d'Écologie Tropicale, Abidjan) for the permission to conduct research. We are grateful to K. E. Linsenmair (Würzburg) and his working group for support during fieldwork. For the use of herbarium facilities we thank L. Aké Assi (Abidjan) and all collaborating specialists for their determinations: E. Fischer (Bonn), P. Goetghebeur (Gent), C. Puff (Wien) and H. Scholz (Berlin-Dahlem). Soil analyses were carried out by the Institute of soil science at the University of Bonn (W. Brümmer), water analyses were carried out by the Wahnbach-Talsperrenverband (J. Clasen, Siegburg). Comments on the manuscript respectively help in the field were provided by N. Biedinger, A. Gröger, B. Möseler, J.-P. Mund, R. Seine, J. Szarzynski and C. Westerkamp. N. Knittel and C. Fetchenhauer are acknowledged for a critical review of the English version of the manuscript.

#### REFERENCES

- ADJANOHOUN, E. (1964). Végétation des savanes et des rochers découverts en Côte-d'Ivoire centrale. *Mémoires ORS-TOM* 7: 1-178.
- AKÉ ASSI, L. (1984). Flore de la Côte-d'Ivoire: Etude descriptive et biogéographique, avec quelques notes ethnobotaniques. Thèse de Doctorat, Université d'Abidjan.
- ANHUF, D. (1994). Zeitlicher Vegetations- und Klimawandel in Côte-d'Ivoire. *In:* LAUER, W. (ed.), *Erdwissenschaftliche Forschung.* Vol. XXX: 7-299.
- BARKMAN, J.J., H. DOING & S. SEGAL (1964). Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. *Acta Bot. Neerl.* 13: 394-419.
- BARTHLOTT, W., A. GRÖGER & S. POREMBSKI (1993). Some remarks on the vegetation of tropical inselbergs: diversity and ecological differentiation. *Biogeographica* 69: 105-124.
- BERG, R. Y. (1983). Plant distribution as seen from plant dispersal: general principles and basic modes of plant dispersal. *In:* KUBITZKI, K. (ed.), *Dispersal and distribution*. Sonderbd. Naturwiss. Vereins Hamburg 7: 13-36.
- BHARUCHA, F. R. & M. Y. ANSARI (1963). Studies on the plant associations of slopes and screes of the Western Ghats, India. *Vegetatio* 11: 141-154.
- BONARDI, D. (1966). Contribution à l'étude botanique des inselbergs de Côte-d'Ivoire forestière. Diplôme d'études supérieures de sciences biologiques, Université d'Abidjan.
- BORNHARDT, W. (1900). Zur Oberflächengestaltung und Geologie Deutsch-Ostafrikas. Reimer, Berlin.
- BRAUN-BLANQUET, J. (1964). Pflanzensoziologie. 3rd ed., Springer, Wien.
- BREMER, H. & J. JENNINGS (eds.) (1978). Inselbergs/Inselberge. Z. f. Geomorph., N. F., Supplem. 31.
- DÖRRSTOCK, S., R. SEINE, S. POREMBSKI & W. BARTHLOTT (1995). First record of the American *Utricularia juncea* (Lentibulariaceae) from Africa. *Kew Bull*. In press.
- ELDIN, M. (1971). Le Climat. In: Le milieu naturel de la Côte-d'Ivoire. Mémoires ORSTOM 50: 73-108.
- FAEGRI, K. & L. VAN DER PIJL (1979). The principles of pollination ecology. 3rd ed., Pergamon Press, Oxford.
- FAO-UNESCO (1988). Soil map of the world. Rome.
- GUILLAUMET, J.-L. (1967). Recherches sur la végétation et la flore de la région du Bas-Cavally (Côte-d'Ivoire). Mémoires ORSTOM 20: 1-247.
- GUILLAUMET, J.-L. & E. ADJANOHOUN (1971). La végétation de la Côte-d'Ivoire. *In:* Le milieu naturel de la Côte-d'Ivoire. *Mémoires ORSTOM* 50: 157-263.
- HAMBLER, D. J. (1964). The vegetation of granitic outcrops in Western Nigeria. J. Ecol. 52: 573-594.
- HINTIKKA, V. (1990). Germination ecology and survival strategy of *Rumex acetosella* (Polygonaceae) on drought-exposed rock outcrops in South Finland. *Ann. Bot. Fennici* 27: 205-215.
- JAEGER, P. & D. WINKOUN (1962). Premier contact avec la flore et la végétation de Bandiagara. *Bull. IFAN (sér. A)* 24: 69-111.
- JENNY, M. (1994). Diasporenausbreitung in Pflanzengemeinschaften. Beitr. Biol. Pflanzen 68: 81-104.
- KEITA, A., C. DIAKITÉ, D. DIALLO, L. AKÉ ASSI & N. DIARRA (1993). Contribution aux études ethnobotaniques et floristiques. 25-115. In: KEITA, A. & P. COPPO (eds.), Plantes et remèdes du plateau Dogon. Éditions CRMT/PSMTM, Bandiagara, Perugia.
- KILLIAN, C. (1953). Contribution à l'étude de la biologie de quelques *Utricularia* tropicaux. *Bull. IFAN (sér. A)* 15: 72-82.

- KNAPP, R. (1966). Höhere Vegetations-Einheiten von West Afrika unter besonderer Berücksichtigung von Nigeria und Kamerun. 2nd ed., *Geobotanische Mitteilungen* 34.
- LENEUF, N. (1959). L'altération des granites calco-alcalins et des granodiorites en Côte-d'Ivoire forestière et les sols qui en sont dérivés. Thèse de Doctorat d'Etat, Université Paris.
- LÉTOUZEY, R. (1968). Étude phytogéographique du Cameroun. Lechevalier, Paris.
- MEYER, S. E. & S. G. KITCHEN (1992). Cyclic seed dormancy in the short-lived perennial *Penstemon palmeri*. *J. Ecol.* 80: 115-122.
- MILDBRAED, J. (1922). Wissenschaftliche Ergebnisse der Zweiten Deutschen Zentral-Afrika-Expedition 1910-1911. Band 2, Botanik. Klinkhardt & Biermann, Leipzig.
- MONNIER, Y. (1983). Végétation. *In:* VENNETIER, P. & G. LACRAVERE (ed. 2, *Atlas de la Côte-d'Ivoire*. Editions Jeune Afrique, Paris.
- MOOR, M. (1936). Zur Soziologie der Isoetalia. Beitr. Z. Geobot. Landesaufnahme Schweiz 20: 1-148.
- ORNDUFF, R. (1987). Islands on Islands: Plant Life on the Granite Outcrops of Western Australia. University of Hawaii, H. L. Lyon Arboretum Lecture No. 15. University Press of Hawaii, Honolulu.
- PIETSCH, W. (1973). Beitrag zur Gliederung der europäischen Zwergbinsengesellschaften (*Isoeto-Nanojuncetea* Br.-Bl. & Tx. 1943). *Vegetatio* 28: 401-438.
- POREMBSKI, S. (1996). Notes on the Vegetation of Inselbergs in Malawi. Flora 191: 1-8.
- POREMBSKI, S. & W. BARTHLOTT (1992). Struktur und Diversität der Vegetation westafrikanischer Inselberge. Geobot. Kolloq. 8: 69-80.
- POREMBSKI, S. & W. BARTHLOTT (1993). Ökogeographische Differenzierung und Diversität der Vegetation von Inselbergen in der Elfenbeinküste. 149-158. *In:* BARTHLOTT, W., C. M. NAUMANN, K. SCHMIDT-LOSKE & K.-L. SCHUCHMANN (eds.), *Animal-plant interactions in tropical environments.* Results of the Annual Meeting of the German Society for Tropical Ecology, Bonn 1992.
- POREMBSKI, S., W. BARTHLOTT, S. DÖRRSTOCK, & N. BIEDINGER (1994). Vegetation of rock outcrops in Guinea: granite inselbergs, sandstone table mountains and ferricretes-remarks on species numbers and endemism. *Flora* 189: 315-326.
- POREMBSKI, S., G. BROWN & W. BARTHLOTT (1995). Species diversity of plant communities in shallow depressions on Ivorian inselbergs. *Vegetatio* 117: 151-163.
- POREMBSKI, S. & G. BROWN (1995). The vegetation of inselbergs in the Comoé National Park (Ivory Coast). Candollea 50: 351-365.
- POTT, R. (1992). Die Pflanzengesellschaften Deutschlands. Ulmer, Stuttgart.
- RAUH, W. (1973). Über die Zonierung und Differenzierung der Vegetation Madagaskars. Akad. Wiss. Lit. Mainz. Trop. Subtrop. Pflanzenwelt 1: 1-146.
- RAUH, W., W. BARTHLOTT & N. EHLER (1975). Morphologie und Funktion der Testa staubförmiger Flugsamen. *Bot. Jahrb. Syst.* 96: 353-374.
- RAUNKIAER, C. (1934). The life-forms of plants and statistical plant geography. Oxford University Press, London.
- REITSMA, J. M., A. M. LOUIS & J. J. FLORET (1992). Flore et végétation des inselbergs et dalles rocheuses: première étude au Gabon. *Bull. Mus. Nat. Hist. Nat., B, Adansonia* 14: 73-97.
- RICHARDS, P. W. (1957). Ecological notes on West African Vegetation I. The plant communities of the Idanre Hills, Nigeria. *J. Ecol.* 45: 563-577.
- RYDING, O. (1986). The genus Aeollanthus s. lat. (Labiatae). Acta Univ. Ups. Symb. Bot. Ups. 26: 1-152.
- SCHMITZ, A. (1988). Révision des groupements végétaux décrits du Zaire, du Rwanda et du Burundi. Musée Royal de l'Afrique Centrale, Tervuren. *Annales Sciences Économiques* 17: 1-315.
- SCHNELL, R. (1952). Végétation et flore de la région montagneuse du Nimba. Mém. IFAN 22: 1-45.
- STÜTZEL, T. (1981). Zur Funktion und Evolution köpfchenförmiger Blütenstände, insbesondere der Eriocaulaceen. *Beitr. Biol. Pflanzen* 56: 439-468.
- TAYLOR, P. (1989). The genus Utricularia. Kew Bull. Additional Series 14. HSMO. London.
- VAN DER PIJL, L. (1982). The principles of dispersal in higher plants. 3rd ed., Springer, Berlin.
- VILLIERS, J. F. (1981). Formations climaciques et relictuelles d'un inselberg inclus dans la forêt dense camerounaise. Thèse de Doctorat d'Etat, Université Paris 6.
- WILMANNS, O. (1984). Ökologische Pflanzensoziologie. 2nd ed., Quelle & Meyer, Heidelberg.

Address of the authors: Botanisches Institut und Botanischer Garten der Universität, Meckenheimer Allee 170, D-53115 Bonn, Germany.

