

Zeitschrift: Candollea : journal international de botanique systématique =
international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 52 (1997)

Heft: 2

Artikel: A new species of Inula L. (Compositae-Inuleae) from Kashmir

Autor: Dawar, Rubina / Qaiser, Mohammad

DOI: <https://doi.org/10.5169/seals-879440>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A new species of *Inula* L. (Compositae-Inuleae) from Kashmir

RUBINA DAWAR

&

MOHAMMAD QAISER

RÉSUMÉ

DAWAR, R. & M. QAISER (1997). Une espèce nouvelle du genre *Inula* L. (Compositae-Inuleae) décrite du Cachemire. *Candollea* 52: 281-285. En anglais, résumés français et anglais.

Inula koelzii Dawar & Qaiser, nouvelle espèce du Cachemire, est décrite et illustrée.

ABSTRACT

DAWAR, R. & M. QAISER (1997). A new species of the genus *Inula* L. (Compositae-Inuleae) from Kashmir. *Candollea* 52: 281-285. In English, French and English abstracts.

Inula koelzii Dawar & Qaiser, new species from Kashmir, is described and illustrated.

KEY-WORDS: COMPOSITAE – *Inula* – Taxonomy – Kashmir.

During the generic revision of *Inula* L. from Pakistan and Kashmir, it was noted that STEWART (1972, p. 753), raising *Inula obtusifolia* var. *clarkei* Hook. f. to specific level, recognized *Inula clarkei* (Hook. f.) R. R. Stewart and treated *I. karakorensis* Clarke MSS. as the synonym of his new species. He has also cited several specimens under *I. clarkei*. The present authors had an opportunity to study most of the specimens and a careful examination of these specimens (viz. Baltistan: Ashkoley, *Clarke* 30418; Braldo river, *Hans Hartmann* s.n.; Dras valley, *Duthie* 13694; Ladakh: Kangi Nullah, *Koelz* 2818, *Koelz* 2827a; Zanskar: Rangdum, *Koelz* 2900a, *Koelz* 2900b) clearly indicated that it was a mixture and belonged to three different taxa. Some of the specimens, viz. Baltistan: Ashkoley, *Clarke* 30418 (BM!); Braldo river, *Hans Hartmann* s.n. (RAW!) belong to true *Inula clarkei*. They are characterized by the presence of 2-3 mm long, dense sericeous cypsellae, lower leaves sessile (shortly attenuate), stem and lower surface of leaves tomentose, outer involucral bracts densely pubescent, 8-14 mm long and uniseriate papillus bristles. While *Duthie* 13694 (BM!, E!) and *Koelz* 2900b (NY!) could not be placed within *I. clarkei*, as in these specimens stem and leaves are papillate-glandular and pilose with 4-6 mm long outer involucral bracts which are scarcely pubescent and glandular. Thus these specimens belong to *Inula obtusifolia* Kern. However, in the remaining three specimens: *Koelz* 2818, *Koelz* 2900a (RAW!, NY!) and *Koelz* 2827a (NY!), the lower leaves are long winged petiolate, cypsellae glabrous, 3-4 mm long and pappus with biseriate bristles. Therefore, these specimens do not correspond with *I. clarkei* or any other known species of *Inula* L. In order to accommodate them a new species is described here.




Fig. 1. – *Inula koelzii* Dawar & Qaiser: A, habit; B, involucral bracts (outer to inner); C, ray floret; D, disc floret; E, stamens, stigma & style; F, style showing stigmatic area and distribution of acute sweeping hairs.




Fig. 2. – Map showing the distribution of *Inula koelzii* Dawar & Qaiser (●); *I. clarkei* (Hook. f.) R. R. Stewart (○); *I. obtusifolia* Kern. (■) and *I. royleana* DC. (▲) in Pakistan and Kashmir.

Inula koelzii Dawar & Qaiser, spec. nova (Fig. 1).

Herba 23-24 cm alta, lax hirsuta et ± papilloso-glandulosa. Folia elliptica, marginibus serrato-glandulosis, superiora sessilia semiamplexicaulia basi auriculata apice acuminate acuta, inferiora longe petiolata obtuse acuta 11-15 × 3-5 cm. Capitula axillaria solitaria vel terminalia solitaria, 2-5 cm diametro. Involucrum 1.8-2.0 cm longum; bracteae 4-seriatae imbricatae, exteriores foliosae lanceolatae vel oblanceolatae acuminatae hirsutae, 18 × 3 mm, interiores membranaceae angustiores dorsaliter glandulosae marginibus puberulis. Cypsellae 3-4 mm longa oblonga quadrangularia glabra. Pappus aureus biseriatus basaliter connatus pilis numero 30-52, 5 mm longis.

Herb, 23-24 cm tall, lax hirsute and ± papillate-glandular. Leaves elliptic, margins serrate-glandular; upper sessile semiamplexicaul, base auricled, apex acuminate-acute; lower long petiolate, obtuse-acute, 11-15 × 3-5 cm. Capitula axillary solitary or solitary terminal, 2-5 cm in diameter. Involucrum 1.8-2.0 cm long, bracts 4-seriate, imbricate, outer leafy, lanceolate or oblanceolate, acuminate, hirsute, 18 × 3 mm, inner membranous, narrower, dorsally glandular with puberulous margins. Cypsellae 3-4 mm long, oblong, quadrangular, glabrous. Pappus golden yellow, biseriate, basally connate, 30-52 in number, 5 mm long.

Type: Kashmir: Kanga Nulla, below Kangi, Ladakh, in river beds, alt. 12.500', Sept. 5, 1931, Koelz 2818 (Holo- RAW!; Iso- NY!).

Paratypes: Kashmir: Kanga Nulla, above Kangi, Ladakh, alt. 13.000', Sept. 6, 1931, Koelz 2827a (NY!); Rangdam, Zanskar, on semi-arid slope, alt. 11.000', Sept. 12, 1931, Koelz 2900a (KUH!; NY!).

Etymology. – This species is named after the collector, Dr. Walter Norman Koelz, who has collected all the specimens known of this species.

Distribution. – Known from Ladakh (Kashmir) only (Fig. 2).

Ecology. – Grows on the semi-arid slopes and river beds at an altitude from 11.000-13.000'.

Fl. Per. – September.

Inula koelzii Dawar & Qaiser is closely related to *Inula royleana* DC. in having semiamplexicaul upper and long petiolate lower leaves; 1 or 2 capitula per plant; 3-4 mm long, glabrous and oblong cypellae. But it differs from *I. royleana* by having capitula upto 5 cm in diameter; outer involucral bracts lanceolate or oblanceolate, upto 3 mm wide and biseriate pappus bristles. Whereas, in *I. royleana* the capitula are 6-10 cm in diameter; outer involucral bracts ovate-obovate, 5-10 mm wide and pappus bristles uniserial. On the other hand the new species also relates with *I. clarkei* (Hook. f.) R. R. Stewart and *I. obtusifolia* Kern. by having solitary terminal capitulum, which is 2-5 cm in diameter. Outer involucral bracts lanceolate or oblanceolate, upto 3 mm wide and oblong cypellae. But it clearly differs from both species by having long petiolate lower leaves, 3-4 mm long and glabrous cypellae and biseriate pappus bristles. While in *I. clarkei* and *I. obtusifolia* lower leaves are sessile (shortly attenuate), cypellae 2-3 mm long and villosus-dense sericeous and pappus with uniserial bristles.

A synoptic key is given below for all the four related species.

1. Lower leaves long winged petiolate. Cypellae glabrous, 3-4 mm long 2
- 1a. Lower leaves without winged petiole. Cypellae villous-dense sericeous, 2-3 mm long .. 3
2. Outer involucral bracts ovate-obovate, 5-10 mm broad. Pappus bristles uniserial *I. royleana*
- 2a. Outer involucral bracts lanceolate or oblanceolate, upto 3 mm broad. Pappus bristles biseriate *I. koelzii*
3. Stem and lower surface of leaves tomentose. Outer involucral bracts 8-14 mm long, densely pubescent *I. clarkei*
- 3a. Stem and leaves papillate-glandular and pilose. Outer involucral bracts 4-6 mm long, scarcely pubescent and glandular *I. obtusifolia*

Inula koelzii seems to be confined to Ladakh only, whereas *I. obtusifolia* is widely distributed in Gilgit, Hazara, Ladakh and Kashmir. *I. clarkei*, *I. obtusifolia* and *I. royleana* are sympatric in distribution in most of the regions of Gilgit, Hazara and Kashmir, but *I. clarkei* and *I. royleana* do not reach upto Ladakh. Whereas, *I. obtusifolia* occupies a larger area upto Ladakh, where it is sympatric with *I. koelzii* (Fig. 2).

ACKNOWLEDGEMENTS

We are grateful to the directors/curators of BM, E, NY and RAW herbaria for sending the material on loan. Thanks are also due to Dr. Saood Omer (KUH) for his valuable suggestions and Dr. R. R. Mill, Royal Botanical Garden Edinburgh, for latin translation of the description.

REFERENCES

- STEWART, R. R. (1972). An annotated catalogue of the vascular plants of West Pakistan and Kashmir. In: NASIR, E. & S. I. ALI (eds.), *Flora of West Pakistan*. Karachi.

