

**Zeitschrift:** Candollea : journal international de botanique systématique = international journal of systematic botany  
**Herausgeber:** Conservatoire et Jardin botaniques de la Ville de Genève  
**Band:** 55 (2000)  
**Heft:** 1

**Artikel:** Nasturtium macrocarpum Boiss. transferred to Barbarea (Brassicaceae)  
**Autor:** al-Shehbaz, Ihsan A. / Jacquemoud, Fernand  
**DOI:** <https://doi.org/10.5169/seals-879515>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.05.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Nasturtium macrocarpum Boiss. transferred to Barbarea (Brassicaceae)

IHSAN A. AL-SHEHBAZ  
&  
FERNAND JACQUEMOUD

## ABSTRACT

AL-SHEHBAZ, I. A. & F. JACQUEMOUD (2000). *Nasturtium macrocarpum* Boiss. transferred to *Barbarea* (Brassicaceae). *Candollea* 55: 201-203. In English, English and French abstracts.

*Nasturtium macrocarpum* Boiss. is transferred to *Barbarea* as *Barbarea macrocarpa* (Boiss.) Al-Shehbaz & Jacquemoud. The generic boundaries between *Barbarea* and *Rorippa* are discussed.

## RÉSUMÉ

AL-SHEHBAZ, I. A. & F. JACQUEMOUD (2000). *Nasturtium macrocarpum* Boiss. transferred to *Barbarea* (Brassicaceae). *Candollea* 55: 201-203. En anglais, résumés anglais et français.

*Nasturtium macrocarpum* Boiss. est transféré dans le genre *Barbarea* sous le nom *Barbarea macrocarpa* (Boiss.) Al-Shehbaz & Jacquemoud, et ses affinités sont exposées. Les limites génériques entre *Nasturtium* et *Barbarea* sont discutées.

**KEY-WORDS:** *Nasturtium macrocarpum* – *Barbarea* – *Ceriosperma* – BRASSICACEAE – Lebanon.

The species in question was originally described by BOISSIER (1849) as *Nasturtium macrocarpum* Boiss. and was kept in *Nasturtium* R. Br. by BOISSIER (1867), POST (1896), POST & DINSMORE (1932), BOULOUMY (1930), THIEBAUT (1936), and SCHULZ (1933, 1936). It was transferred to *Rorippa* Scop. by MOUTERDE (1970) and retained in this genus by ZOHARY & al. (1980). Finally the species was transferred by GREUTER & BURDET (1983) to the genus *Ceriosperma* (O. E. Schulz) Greuter & Burdet and was maintained there by GREUTER & al. (1986, 1993), without any indication about the definition and typification of this genus. For the reasons discussed below, we believe that *Nasturtium macrocarpum* is best accommodated in *Barbarea* R. Br., and it does not belong to *Nasturtium*, *Rorippa*, or *Ceriosperma*.

*Barbarea macrocarpa* is characterized by having woody caudex with petiolar remains of previous seasons, pinnatifid basal leaves with terminal lobe considerably larger than lateral ones, auriculate, often subleathery cauline leaves, bright yellow flowers with ascending-spreading parts, stout fruiting pedicels more than or about half as wide as the fruit, conspicuous midvein extending the full length of the fruit valve, narrowly linear fruits attenuate to apex gradually ending in style, minute stigma narrower than the style, and uniseriate, dark reddish brown to blackish, and coarsely reticulate seeds. This combination of characters is found in almost all species of *Barbarea*.

Although several species of *Rorippa* resemble *Barbarea* in aspects of the foliage and flowers above, none has a woody caudex with petiolar remains, and all have slender fruiting pedicels mostly much narrower than half of the fruit width, fruit apex abruptly terminated into style, large stigmas wider than style, obscurely veined or veinless valves, and foveolate or rarely minutely reticulate, often yellowish to light brown seeds. The vast majority of *Rorippa* has globose to oblong or ovoid fruits with biseriate seeds, and narrowly linear fruits with uniseriate seeds are quite rare. In fact, some species of *Rorippa* that deviate from the above combination of characters are anomalous in the genus.

By assigning *Nasturtium macrocarpum* to any genus other than *Barbarea*, one immediately runs into the problem of finding its closest relative. In fact, BOISSIER (1867) pointed out (p. 182) that "Planta nexum inter *Nasturtium* et *Barbaream* praebens," but apparently subsequent authors overlooked this keen observation by Boissier. The species is most closely related to *B. minor* K. Koch and, to a lesser extent, to *B. brachycarpa* Boiss. On the basis of its treatment by COODE & CULLEN (1965), GREUTER & al. (1986) regard the two epithets as specific synonyms, *B. brachycarpa* Boiss. being the correct name, due to anteriority. Almost all other Middle Eastern and Caucasian works do not mention the species in their synonymy of *B. minor* (e.g., AVETISIAN, 1966; BOULOUMY, 1930; GROSSHEIM, 1950; HEDGE, 1968; MARDALEJ-SCHVILI, 1979; MOUTERDE, 1970; POST, 1896; POST & DINSMORE, 1932; RAGIMOV, 1953; THIEBAUT, 1936; ZOHARY & al., 1980. Neither did BUSCH (1909, 1939), describing a *B. minor* var. *eriopoda*, which COODE & CULLEN (1965) consider as a synonym of *B. brachycarpa*. Recently, HARTVIG & STRID (1987) added another variety to *B. minor* without taking *B. brachycarpa* in consideration in their comments. In our opinion, though very close, *B. minor* and *B. brachycarpa* may be distinguished, principally by their size, by the shape, pubescence and venation's patterns of their radical leaves, and their siliqua (rarely more than 4(5) loculate in *B. brachycarpa*, more than 5 to 7-8-10(14) loculate in *B. minor*).

Both *Barbarea macrocarpa* and *B. minor* are perennials with a distinct caudex, auriculate leaves, compressed fruits, and attenuate fruit apex terminated into a style to 3 mm long. From *B. minor*, *B. macrocarpa* is readily distinguished by having pectinate to finely pinnatisect cauline leaves, slender and flexuous infructescence rachises, and fruits (2.5-)4-6 cm long. By contrast, *B. minor* has toothed cauline leaves, stout and straight infructescence rachises, and fruits 1-2(-2.5) cm long.

*Barbarea macrocarpa* has no relatives whatsoever in *Rorippa* or *Nasturtium*. The latter genus consists of five species of aquatic plants with white flowers and lower nodes producing adventitious roots (AL-SHEHBAZ & PRICE, 1998).

*Ceriosperma* was initially described by SCHULZ (1933) as a section of *Nasturtium* with eight species all, except *N. macrocarpum*, restricted to Australia, Papua New Guinea, and Indonesia. SCHULZ (1933, 1936) did not typify the section, which was raised to the generic rank by GREUTER & BURDET (1983) without comments. Only *N. macrocarpum* was transferred by Greuter & Burdet as *C. macrocarpa* (Boiss.) Greuter & Burdet. In our opinion, JONSELL (1988) and HEWSON (1982) have correctly treated the members of sect. *Ceriosperma* (as enumerated by SCHULZ, 1933), excluding *N. macrocarpum*, in *Rorippa*. SCHULZ (1936) united *Rorippa* and *Nasturtium*, but he adopted that later-published *Nasturtium* for the combined genus.

***Barbarea macrocarpa* (Boiss.) Al-Shehbaz & Jacquemoud, comb. nova**

≡ *Nasturtium macrocarpum* Boiss., Diagn. Pl. Orient. 8: 18. 1849.

≡ *Ceriosperma macrocarpa* (Boiss.) Greuter & Burdet in Willdenowia 13: 86. 1983.

**Type:** [Lebanon], "Antilibani inter Hasbeya et Rascheya," May 1846, *E. Boissier s.n.* (Holo-, G-BOIS!; iso-, G-BOIS!, G!).

## ACKNOWLEDGEMENTS

The authors thank Patrick Perret and the anonymous reviewer for their valuable comments.

## REFERENCES

- AL-SHEHBAZ, I. A. & R. A. PRICE (1998). Delimitation of the genus *Nasturtium* (Brassicaceae). *Novon* 8: 124-126.
- AVETISIAN, V. E. (1966). Brassicaceae. In: TAKHTAJAN, A. L. (ed.), *Fl. Armenia* 5: 61-301. Yerevan.
- BOISSIER, E. (1849). *Diagnoses plantarum orientalium novarum* ser. 1, 8: 1-128. Paris.
- BOISSIER, E. (1867). *Flora orientalis*. Vol. 1. Basileae, Genevae.
- BOULOUMOY, L. (1930). *Flore du Liban et de la Syrie*. Paris.
- BUSCH, N. A. (1909). *Barbarea*. In: KUSNEZOW, N., N. BUSCH & A. FOMIN (ed.), *Fl. Caucas. Crit.* 3: 308-319. Urev.
- BUSCH, N. (1939). *Barbarea*. In: KOMAROV, V. L. (ed.), *Fl. CCCP* 8: 130-135. Moscow & Leningrad.
- COODE, M. J. E. & J. CULLEN (1965). *Barbarea*. In: DAVIS, P. H. (ed.), *Fl. Turkey* 1: 433-438. Edinburgh.
- GREUTER, W. & H. BURDET (1983). *Ceriosperma* (O. E. Schulz) Greuter & Burdet. In: GREUTER, W. & T. RAUS (eds.) - *Med-checklist notulae*, 7. *Willdenowia* 13: 79-99.
- GREUTER, W., H. BURDET & G. LONG (eds.) (1986). *Med-Checklist*. Vol. 3. Genève.
- GREUTER, W., R. K. BRUMMITT, E. FARR, N. KILIAN, P. M. KIRK & P. C. SILVA (eds.) (1993). Names in current use for extant plant genera. *Regnum Veg.* 129. Königstein.
- GROSSHEIM, A. A. (1950). *Flora Kavkaza*, ed. 2. Moscow & Leningrad.
- HEDGE, I. C. (1968). Cruciferae - Arabideae. In: RECHINGER, K. H. (ed.), *Fl. Iran*. 57: 193-218. Graz.
- HARTVIG, P. & A. STRID (1987). New taxa and new records from the mountains of SW and SC Turkey. *Bot. Jahrb. Syst.* 66: 91-102.
- HEWSON, H. J. (1982). Brassicaceae. In: BRIGGS, B. G. & al. (eds.), *Fl. Australia* 8: 231-357. Canberra.
- JONSELL, B. (1988). Cruciferae. In: VAN STEENIS, C. G. G. J. (ed.), *Fl. Males.*, ser. 1: 10: 541-560. Dordrecht.
- MARDALEJSCHVILI, T. K. (1979). *Barbarea*. In: KETSKHOVELI, N. N. (ed.), *Fl. Gruzii*, ed. 2, 5: 50-54. Tbilisi.
- MOUSTERDE, P. (1970). *Nouvelle flore du Liban et de la Syrie*. Vol. 2. Beirut.
- POST, G. E. (1896). *Flora of Syria, Palestine, and Sinai*. Beirut.
- POST, G. E. & J. E. DINSMORE (1932). *Flora of Syria, Palestine, and Sinai*. Ed. 2, vol. 1. Beirut.
- RAGIMOV, M. A. (1953). *Barbarea*. In: KARIAGYN, I. I. (ed.), *Fl. Azerbaijan* 4: 240-242. Baku.
- SCHULZ, O. E. (1933). Kurze Notizen über neue Gattungen, Sektionen und Arten der Cruciferen. *Bot. Jahrb. Syst.* 66: 91-102.
- SCHULZ, O. E. (1936). Cruciferae. In: ENGLER, A. & K. PRANTL (eds.), *Nat. Pflanzenfam.*, ed. 2, 17B: 227-658. Verlag von Wilhelm Engelmann, Leipzig.
- THIEBAUT, J. (1936). *Flore Libano-Syrienne*. Vol. 1. Cairo.
- ZOHARY, M., C. C. HEYN & D. HELLER (1980). *Conspectus Florae Orientalis*. Fascicle 1. Jerusalem.

---

Addresses of the authors: AS: Missouri Botanical Garden, P.O. Box 299, St. Louis, MO63166-0299, U.S.A.

FJ: Herbarium, Conservatoire et Jardin botaniques de la Ville de Genève, Case Postale 60, CH-1292 Chambésy/Genève, Switzerland.

