

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	58 (2003)
Heft:	2
Artikel:	Taxonomic novelties in the neotropical genus Quiina Aubl. (Quiinaceae)
Autor:	Schneider, Julio Valentin / Zizka, Georg
DOI:	https://doi.org/10.5169/seals-879313

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Taxonomic novelties in the neotropical genus *Quiina* Aubl. (Quiinaceae)

JULIO VALENTIN SCHNEIDER

&

GEORG ZIZKA

ABSTRACT

SCHNEIDER, J. V. & G. ZIZKA (2003). Taxonomic novelties in the neotropical genus *Quiina* Aubl. (Quiinaceae). *Candollea* 58: 461-471. In English, English and French abstracts.

During a forthcoming familial revision, five new species (*Q. attenuata* J. V. Schneid. & Zizka, *Q. berryi* J. V. Schneid. & Zizka, *Q. cidiana* J. V. Schneid. & Zizka, *Q. gentryi* J. V. Schneid. & Zizka, *Q. piresii* J. V. Schneid. & Zizka) of the neotropical genus *Quiina* Aubl. (Quiinaceae) are described and illustrated.

RÉSUMÉ

SCHNEIDER, J. V. & G. ZIZKA (2003). Nouveautés taxonomiques dans le genre néotropical *Quiina* Aubl. (Quiinaceae). *Candollea* 58: 461-471. En anglais, résumés anglais et français.

Dans le cadre d'une révision de la famille des Quiinaceae, cinq espèces nouvelles (*Q. attenuata* J. V. Schneid. & Zizka, *Q. berryi* J. V. Schneid. & Zizka, *Q. cidiana* J. V. Schneid. & Zizka, *Q. gentryi* J. V. Schneid. & Zizka, *Q. piresii* J. V. Schneid. & Zizka) du genre néotropical *Quiina* Aubl. sont décrites et illustrées.

KEY-Words: Malpighiales – Taxonomy.

Introduction

Quiinaceae is a small neotropical family of trees and shrubs, encompassing four genera (*Froesia*, *Lacunaria*, *Quiina*, *Touroulia*) and about 51 species. *Quiina* Aubl. is the largest genus of the family, comprising 31 species distributed from Belize to southern Brazil and Bolivia, including the Caribbean islands Jamaica and Trinidad, with the bulk of the species occurring in the Guayana and Amazon region. They are generally confined to lowland rain forests, often being locally abundant in the forest understorey, and rarely exceed altitudes of 1000 m.

Quiina is circumscribed and distinguished from the other genera by a combination of characters: simple, opposite leaves (except *Q. pteridophylla* (Radlk.) Pires with whorled leaves) with well developed intersecondary and plumose-reticulate tertiary veins (FOSTER, 1950), paired interpetiolar stipules, axillary inflorescences, androdioecious sex distribution, a syncarpous gynoecium, and a usually 2-locular ovary (SCHNEIDER, 1998; SCHNEIDER & al., 2002).

CODEN: CNDLAR

58(2) 461 (2003)

ISSN: 0373-2967

CONSERVATOIRE ET JARDIN

BOTANIQUES DE GENÈVE 2003

Material and methods

For the study of the plant material, specimens from the herbaria A, AAU, B, BM, BR, CEN, COL, F, G, GH, IAN, K, L, LPB, M, MO, NY, P, PORT, QCA, R, RB, S, SP, U, UB, VEN, W, WAG were investigated. The applied terminology is based on STEARN (1992) for general morphology and on HICKEY (1973) for leaf venation types.

Descriptions

Quiina attenuata J. V. Schneid. & Zizka, spec. nova (Fig. 1)

Type: PERU: Loreto, Prov. Requena, ad flumen Ucayali, 4°55'S 73°45'W, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200594) (arbre no. 5-133) (Holo-: G).

Affinis Q. parvifolia Lanj. & Heerdt, Q. berryi J. V. Schneid. & Zizka et Q. leptoclada Tul. sed ab Q. parvifolia stipulis angustioribus differt, ab Q. berryi sepalis majoribus, petiolo canaliculato differt, ab Q. leptoclada inflorescentiis longioribus, floribus pluribus, fructu elliptico vel globoso differt.

Trees, to 10-14 m tall; terminal internodes 0.3-1.3 × 0.1-0.2 cm, ± laterally compressed, longitudinally striate, densely pilose (trichomes < 0.1-0.5 mm long), glabrescent. Leaves opposite,




Fig. 1. – *Quiina attenuata* J. V. Schneid. & Zizka

A, terminal branchlet with fruiting inflorescences. **B**, single stipule with inconspicuous midvein. **C**, part of inflorescence with pedicel and flower bud. **D**, mature fruit with pedicel and part of peduncle. Figure drawn from *Bernardi s. n.* (G 200596) by Julio V. Schneider.

petiolate; stipules free, $3.5\text{-}9.5 \times 0.3\text{-}0.8$ mm, subulate (to elliptical-lanceolate), acute, \pm glabrous or inconspicuously pilose, \pm persistent, scars at subterminal node $0.3\text{-}0.4 \times 0.4\text{-}0.55$ mm, suborbicular-elliptical, joint or separated $< 0.1\text{-}0.3$ mm; petiole $1.0\text{-}3.0 \times 0.7\text{-}1.5$ mm, subcylindrical, dark, sparsely hairy, glabrescent; lamina $3.5\text{-}9.3(-14.0)$, juv.) $\times 1.3\text{-}2.4$ cm, chartaceous to subcoriaceous, elliptical to elliptical-obovate, apically shortly (sub-) acuminate, basally attenuate, adaxially glabrous, abaxially glabrous or along midvein sparsely pilose; margin minutely serrulate, inconspicuously revolute; secondary veins $6\text{-}10(-17)$, juv.) per side, on both sides \pm flat or prominulous (slightly more on adaxial surface), $0.2\text{-}1.2$ cm distant at about middle of lamina, separated by $0.2\text{-}(5)$ intersecondary veins. Hermaphroditic inflorescence, botryoid, 5-19-flowered, male 35-70-flowered, with partial inflorescences 3-6-flowered; peduncle $1.0\text{-}3.3$ cm $\times 0.6\text{-}0.9$ mm, (in flower) laterally compressed, inconspicuously longitudinally furrowed, shortly pilose (trichomes to 0.1 mm long), subcylindrical; pedicels articulated at base (up to 0.3 mm from base), the upper part $4.5\text{-}6.5 \times c. 0.3$ mm (in fruit thickened, 1-1.2 mm diameter), laterally compressed or cylindrical, glabrous; bracts to 0.5×0.3 mm, ovate, acute-acuminate, \pm glabrous. Flower c. 3 mm in diameter, globose in bud; sepals imbricate, 5, $0.9\text{-}1.0 \times 0.8\text{-}1.2$ mm, ovate to elliptical, rounded, ciliolate; petals imbricate, 5, $1.8\text{-}2.0 \times c. 2.0$ mm, suborbicular, glabrous, imbricate; stamens in hermaphroditic flowers 10, filaments c. 1.1 mm long, free, anthers c. 0.25×0.3 mm, subglobose; ovary 2-locular, ovules 2 per locule, basal-axile, styles 2, 1.0-1.2 mm long, \pm free, stigma oblique-subpeltate. Fruiting pedicel thickened. Fruits $1.4\text{-}1.5 \times 1.1\text{-}1.2$ cm, ellipsoid to subglobose, apically rounded, conspicuously longitudinally furrowed, glabrous; seeds 1 per fruit, c. 0.8×0.7 cm, subglobose, densely villous (trichomes 0.2-0.3 mm long).

Etymology. – The specific epithet refers to the attenuate leaf base of the species.

Distribution. – The species is only known from two localities, one in a lowland rain forest of Peru and one in lowland rain forests of the blackwater region of the Río Negro in southern Venezuela.

Ecology. – Associated with blackwater rivers.

Phenology. – No information available.

Additional material examined. – PERU: Loreto, Prov. Requena, ad flumen Ucayali, $4^{\circ}55'S$ $73^{\circ}45'W$, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200592) (arbre no. 6-R-96); Prov. Requena, ad flumen Ucayali, $4^{\circ}55'S$ $73^{\circ}45'W$, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200593) (arbre no. 2-10-6); Prov. Requena, ad flumen Ucayali, $4^{\circ}55'S$ $73^{\circ}45'W$, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200593; the same herbarium-no. as above, but from a different tree!) (arbre no. 7-15-4); Prov. Requena, ad flumen Ucayali, $4^{\circ}55'S$ $73^{\circ}45'W$, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200596) (arbre no. 5-133); Prov. Requena, ad flumen Ucayali, $4^{\circ}55'S$ $73^{\circ}45'W$, 120 m, VIII-IX.1976, L. Bernardi s. n. (G 200604) (arbre no. 3-R-202).

VENEZUELA: T. F. Amazonas, Dept. Casiquiare, Caño San Miguel, $2^{\circ}35'N$ $67^{\circ}07'W$, 160 m, 18.IV.1991, G. Aymard 8898 (MO); Dept. Río Negro, San Carlos de Río Negro, c. 20 Km S of confluence of Río Negro and Brazo Casiquiare, along Caño Cuweje, 32 Km S of San Carlos, $1^{\circ}56'N$ $67^{\circ}03'W$, 119 m, 25.IX.1980, H. L. Clark 7678 (MO, NY).

All Peruvian collections are from the same locality. The enumeration of the specimens is somehow misleading, because there is no collection number, and tree numbers and herbarium numbers are not unique to one collection. There are two collections of *Bernardi* s. n. with tree number 5-133. The holotype (G 200594) is a fruiting specimen, while the specimen G 200596 has flower buds. From the herbarium number G 200593 two collections of different tree numbers are known. The flowering specimen Clark 7678 (male flower buds) only differs slightly in larger bracts.

Similar species are *Q. parvifolia* Lanj. & Heerdt, *Q. leptoclada* Tul., and *Q. berryi* J. V. Schneid. & Zizka (see below). In *Q. parvifolia* stipules are elliptical and much broader (> 1.5 mm wide), and leaf blades are generally subcordate to acute, rarely attenuate. *Quiina berryi* is characterised by constantly 4 (larger) sepals, and an obconically widened pedicel in fruiting plants. *Quiina leptoclada* is distinguished by its short, mostly 3-flowered inflorescences, small obovoid fruits and conspicuously chartaceous leaf blades (even in adult plants).

***Quiina berryi* J. V. Schneid. & Zizka, spec. nova (Fig. 2)**

Type: BRAZIL: Amazonas, Km 159, BR 174, Manaus-Caracaraí Road, 20.IX.1974, G. T. Prance & al. 22717 (Holo-: R; iso-: F, K, NY, photo FR).

Affinis Q. amazonica A. C. Sm. et Q. negrensis A. C. Sm., sed ab Q. amazonica inflorescencia breviore foliis crassioribus differt, ab Q. negrensis folia nervis secundariis supra planis vel impressis, flore masculino trifasciculato.

Tree, to 15(-20) m tall; terminal internodes $0.3\text{-}8.7 \times 0.08\text{-}0.15\text{(-}0.2)$ cm, laterally compressed or subcylindrical, longitudinally furrowed, \pm pilose-pubescent (trichomes to 0.8 mm long). Leaves opposite, petiolate; stipules free, $(0.18\text{-})0.25\text{-}1.0\text{(-}1.7)$ \times 0.03-0.1 cm, subulate (to subulate-elliptical), acute, adaxially \pm glabrous, rarely puberulent, abaxially glabrous or sparsely to densely pilose-pubescent, with trichomes to 0.6(-1.0) mm long, scars at subterminal node $0.3\text{-}0.5 \times 0.4\text{-}0.8$ mm, elliptical, joint or separated to 0.3 mm; petiole $0.1\text{-}0.6 \times 0.07\text{-}0.15\text{(-}0.2)$ cm, canaliculate or not; lamina $3.3\text{-}12.5 \times 1.3\text{-}5.3$ cm, chartaceous to coriaceous, elliptical (to elliptical-obovate), apically (caudate-) acuminate, basally acute, cuneate or attenuate, adaxially glabrous, abaxially glabrous or pubescent along midvein, rarely covering the whole surface; margin minutely serrulate, rarely entire, flat or inconspicuously revolute; secondary veins 7-15 per side, adaxially level with surface or impressed, rarely prominulous, abaxially prominulous, 0.2-1.7 cm distant at about middle of lamina, separated by 0-2(-4) intersecondary veins. Hermaphroditic inflorescence botryoid, 1-13-flowered, male inflorescence 10-24-flowered, with partial inflorescences 3-flowered; peduncle $0.1\text{-}1.4 \times 0.03\text{-}0.06$ cm, laterally compressed or cylindrical, longitudinally inconspicuously furrowed-costate, glabrous or sparsely pubescent; bracts $0.7\text{-}1.4\text{(-}2)$ $\times 0.5\text{-}1.0$ mm, \pm ovate, acute; pedicel below articulation to 0.2-1.4 mm long, above articulation $2.2\text{-}9 \times 0.2\text{-}0.6$ mm, (sub-)cylindrical, distally inconspicuously to obconically widened, glabrous. Flowers 3 mm in diameter; flower bud subglobose; sepals imbricate, 4, $0.9\text{-}2.2 \times 1.0\text{-}2.3$ mm, ovate, elliptical, suborbicular or obovate, apically obtuse to rounded, marginally ciliolate, adaxially and abaxially glabrous, coriaceous; petals imbricate, free, in hermaphroditic flowers rarely connate at base, 4, $1.9\text{-}3.5 \times 1.2\text{-}1.9$ mm, obovate to elliptical-obovate, apically rounded, glabrous, membranous; stamens in hermaphroditic flowers 8-10, in male flowers 12-29, filaments $(0.4\text{-})1\text{-}1.8$ mm long, in hermaphroditic flowers sometimes adnate to petals, anthers $0.3\text{-}0.4 \times 0.3\text{-}0.4$ mm, ellipsoid, subglobose; ovary 2-locular, ovules 2 per locule, placentation basal-axile, styles 2, free, 0.9-2.0 mm long. Fruiting peduncle thickened, to 1.5 mm in diameter; fruiting pedicel thickened, apically obconically widened, to 1.2 mm in diameter. Fruit $1.1\text{-}2.1 \times 0.8\text{-}1.5$ cm, broadly ellipsoid to subglobose, longitudinally furrowed by underlying lacunes, glabrous, pericarp fibrous; seeds 1-2 per fruit, $0.9\text{-}1.25 \times 0.5\text{-}0.7$ cm, ellipsoid, densely villous, trichomes 0.2-1.5 mm long.

Etymology. – This species is named in honour of the U.S. botanist Paul E. Berry.

Distribution. – Widespread throughout the Brazilian Amazon region.

Ecology. – Found in the understorey of terra firme or caatinga lowland forests on sandy-clayey, yellow lateritic soils.

Phenology. – The principal flowering period lasts from July to December, while fruiting is throughout the year.

Additional material examined. – **BRAZIL:** Amazonas, Município de Novo Aripuña, BR 230, Rod. Transamazônica a 400 km de Humaitá, Rio Juma, vicinal Corujá a 30 km da Transamazônica, $7^{\circ}15'S$ $60^{\circ}00'W$, 3.V.1985, Cid Ferreira 5999 (FR, NY, UB); Município Pres. Figueiredo, Canteiro de obras da Hidrelétrica de Balbina, rio Uatumá, $1\text{-}2^{\circ}S$ $59\text{-}60^{\circ}W$, 12.III.1986, Cid Ferreira & al. 6759 (NY); Município Presidente Figueiredo, Canteira de obras de UHE de Balbina, $1^{\circ}30'\text{-}2^{\circ}00'S$ $59^{\circ}30'\text{-}60^{\circ}00'W$, 11.IX.1986, Cid Ferreira & al. 8075 (FR, NY); Distrito Agropecuário, Reserva 1501-Km 41 da Smithsonian/INPA, $2^{\circ}24'26''\text{-}2^{\circ}25'31''S$ $59^{\circ}43'40''\text{-}59^{\circ}45'50''W$, 50-125 m, 16.VII.1990, Freitas & al. 9 (NY); Rio Urubu, São José das Pedras, 22.II.1950, R. L. Fróes 26066 (IAN); Região do Rio Madeira, Rio Canuma, 5.XI.1957, R. L. Fróes 33770 (UB); Município de Borba, BR 230 Estrada Transamazonica, 5 km E of Sucunduri, $6^{\circ}50'S$ $59^{\circ}00'W$, 7.V.1985, Henderson & al. 338 (F, K, NY); Município de Manaus, c. 90 km NNE de Manaus, Distrito Agropecuário da Suframa, $2^{\circ}24'26''\text{-}2^{\circ}25'31''S$ $59^{\circ}43'40''\text{-}59^{\circ}45'50''W$, 50-125 m, 10.XI.1989, Kukle 22 (K, NY); Estrada Manaus-Itacoatiara, Km 108, 20.XI.1968, Lima s. n. (UEC); Estrada Manaus-Caracaraí, km 132, 13.II.1974,




Fig. 2. – *Quiina berryi* J. V. Schneid. & Zizka

A, terminal branchlet with fruiting inflorescence. **B**, detail of leaf margin with minute serrulation. **C**, stipules at terminal node. **D**, inflorescence with flower buds. **E**, mature fruit. Figures **A**, **B**, **C**, **E** drawn from Santos & Silva 4651, **D** from Fróes 33770 by Julio V. Schneider.

Loureiro & al. s. n. (INPA-No. 47898) (F); Distrito Agropecuário, Reserve 1501 (km 41) of the Smithsonian/INPA Biological Dynamics of Forest Fragments Project, 2°24'26"-2°25'31"S 59°43'40"-59°45'50"W, 50-125 m, 26.VII.1990, *Mori & Costa Lima Assunção* 21488 (FR, NY); Distrito Agropecuário, Reserva 1501 (Km 41) of the WWF/INPA MCS Project, 2°24'26"-2°25'31"S 59°43'40"-59°45'50"W, 50-125 m, 27.VII.1989, *Mori & al.* 20645 (FR, NY); Distrito Agropecuário, 90 Km NNE de Manaus, Reserva 1501(Km 41) da Smithsonian/INPA, Projeto Dinâmica Biológica dos Fragmentos Florestais, 2°24'26"-2°25'31"S 59°43'40"-59°45'50"W, 50-125 m, 3.XII.1991, *Oliveira & al.* 253 (NY); Distrito Agropecuário, 90 km NNE de Manaus, Reserva 1501 (km 41), Projeto Dinâmica Biológica de Fragmentos Florestais, 2°24'26"-2°25'31"S 59°43'40"-59°43'50"W, 50-125 m, 11.I.1992, *Oliveira & al.* 303 (NY); Distrito Agropecuário, 90 km NNE de Manaus, Reserva 1501 (km 41), 2°24'26"-2°25'31"S 59°43'40"-59°43'50"W, 50-120 m, 10.II.1993, *Oliveira & al.* A979 (NY); Distrito Agropecuário, Fazenda Porto Alegre, Reserva 3402 (Cabo Frio) of the WWF/INPA MCS project, 2°25'25"S 59°54'38"W, 50-125 m, 28.I.1989, *Pacheco & al.* 157 (K, NY); Manaus-Itacoatiara Highway, Reserva Florestal Ducke, km 26, 9.IX.1967, *G. T. Prance & al.* 2235 (FR, IAN, NY, R); Manaus-Itacoatiara Highway, Reserva Florestal Ducke, km 28, 20.III.1967, *G. T. Prance & al.* 4694 (FR, IAN, NY, R); Vicinity of Manaus, Reserva Florestal Ducke, Manaus Itacoatiara road, km 27, 13.III.1969, *G. T. Prance & al.* 10414 (F, IAN, K, NY, R); Reserva Florestal Ducke, 24.VII.1963, *Rodrigues* 5461 (F); Reserva Florestal Ducke, 28.IV.1964, *Rodrigues & Loureiro* 5802 (F); Reserva Florestal Ducke, 13.XII.1963, *Rodrigues & Coêlho* 5602 (F); Reserva Florestal Ducke, igarapé do Barro Branco, 13.V.1963, *Rodrigues & Coêlho* 5206 (NY); **Pará**, Tapajós, Boa Vista, 11.IV.1933, *Capucho* 569 (F, IAN); Santarém, km 67 da estrada para a Cachoeira do Palhão no rio Curuá Una, 1.XII.1966, *Ca valcante & Silva* 1494 (IAN); Município Obidos, rio Jaramacaru, próximo aos Campos do Ariramba, 1°10'S 55°35'W, 5.XII.1987, *Cid Ferreira* 9781 (F, GH, NY); Obidos, 15.VII.1918, *A. Ducke s. n.* (RB-No. 18131) (K, P); Rio Curuaúna, região do planalto de Santarém, 14.XI.1954, *R. L. Fróes* 31390 (IAN); Beira do Rio Curuatinga, Planalto de Santarém, 14.II.1955, *R. L. Fróes* 31592 (IAN); Igarapé Cuçári, região do Planalto de Santarém, IV.1955, *R. L. Fróes* 31757 (IAN); Região do Planalto de Santarém, V.1955, *R. L. Fróes* 31837 (IAN, NY, P); Cabeceiras do Rio Urucá, Município de Prainha, V.1955, *R. L. Fróes* 31857 (IAN); BR 230, Transamazon Highway, 53 km NE of Itaituba, 28.XI.1977, *G. T. Prance & al.* 25835 (NY, RB); Rio Trombetas, planalto Saracá, 27.V.1978, *Silva & Santos* 4651 (F, NY, RB); Rondonia, Eixo JP-14 e arredores, 10.X.1986, *Romanuc Neto & al.* 562 (SP).

This species is most similar to *Q. amazonica* A. C. Sm. and *Q. negrensis* A. C. Sm. The first is distinguished by larger inflorescences usually with > 10 flowers in hermaphroditic and > 30 flowers in male inflorescences, and the peduncle being generally > 1.5 cm long. Additionally, its fruits tend to be obovoid rather than ellipsoid, the leaf apex is more conspicuously caudate, and the leaf texture is consistently chartaceous. *Q. negrensis* differs in adaxially more prominent secondary veins and in staminate flowers being disposed in fascicles of 5-11. For the discrimination of *Q. attenuata* see above.

Quiina cidiana J. V. Schneid. & Zizka, spec. nova (Fig. 3)

Type: BRAZIL: Amazonas, Alto rio Solimões, Município de São Paulo de Olivença, platô ao sul da cidade, estrada para a localidade Bom Fim, 24.XI.1986, *Cid Ferreira & al.* 8513 (Holo-: NY, photo FR; iso-: F).

Propter folia petiolis brevibus Q. tinifolia Planch. & Triana affinis, sed foliis rigidis, inflorescentia fructibus majoribus differt.

Tree, 12-18 m tall; terminal internodes 0.5-1.2(-2.5) × 0.2-0.4 cm, laterally compressed or subcylindrical, even or longitudinally furrowed, glabrous or with scattered trichomes (0.2-0.5 mm long); leaves opposite, petiolate; stipules free, 0.5-1.5 × 0.1-0.55 cm, narrowly elliptical to ovate, acute, basally acute, thickened, glabrous or abaxially with scattered trichomes; petiole 0.15-0.9 × 0.13-0.3 cm, subcylindrical, thickened towards base; lamina 3.2-15.8 × 1.7-6.6 cm, coriaceous, stiff, obovate, apically shortly sub acuminate, obtuse, rounded or emarginate, basally (narrowly) rounded to acute, adaxially glabrous, abaxially glabrous or with scattered trichomes (0.2-1.0 mm long) along midvein; margin entire or minutely serrulate, revolute; secondary veins 9-16 per side, adaxially prominent to prominulous, abaxially prominent, 0.2-1.8 cm distant at about middle of lamina, separated by 0-1(-3) intersecondary veins. Hermaphroditic inflorescence botryoid, 22-25-flowered, male inflorescence not seen; peduncle 5.0-10.7 × 0.15-0.21 cm, laterally compressed or subcylindrical, longitudinally furrowed, densely pilose; bracts to 0.6 × 0.6 mm, ovate, sub acuminate to acute; pedicel articulated, below articulation to 0.8-1.9 mm long, above articulation 6-12 × 1.2-1.3 mm, cylindrical, distally narrowly widened, glabrous or densely pilose. Flowers not seen; sepals 4-5, 0.8-1.2 × 1.1-1.9 mm, ovate to elliptical, obtuse to rounded, marginally ciliolate, abaxially glabrous or sparsely pilose, coriaceous; petals not seen; stamens




Fig. 3. – *Quiina cidiana* J. V. Schneid. & Zizka

Terminal branchlet with fruiting inflorescence. Figure drawn from Cid & al. 8513 by Julio V. Schneider.

not seen; ovary 2-locular; styles 2. Fruit berrylike, 2.3-2.7 × 1.3-1.6 cm, ellipsoid-ovoid, apically acute, longitudinally furrowed by underlying lacunes, glabrous, pericarp fibrous; seeds not seen.

Etymology. – This species is dedicated to the Brazilian botanist C. A. Cid Ferreira, explorer of the Amazon flora.

Distribution. – Only known from the Brazilian Amazon region.

Ecology. – Known from Caatinga forests on white sand.

Phenology. – No information available.

Additional material examined. – BRAZIL: Amazonas, Upper Rio Solimões, Município Jutaí, Jutaí, 10 km S of town center, 2°45'S 66°45'W, 29.X.1986, Daly & al. 4181 (NY).

This species resembles *Q. tinifolia* Planch. & Triana in its shortly petiolated leaves. But the unique combination of stiffly coriaceous leaf texture, ovate(-elliptical) stipules and densely hairy, long inflorescences with large fruits distinguish it clearly from other species.

Quiina gentryi J. V. Schneid. & Zizka, spec. nova (Fig. 4)

Type: PERU: Madre de Dios, Prov. Manu, Parque Nacional del Manu, Río Manu, Cocha Cashu Station, 350 m, 26.III.1981, Foster & Janson 8327 (Holo-: F, no. 1945128, 1945129, two sheets).

Affinis Q. grandifolia Tessmann, sed ab *Q. grandifolia* inflorescentia axe singulari differt.

Trees or shrubs, 1.5-15 m tall; terminal internodes 5.8-6.8 × 0.33-0.5 cm, laterally compressed or subcylindrical, longitudinally furrowed, glabrous or densely pilose (trichomes 0.1 mm long); leaves opposite, petiolate; stipules free, ± caducous, 1.8-3.2 × 0.65-1.4 cm, ovate, acute to




Fig. 4. – *Quiina gentryi* J. V. Schneid. & Zizka

A, branchlet with leaf and interpetiolar stipules. **B**, apical part of inflorescence with young fruit. **C**, mature fruit. **D**, stipule with venation. Figures **A**, **D** drawn from Foster & Janson 8327, **B** from Gentry & al. 27081, **C**, from Dodson & Gentry 6591 by Julio V. Schneider.

acuminate, basally rounded, asymmetrical, with conspicuous lateral veins, on both surfaces glabrous or basally pilose; petiole $0.4-1.2 \times 0.3-0.5$ cm, subcylindrical, canaliculate or not, \pm thickened; lamina $26.5-34 \times 10-12.8$ cm, (chartaceous-) subcoriaceous, elliptical, apically acuminate, basally acute or narrowly rounded, adaxially glabrous or sparsely pilose along midvein, abaxially sparsely pilose-pubescent along midvein; margin entire or minutely serrulate, flat; secondary veins 14-29 per side, adaxially prominent to prominulous, abaxially prominent, 0.8-3.7 cm distant at about middle of lamina, separated by 0-1(-2) intersecondary veins. Hermaphroditic inflorescence botryoid, 29-36-flowered, with flowers multiseriate arranged, male inflorescence not seen; peduncle $3.4-6.7 \times 0.14-0.5$ cm, cylindrical, longitudinally furrowed, densely pilose; bracts $1.1-1.2 \times 0.7-0.8$ mm, ovate, acute; pedicel articulated, below articulation to 2-4.5 mm long, above articulation $2-4 \times 0.9-2.3$ mm, cylindrical, glabrous. Flowers not seen; sepals 4-5, $1.0-2.4 \times 1.0-2.8$ mm, ovate to elliptical, obtuse to rounded, marginally ciliolate, coriaceous; petals not seen; stamens not seen; ovary 2-locular, ovules 2 per locule, placentation basal-axile; styles 2; fruiting peduncle conspicuously thickened, to 5 mm in diameter; fruiting pedicel thickened, to 2.3 mm in diameter. Fruit berrylike, $(2.3)-3.3 \times (0.9)-1.6$ cm, ellipsoid-ovoid to ellipsoid, apically obtuse, longitudinally furrowed, glabrous, pericarp fibrous; seeds 1-2 per fruit, 1.6×1.15 cm, ellipsoid, densely villous, trichomes 0.2-0.6 mm long.

Etymology. – This species is dedicated to Alwyn H. Gentry (1945-1993).

Distribution. – Distributed in lowland to premontane rain forests in Peru and Ecuador.

Ecology. – No information available.

Phenology. – No information available.

Additional material examined. – PERU: Madre de Dios, Prov. Manu, Cocha Cashu Camp, Manu National Parc, 380 m, 21.X.1979, Gentry & al. 27081 (F).

ECUADOR: Los Ríos, Km 12 road from Patricia Pilar to 24 de Mayo, Montañas de Ila, 540 m, 7.X.1976, Dodson & Gentry 6591 (NY).

This species is closely related to *Q. grandifolia* Tessmann, but differs in adjacent stipules being separated (not fused into one as in *Q. grandifolia*). Both species most likely differ also in inflorescence characters, but so far only hermaphroditic inflorescences have been seen in *Q. gentryi*, only male ones in *Q. grandifolia*. In the first, flowers are borne along a long peduncle, while in the second, flowers appear on much shorter axes with the inflorescence looking circular in outline. Since heteromorphy between different sexual morphs is known in *Quiina*, distinction of the species based on inflorescences may be premature, albeit such a strong contrast in inflorescence structure has not been observed between different sexes in a species of the genus up to now.

The Ecuadorian collection (Dodson & Gentry 6591) differs slightly in having unthickened fruiting pedicels and more secondary veins. But for the overall habit it is similar and included in the diagnosis.

Quiina piresii J. V. Schneid. & Zizka, spec. nova (Fig. 5)

Type: BRAZIL: Amazonas, Município de Manaus, Distrito Agropecuário da SUFRAMA, Rodovia BR-174, km 72 depois 6 km oeste da BR, Fazenda Dimona, Terra firme, $2^{\circ}19'S$ $60^{\circ}05'W$, 13.V.1985, Pereira & al. s. n. (INPA/WWF 2303.1664) (Holo-: NY, photo FR).

Affinis Q. yatuensis J. V. Schneid. & Zizka, Q. integrifolia Pulle et Q. obovata Tul., sed ab Q. yatuensis folia nervis secundariis infra prominentibus, margine minutissime serrulata, fructu angustae obovoideo differt, ab Q. integrifolia inflorescentia pilosa, foliis margine minutissime serrulata differt, ab Q. obovata stipulis brevibus, triangulo-ovatis differt.

Tree to 11-15 m tall; terminal internodes $0.6-5.9 \times 0.15-0.25$ cm, subcylindrical, \pm even or inconspicuously furrowed, sparsely pilose (trichomes 0.1-0.5 mm long), glabrescent. Leaves opposite; stipules $5.0-8.0 \times 1.2-1.5$ mm, triangular-ovate, acute, thickened at base, ciliolate, caducous, scars at subterminal node $1.3-1.7 \times 1.2-1.4$ mm, elliptical to suborbicular, joint;




Fig. 5. – *Quiina piresii* J. V. Schneid. & Zizka

A, terminal branchlet with fruiting inflorescences. B, young fruit with style remnants. C, flower bud with ciliolate sepals. D, stipules at terminal node. Figure drawn from Freitas & al. 67 by Julio V. Schneider. Scale bars in A = 1 cm, B = 5 mm, C = 1 mm, D = 5 mm.

petiole $0.15\text{-}0.8 \times 0.13\text{-}0.3$ cm, subcylindrical, basally thickened, dark, when old often transversally furrowed; lamina $(2.8\text{-})5.0\text{-}16.2 \times (1.4\text{-})1.8\text{-}4.4$ cm, coriaceous, (elliptical-) obovate, apically (sub-) acuminate, basally tapering, adaxially glabrous, abaxially pilose along midvein, glabrescent; margin minutely serrulate, revolute; secondary veins 6-18 per side, adaxially prominent, abaxially flat or prominulous, 0.3-1.6 cm distant at about middle of lamina, separated by 0-1(-3) conspicuous intersecondary veins. Hermaphroditic inflorescence botryoid, 5-9-flowered, often basally branched, male inflorescence not seen; peduncle $5\text{-}20 \times 0.55\text{-}0.8$ mm, laterally compressed (to subcylindrical), inconspicuously furrowed, densely pilose; pedicels articulated, below articulation to 1.3 mm long, above articulation $5\text{-}8 \times 0.25\text{-}0.35$ mm, subcylindrical, glabrous; bracts to $1.9\text{-}4.0 \times 0.5\text{-}0.6$ mm, (elliptical-) ovate, acute, pilose. Flowers globose in

bud; sepals imbricate, (4-)5, 0.9-1.9 × 0.6-1.9 mm, ovate to suborbicular, rounded to obtuse, ciliolate, abaxially pilose or glabrous, coriaceous; petals (only seen in bud) 4-5 (?), ± obovate, apically rounded, membranous; stamens in male flowers c. 22, in hermaphroditic flowers 12-14, filaments free, anthers subglobose, c. 0.4-0.5 × 0.4-0.5 mm; ovary 2-locular, ovules 2 per locule, placentation basal-axile, styles 2, c. 1.4 mm long, ± free, stigma oblique-subpeltate, reniform. Fruit (inmature) to 1.0 × 0.4 cm, narrowly oblong-obovoid, longitudinally striate, glabrous; seed 1 per fruit, ovoid, densely villous.

Etymology. – This species is named in memory of the Brazilian botanist João Murça Pires (1917-1994) for his significant contributions to our knowledge on *Quiinaceae*.

Distribution. – Scattered in the Amazon region.

Ecology. – This species grows in lowland terra firme forests on yellow lateritic soils.

Phenology. – No information available.

Additional material examined. – BRAZIL: Amazonas, Distrito Agropecuário, Reserva 1501, km 41 da Smithsonian/INPA, 2°24'26"–2°25'31"S 59°43'40"–59°45'50"W, 50-125 m, 15.IX.1990, Freitas & al. 67 (NY); Município de Manaus, Distrito Agropecuário da SUFRAMA, Rodovia BR-174, km 64 depois 21 km leste na ZF3, Fazenda Porto Alegrê, 2°00'S 59°00'W, 18.III.1987, Sothers & al. s. n. (INPA/WWF 3402.598.2) (NY).

Quiina yatuensis J. V. Schneid. & Zizka is similar, but differs in secondary veins which are more prominent on the abaxial surface, the entire leaf margin, a more regular, opposite arrangement of the flower bracts, and more finely striate internodes. Additionally, fruits are broadly obovoid rather than narrowly oblong-obovoid as in *Q. piresii*. *Quiina integrifolia* Pulle can be of similar leaf shape, but is characterised by densely tomentose inflorescences, entire leaf margin, and ± globose fruits. Small-leaved *Q. obovata* Tul. is distinguished by long, narrowly elliptical to subulate, persistent stipules.

ACKNOWLEDGEMENTS

The *Quiinaceae* studies have been partly supported by a grant from the Hermann-Willkomm-Foundation and by the Deutsche Forschungsgemeinschaft (Zi 557/3-1). We further would like to thank the directors of the following herbaria for the loan of their specimens: A, AAU, B, BM, BR, CEN, COL, F, G, GH, IAN, K, L, LPB, M, MO, NY, P, PORT, QCA, R, RB, S, SP, U, UB, VEN, W, WAG.

BIBLIOGRAPHY

- FOSTER, A. S. (1950). Morphology and venation of the leaf in *Quiina acutangula* Ducke. *Amer. J. Bot.* 37: 159-171.
 HICKEY, L. J. (1973). Classification of the architecture of dicotyledonous leaves. *Amer. J. Bot.* 60: 17-33.
 SCHNEIDER, J. V. (1998). El género *Quiina* (*Quiinaceae*) con especial referencia a las especies de Venezuela. *Acta Bot. Venez.* 21: 1-74.
 SCHNEIDER, J. V., U. SWENSON & G. ZIZKA (2002). Phylogenetic reconstruction of the neotropical family Quiinaceae (Malpighiales) based on morphology and some remarks on the evolution of an androdioecious sex distribution. *Ann. Missouri Bot. Gard.* 89: 64-76.
 STEARN, W. T. (1992). *Botanical Latin: History, Grammar, Syntax, Terminology and Vocabulary*. Davis & Charles, London.

