

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 59 (2004)

Heft: 2

Artikel: Plant diversity in Marmarica (Libya & Egypt) : a catalogue of the vascular plants reported with their biology, distribution, frequency, usage, economic potential, habitat and main ecological features, with an extensive bibliography

Autor: Houérou, Henry Noël le

DOI: <https://doi.org/10.5169/seals-879332>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Plant diversity in Marmarica (Libya & Egypt): a catalogue of the vascular plants reported with their biology, distribution, frequency, usage, economic potential, habitat and main ecological features, with an extensive bibliography

HENRY NOËL LE HOUÉROU

ABSTRACT

LE HOUÉROU, H. N. (2004). Plant diversity in Marmarica (Libya and Egypt): a catalogue of the vascular plants reported with their biology, distribution, frequency, usage, economic potential, habitat and main ecological features, with an extensive bibliography. *Candollea* 59: 259-308. In English, English and French abstracts.

Marmarica, as it is called since antiquity, is the natural region located between the Jebel Lakhdar of Cirenaica and the Nile Delta over an E-W distance of 750 km. It covers an area of 220 000 km² along the southern shore of the Mediterranean Sea between longitudes 23°E and 30°E. It is a typically Mediterranean arid zone in the northern fringe along the shoreline, shifting slowly to absolute desert southwards on a depth of ca. 300 km. Mean annual rainfall is slightly below 200 mm in the northernmost sites dropping to 10 mm at the oases of Jarabub and Siwa 300 km further south. The flora belongs to the Ibero-Maghribian entity of the Mediterranean phytogeographic region with some Saharo-Arabian and Irano-Turanian elements. It includes an overall 1015 vascular taxa representing 48% of the Egyptian flora and 53% of the Libyan flora. Eighteen taxa are endemic to the region. Flora and vegetation are very homogenous and strongly influenced by 3 types of farming: 1) irrigation between Alexandria and El Alamein (ca. 100 km W of the Nile); 2) runoff farming in the valleys along the wadis of the coastal plains with episodic mixed dry farming of cereals; 3) grazing on fallow, stubble and the depleted steppes on the interflaves. Vegetation and flora are strongly influenced by the impact of man: virtually all of the pristine steppe vegetation still present, and mapped in the years 1960's have now been cleared for opportunistic farming. This situation has heavily imprinted on the flora and vegetation now dominated by «weeds». The present study presents a catalogue of the native taxa reported in both the Libyan and Egyptian part of Marmarica with their biology, abundance, frequency of occurrence, geographic distribution, economic and grazing potential, utilization and ecological requirement traits in terms of climate and soils. An extensive bibliography is provided.

RÉSUMÉ

LE HOUÉROU, H. N. (2004). Diversité botanique de la Marmarique (Libye et Egypte): catalogue des plantes vasculaires inventoriées avec leur biologie, répartition, fréquence, usage économique, habitat et caractéristiques écologiques principales, suivi d'une bibliographie étendue. *Candollea* 59: 259-308. En anglais, résumés anglais et français.

La région naturelle de Marmarique, connue sous ce nom depuis l'Antiquité, est située entre le Jebel Lakhdar de Cyrénaïque et le delta du Nil sur une distance O-E de 750 km. Elle couvre une aire de 220 000 km² le long de la côte sud de la Méditerranée entre les longitudes 23°E et 30°E. Elle se présente comme une zone aride typiquement méditerranéenne dans sa partie nord le long de la côte, évoluant progressivement vers un désert absolu sur une distance d'environ 300 km. La pluviosité

moyenne annuelle est légèrement au-dessous de 200 mm dans les sites situés le plus au nord pour descendre progressivement à 10 mm dans les oasis de Jaraboub et de Siwa, situées à 300 km plus au sud. La flore appartient à l'entité phytogéographique ibéro-maghrébiennne de la région botanique méditerranéenne avec quelques éléments saharo-arabiques et irano-touraniens. Elle comprend un total de 1015 taxa vasculaires représentant 48% de la flore égyptienne et 53% de la flore libyenne. Dix-huit taxa sont endémiques de la région. La flore et la végétation sont zonalement très homogènes et fortement influencées par la culture. Les cultures sont de trois types: 1) cultures irriguées d'Alexandrie à El Alamein à environ 100 km à l'O du Nil; 2) cultures arbustives avec utilisation du ruissellement le long des oueds dans la plaine côtière; 3) cultures sèches épisodiques de céréales avec des jachères pâturées sur les interfluves jusqu'à environ 50 km au S de la côte, là où le sol le permet. Les sols squelettiques superficiels à croûte calcaire constituent un pâturage généralement très dégradés. La végétation originelle, encore présente dans les années 1960, a maintenant été défrichée au profit de cultures céréalières opportunistes. Cette situation se répercute sur la flore et la végétation maintenant largement dominées par des «mauvaises herbes». La présente étude est un catalogue des taxa présents en Marmarique tant libyenne qu'égyptienne, avec des indications sur leur biologie, répartition géographique, fréquence, et leurs principales exigences écologiques en matière de climat et de sols, leur utilisation actuelle, leur potentiel économique et pastoral. Une bibliographie étendue est fournie.

KEY-WORDS: Mediterreanean – Arid lands – Land-use – Steppe – Rangeland – Desert – Sahara – Libya – Egypt – Phytogeography – Flora – Vegetation – Ethnobotany

1. Introduction : the geographic setting

The geographic natural region known since antiquity as Marmarica encompasses the land located immediately south of the Mediterranean Sea between longitudes 23° and 30°E, i.e. from 40 km E of Derna and Cyrenaïca's highlands to Alexandria and the Nile Delta. The region extends from the shore line (lat 31° to 32.5°N) to the oases of Jarabub and Siwa (ca. lat 29°N) some 300 km further south in the inland desert (Map 1).

Marmarica is a very homogenous region on many accounts from the stand points of climate, geology, geomorphology, flora and vegetation. Marmarica thus appears as a self contained ecologic and biogeographic unit split in two uneven parts by the political border between Libya and Egypt. The region as a whole covers ca. 220 000 km², 2/3 of which are in Egypt and 1/3 in Libya. A detailed zoning is shown in § 4.2, 'area of distribution'.

From the geomorphic standpoint one may distinguish:

The shoreline generally covered with marine white dunes lying on a upper Pliocene and Holocene more or less consolidated beach of *Strombus bubonius*. The width of the dune zone may reach one km and locally more. Their height may reach 20 m and locally more. Some salt marshes may interrupt the duney offshore lines. The salt marshes are generally tied to the lower reaches of the main wadis. The area covers an area of some 750 km².

The coastal plain, with its valleys and wadi terraces extends from the dune zone to the first escarpment of the inland plateau. It is mainly formed of silty soils, locally moderately saline, sandier to the east, and generally cultivated either to cereals or to tree crops: fig-trees, olive, almond, grapes, date palm, etc. the width of the coastal plain varies from ca. 1 km to ca. 10 km. In the coastal plain also emerge a number (1 to 5, usually 3) ridges parallel to the shoreline made of soft oolitic Tyrrhenian (5000-15000 yrs old) sandstone/limestone, very much in demand as building material and heavily quarried between Borg et Arab and El Alamein. This Tyrrhenian oolitic calcareous sandstone bright white when fresh, turning to an ivory colour when exposed or wheathered, is common along the southern shores of the south eastern Mediterranean from Cap Bon in Tunisia to S Turkey and everywhere popular as building material. The area covered by the coastal plain is of ca. 3750 km².

The inland plateau is made of Mio-Pliocene outcrops of sandy limestone interspersed with sandstones, shales and occasionally unconsolidated interbedded sands (CONANT & GUDARZI, 1964). Some upper Miocene limestones may be fairly saline containing as they do 1-5% NaCl. This is quite unusual as the conditions of sedimentology of CaCO₃ and NaCl are normally not compatible. The salinity of these limestones is thus probably secondary, i.e. post-depositional. These saline limestones, wherever they outcrop directly, are generally covered with a halophytic steppe, often monospecific populations of *Suaeda pruinosa* and sometimes with *Limonium pruinatum* and/or *L. tubiflorum* as co-dominant companions. This is also a very unusual fact of typical perennial halophytic vegetation straight on well drained hard limestone geologic outcrops, with, of course, the exception of Triassic and other diapirs on which they are commonplace, but there are no diapirs in the region.

The inland plateau is generally covered with an upper and mid-Pleistocene calcrete 30-50 cm thick, very hard, salmon coloured with many *Helix* spp. inclusions, hence the often given name of «salmon crust of Helicidae» by the French speaking geologists and soil scientists. This calcrete is often quarried or bulldozed as a building material and/or for road construction.

One may distinguish three levels of the inland plateau raising respectively to ca. 50, 100 and ca. 200 m a.s.l., then softly declining southward (ca. 0.3% slope) to sea level or slightly below towards the oases of Siwa and Jarabub, and to -134 m at the bottom of the Qattara depression. The highest level is reached towards 50 km south of the shoreline. The inland plateau occupies an area of ca. 195 500 km².

The Qattara depression extends to some 25 000 km² in the SE corner of the region north east of Siwa to the Fayoum depression from sea level down to -134 m, and -40 m in the Fayoum. The depression is covered with dry salt marshes and locally with halophytic vegetation (*Arthrocnemum*, *Atriplex*, *Chenolea*, *Halocnemum*, *Halopeplis*, *Imperata*, *Limonium*, *Limoniumstrum*, *Nitraria*, *Phragmites*, *Salicornia*, *Sarcocornia*, *Salsola*, *Seidlitzia*, *Suaeda*, etc.).

The climate is arid Mediterranean on the north to 50 km from the shoreline (REGNER, 1997), hyper-arid Mediterranean from 50 to 100 km southwards and Eremaean-Mediterranean (the scanty rain always occurs in the winter season) further south. Details are shown in the ombrothermal digrammes (LE HOUÉROU, 1989) and in the data and bioclimatic classification (FANTOLI, 1952; MAGGAZINI, 1963; FATHY & MEHANNA, 1980; LE HOUÉROU, 1984; AYYAD & VAN KREULEN, 1987; LE HOUÉROU, 1989; LE HOUÉROU, 1997b). The highest mean annual rainfall is slightly below 200 mm (Alexandria, Dekheila, Ras el Hekma, Ftaiah, Martuba) to the north dropping to 10 mm in Jarabub and Siwa.

2. The sources

The present catalogue results from the following information and data base:

1. Durand and Baratte's *Plantae Libycae Prodromus* (DURAND & BARATTE, 1910) contains a synoptic table with 303 taxa from the region called Turkish Marmarica (i.e. Libyan Marmarica).
2. The *Student's flora of Egypt* by Vivi Täckholm (TÄCKHOLM, 1974) includes 475 references to the flora of Marmarica, to which I added 155 taxa.
3. The *Flora of Egypt. Checklist* by L. Boulos (BOULOS, 1995) contains reference to 960 taxa from Marmarica, to which I added 30 taxa.
4. The *Flora of Libya* (ALI & JAFRI, 1976-1977; JAFRI & EL-GADI, 1977-1989).
5. The *Preliminary check list of the Libyan flora* by H. G. Keith (KEITH, 1965) and *Check list of the Libyan flora* by L. Boulos (BOULOS, 1977, 1979).

6. My book on the bioclimatology and biogeography of the arid steppeland north of the Sahara (LE HOUÉROU, 1995) lists 1178 species and subspecies of vascular plants in the Egyptian and Libyan steppes.
7. Data from the 146 scientific publications on the vegetation of NW Egypt, listed in the bibliography; many of these phytosociological surveys provide useful and detailed information on plant distribution in Egyptian Marmarica; all of them are not quoted in the text, tables and figures, they are nonetheless listed as a potential source of information and possible starting points for further investigations.
8. My own unpublished field data which include 250 site-surveys (“relevés” in the phytosociological jargon) in Egyptian Marmarica from 1988 to 1998 and 300 other relevés in Libyan Marmarica from 1964 to 1966 and from 1981 to 1983, which makes a total of 550 site plant-lists for overall Marmarica.
9. A third check list of the Libyan flora is not yet published.

3. General

The following catalogue identifies 1009 species present in Marmarica from Alexandria to Ras el Tin (40 km E of Derna), on a depth of 300 km from the Mediterranean shore inland. There are, in addition, some 6 subspecies, i.e. a total of 1015 epithets of vascular plants (there are only 2 species of ferns and 3 gymnosperms).

The close concurrence between BOULOS (1995) and LE HOUÉROU (1995), and both of them with the following catalogue ought to be underlined.

As suggested above, the following catalogue identifies 1015 taxa; of these, some:

- 534 taxa (53% of the flora) have virtually no grazing value.
- 476 (47% of the flora) are weeds of unknown utilization to me.
- 389 (38% of the flora) have a mediocre grazing value (scoring 1 to 3 on the evaluation scale).
- 96 (10% of the flora) constitute good grazing taxa (scoring 3 to 5 on the evaluation scale).
- 12 (1% of the flora) are excellent fodder taxa, i.e. actual or potential fodder crops, including some shrubs.
- 47 (5% of the flora) are toxic to stock (see § 4.2, ‘grazing value’).
- 98 (10% of the flora) are shrubs and trees (see § 4.2, ‘biological type’).
- 118 (12% of the flora) are halophytes.
- 177 (17% of the flora) are psammophytes.
- 104 (10% of the flora) are hygrophytes.
- 33 (3% of the flora) are hydrophytes.
- 56 (6% of the flora) are nitratophytes.
- 55 (5% of the flora) are chasmophytes.
- 18 (2% of the flora) are parasites or hemi-parasites.
- 138 (14% of the flora) have an actual or potential medicinal usage; for details on these medicinal usages (see BOULOS, 1983).

The systematic order, taxonomy and synonymy utilized follow BOULOS (1995) with few exceptions e.g. *Rhamnus lycioides* subsp. *oleoides* in Boulos is *R. oleoides* here, since, rightly or

wrongly, opposite to Boulos, I consider this taxon as quite a valid species on its own right. Some species have been added (e.g. *Helianthemum kahiricum* and *Periploca angustifolia*) and others removed (e.g. *Medicago sativa*) from the Boulos checklist. Some 12 species are recorded herein for the first time in Marmarica, e.g. *Acacia ehrenbergiana* «Seyal», recently reported by a bedouin from Wadi Iskanderia, ca. 150 km NE of Siwa (100 km E of Bir Nousf) (G. S. Mikhiel, *pers. comm.*). I should specify that I did not see this tree myself. I am only reporting information collected by someone else, which seemed plausible to me, from the description of the tree, of its fruit, and from the vernacular name given to it by nomads. This information requires confirmation, this is why I report it herein.

4. Attributes Coding

4.1 General

Table 3 shows the rating of the 92 parameters mentioned below, in addition to the scientific epithets and vernacular names. They are distributed in 12 groups in Table 3.

1. & 2. Scientific name (genus, species & subspecies).
3. Botanic family.
4. Local vernacular names are, to a large part, taken in TÄCKHOLM (1974), BOULOS (1983) and LE FLOC'H (1983), and from BOULOS & EL HADIDI (1984), with some additions and corrections from my own field experience, transliteration and transcription follow the mentioned sources.
5. Frequency status.
6. Area of distribution in 5 parallel belts from the Mediterranean shore to some 150 km inland, i.e. from a mean annual rainfall (MAR) belt of ca. 150 mm to less than 25 mm, then a sixth zone between 150 and 300 km from the shoreline, corresponding to the Eremaean zone or true desert with 25 to 10 mm of MAR.
7. Biological type.
8. Life span.
9. Habitat.
10. Utilization.
11. Grazing value on an evaluation scale.
12. Biological, ecological and other miscellaneous remarks.

4.2 Codes

Below are presented explanations of codes of Table 3 for 'frequency', 'area of distribution', 'biological type', 'life span', 'habitat', 'utilization', 'grazing value' and 'remarks' respectively.

Frequency status

- CC = Very common
- C = Common
- FC = Fairly common
- FR = Fairly rare
- R = Rare
- RR = Very rare

This rating results from TÄCKHOLM (1974), from the many phytosociological studies published and mentioned in the bibliography and also from my own field notes.

Area of distribution

- I = 0-2 km from Mediterranean shoreline: ca. 750 km²
- II = 1-10 km inland steppe: ca. 3750 km²
- III = 10-25 km inland steppe: ca. 9375 km²
- IV = 25-50 km inland desert: ca. 18750 km²
- V = 50-150 km inland hyper-arid zone: ca. 75 000 km²
- VI = 150-300 km inland Eremaean zone: ca. 112 500 km² (including the Qattara depression: 25 000 km²)

The total area is ca. 220 125 km².

Biological type

- C = Crassulescent (= fleshy halophyte)
- CH = Crassulescent (= fleshy herb)
- Cl = Climber
- F = Fern
- Fr = Frutescent (= herbaceous with a woody base)
- H = Herbaceous (forb & grass)
- Par = Parasite & hemi-parasite
- S = Shrub
- CS = Crassulescent shrub (= fleshy halophyte)
- DS = Dwarf shrub (h < 50 cm)
- HS = Tall shrub (h > 50 cm)
- TS = Trailing shrub
- Su = Succulent (= fleshy glycophyte)
- T = Tree (h > 3 m)

Life span

- A = Annual (= therophyte)
- B = Biennial
- P = Perennial
- Sh = Short-lived perennial (= pauciennial)

Habitat

- Bra = Brackish water
- Can = Canals
- Cli = Cliffs
- Coa = Coastal (= littoral)
- Con = Continental
- Cul = Cultigene, rainfed crops and fallows
- Des = Desert (MAR < 100 mm)
- Dne = Dunes
- Frw = Fresh (sweet) water
- Irr = Irrigated land
- Rge = Rangeland
- Roc = Rocks
- Run = Runnels

Sea = Open sea water
 Sha = Shallow soils
 Snd = Sandy soils
 Slt = Salt marshes
 Ste = Steppe
 Wad = Wadis & topographic depression
 Wel = Wells
 Wet = Wetland
 Wld = Wasteland

Utilization

Cro = Crop
 Fen = Fencing (live)
 Fsh = Fodder shrub
 Fue = Fuel wood
 Gra = Grazing
 Han = Handicraft
 Med = Medicinal & herbal
 Non = None
 Orn = Ornamental
 Sbi = Sand binding
 Unk = Unknown
 Veg = Vegetable, salad, fruit, human food
 Wbr = Windbreak

Grazing value

T = Toxic
 0 = No value
 1 = Poor
 2 = Mediocre
 3 = Fair
 4 = Good
 5 = Excellent (fodder crops, incl. shrubs)

Remarks

Cha = Chasmophyte (craks, crevices, diaclases)
 End = Endemic (restricted to Marmarica)
 Ere = Eremophyte (= desert plant)
 Fce = Fencing material
 Fod = Fodder (actual or potential)
 FoRe = Forest relict
 Gly = Glycophyte (= non tolerant to salinity in generally halophytic groups)
 Hal = Halophyte (salt - tolerant)
 Hyd = Hydrophyte (living in free water)
 Hyg = Hygrophyte (living in wet soils)
 Nat = Naturalized xenophyte
 Nit = Nitratophyte (on nitrates-rich soils)
 Pel = Pelophyte (on clay soils)

- Phr = Phreatophyte (on water table)
 Psa = Psammophyte (on sandy soils)
 Sil = Limonophyte (on silty/loamy soils)
 SO₄ = Gypsophyte (on gypsic soils)
 Sum = Summer growing
 Tac = Tachytherophytes (very short lived annuals: Aacheb – Ghizzu)
 Wee = Weed (in range or crop)

4.3 Interpretation of the Plant Biodiversity Table (Table 3)

4.3.1 Analysis of the flora

Table 3 contains:

86 families of vascular plants, out of a total of 121 for the overall flora of Egypt (71%) and 168 (51%) for Libya.

451 genera out of an overall 742 (61%) for Egypt and 845 (53%) for Libya.

1015 species and subspecies which contain 777 dicots, 233 monocots, 3 gymnosperms and 2 ferns.

The overall vascular flora of Egypt includes 2121 taxa; the flora of Marmarica thus represents nearly half the Egyptian flora in terms of taxa numbers (47.9%). The flora of Libya includes 1930 taxa (2200 with the xenophytes); Marmarica as a whole thus represents the equivalent of some 53% of the country's flora. But this is not a realistic account since the flora of Libyan Marmarica does not exceed 400 taxa and thus the Libyan Marmarica flora is only 21% (= 400/1930) of the national vascular flora.

The largest families are as follows for both Marmarica and Egypt:

Table 1. – Family Richness

Families	Libya		Marmarica		Egypt	
	Nr of spp.	%	Nr of spp.	%	Nr of spp.	%
<i>Compositae</i>	277	14.4	128	12.6	230	10.8
<i>Gramineae</i>	238	12.3	130	12.8	277	13.1
<i>Leguminosae</i>	210	10.9	104	10.2	233	11.0
<i>Cruciferae</i>	116	6.0	62	6.1	102	4.8
<i>Chenopodiaceae</i>	64	3.3	51	5.0	77	3.6
<i>Boraginaceae</i>	53	2.7	32	3.2	55	2.6
<i>Caryophyllaceae</i>	66	3.4	41	4.0	85	4.0
<i>Umbelliferae</i>	85	4.4	30	3.0	51	2.4
<i>Cyperaceae</i>	28	1.5	25	2.5	44	2.1
<i>Convolvulaceae</i>	18	0.9	22	2.2	48	2.3
<i>Labiatae</i>	64	3.3	16	1.6	55	2.6
<i>Scrophulariaceae</i>	38	2.0	13	1.3	62	2.9
Total	1257	65.1	654	64.4	1319	62.2

The 27 larger genera in taxa numbers are shown in Table 2.

Table 2. – Number of taxa in the major genera

Genera	Nr of spp.		
	Libya	Marmarica	Egypt
<i>Euphorbia</i>	19	19	42
<i>Medicago</i>	21	16	18
<i>Plantago</i>	15	14	18
<i>Allium</i>	9	13	22
<i>Silene</i>	19	13	24
<i>Trifolium</i>	22	13	19
<i>Erodium</i>	16	12	14
<i>Convolvulus</i>	16	12	23
<i>Astragalus</i>	29	11	32
<i>Atriplex</i>	14	11	15
<i>Centaurea</i>	8	10	16
<i>Cyperus</i>	6	10	23
<i>Helianthemum</i>	19	10	23
<i>Lotus</i>	17	10	12
<i>Salsola</i>	12	10	20
<i>Reseda</i>	7	9	14
<i>Trigonella</i>	8	9	11
<i>Amaranthus</i>	4	8	11
<i>Bromus</i>	9	8	11
<i>Fumaria</i>	8	8	16
<i>Persicaria</i>	1	8	8
<i>Bellevalia</i>	1	7	8
<i>Echium</i>	13	7	12
<i>Limonium</i>	13	7	7
<i>Scirpus</i>	1	7	7
<i>Suaeda</i>	7	7	9
<i>Vicia</i>	11	7	9
Total n	325	276	444
% of flora	17	27	21

The above tables shows that the flora of Marmarica is, by and large, similar to that of the two countries as a whole in its proportions.

It is worth noting that genera having many taxa of hygrophytes or of halophytes are particularly well represented in Egyptian Marmarica:

Hygrophytes	Halophytes
<i>Cyperus</i>	<i>Atriplex</i>
<i>Persicaria</i>	<i>Limonium</i>
<i>Scirpus</i>	<i>Salsola</i>
<i>Trigonella</i>	<i>Suaeda</i>

The same remark applies to genera with many weeds, nitratophytes and wasteland taxa:

Hygrophytes	Halophytes
<i>Allium</i>	<i>Euphorbia</i>
<i>Amaranthus</i>	<i>Fumaria</i>
<i>Bellevalia</i>	<i>Medicago</i>
<i>Bromus</i>	<i>Reseda</i>
<i>Centaurea</i>	<i>Silene</i>
<i>Echium</i>	

In terms of endemics, BOULOS (1995) recognizes 61 endemic taxa in Egypt of which over half are from Sinai. Marmarica is poorer in this respect since we only record 19 species and subspecies:

<i>Allium blomfeldianum</i> E	<i>Lepidium aucheri</i> E & L
<i>Allium desertorum</i> E & L	<i>Nigella arvensis</i> subsp. <i>taubertii</i> E & L
<i>Allium mareoticum</i> E & L	<i>Pancratium arabicum</i> E & L
<i>Allium myrianthum</i> E	<i>Silene fruticosa</i> E & L
<i>Carduncellus mareoticus</i> E & L	<i>Silene biappendiculata</i> L
<i>Echinops taeckholmianus</i> E	<i>Teucrium brevifolium</i> E & L
<i>Ebenus armitagei</i> E & L	<i>Verbascum letourneuxii</i> E & L
<i>Ferula marmarica</i> E & L	<i>Viola scorpiuroides</i> E & L
<i>Helianthemum sphaerocalyx</i> E	<i>Zilla spinosa</i> subsp. <i>biparmata</i> E & L
<i>Herniaria cyrenaica</i> L	

E = Egyptian endemics; **L** = Libyan endemics.

As one can see most endemics are shared between the two sub-regions.

Microendemism (varieties are not considered in Table 3) is represented by the following:

- *Centaurea conglomerata* var. *claviceps*, *Pituranthos tortuosus* var. *virgatus* (= var. *arcuatus* Le Houérou).
- *Vicia calcarata* var. *marmarica*.

4.3.2 Floristic similarities and differences between Libyan and Egyptian Marmarica

There are some 24 taxa from the Libyan side of Marmarica that have not been reported from the Egyptian part:

Juniperus phoenicea, *Paretaria lusitanica*, *Blitum virgatum*, *Atriplex mollis*, *Diploaxis simplex*, *Sinapis pubescens*, *Euphorbia bivonae*, *Olea europaea* subsp. *oleaster*, *Phyllirea media*, *Pergularia tomentosa*, *Calotropis procera*, *Convolvulus supinus*, *C. tricolor*, *Cuscuta europaea*, *C. epithymum*, *Scaligera cretica*, *Borago officinalis*, *Ballota pseudo-dictamnus*, *Rhanterium suaveolens*, *Asparagus stipularis*, *Castellia tuberculata*, *Melica minuta*, *Elytrigia cristata*, *E. juncea*.

These include a number of taxa linked to the Libyan semi-arid vegetation of Cyrenaica which extends into nearby eastern Marmarica, for reason of vicinity. Conversely there are some 625 taxa present in Egyptian Marmarica which are lacking on the Libyan side of the political border.

These are mostly hydrophytes, hygrophytes, helophytes and wetland taxa tied to canals, ponds, irrigation and agriculture such as, but not limited to, the following 21 genera and 68 species:

Persicaria, 8 species, *Ranunculus*, 4 species, *Lythrum*, 3 species, *Ammania*, 4 species, *Centaurium*, 3 species, *Damasonium*, 1 species, *Najas*, 3 species, *Potamogeton*, 4 species, *Ruppia*, 2 species, *Juncus*, 6 species, *Leersia*, 1 species, *Spirodela*, 1 species, *Lemna*, 1 species, *Pseudowolfia*, 1 species, *Typha*, 1 species, *Carex*, 2 species, *Schoenus*, 1 species, *Fimbristylis*, 2 species, *Scirpus*, 7 species, *Eleocharis*, 3 species, *Cyperus*, 10 species.

4.3.3 Frequency status (see § 4.2 for abbreviations)

Frequency status	Nr of spp.	Subtotal	%	Subtotal
CC	204	489	20.1	48.2
C	168		16.6	
FC	117		11.5	
FR	81	525	8.0	51.8
R	269		26.5	
RR	175		17.3	
Total	1014	1014	100	100

The above table shows that rare taxa are slightly more numerous than the common ones, a fact that could be interpreted as one more evidence of the intensive anthropization of the Marmarica vegetation and flora.

4.3.4 Area of distribution (see § 4.2 for abbreviations)

Area of distribution	Nr of spp.	%	
I	33	3.3	Shoreline
I-II	683	67.3	Shore and coastal plain steppe
I-III	186	18.3	Shore and steppe
I-IV	22	2.2	Shore, steppe and desert
I-VI	10	1.0	Shore, steppe, desert and Eremian zone
II-III	10	1.0	Steppe
II-IV	21	2.1	Steppe and desert
II-VI	4	0.4	Steppe, desert & Eremian
III-IV	9	0.9	Desert and Eremian
III-VI	37	3.6	Desert and Eremian
Total	1015	100	

The above table shows that the shore and steppe zones include some 90% of the taxa while less than 10% are restricted to the desert. The shore and steppe zones include the newly irrigated land west of the Nile Delta (El Amariya, El Nobariya, El Hammam, Borg el Arab, etc.) which significantly contributes to this richness.

This zone includes some 148 taxa tied to irrigation and wetlands and free water areas (canals, ponds).

4.3.5 *Biotypes (see § 4.2 for abbreviations)*

Biological types	Nr of spp.	%
CH	26	2.5
Cl	11	1.1
F	2	0.2
Fr	50	4.8
H	816	79.0
Par	18	1.7
S	3	0.3
CS	25	2.4
DS	20	1.9
HS	37	3.6
TS	10	1.0
Su	6	0.6
T	9	0.9
Total	1033	100

The total number of taxa is superior to 1015, the codes being not mutually exclusive.

Biotypes are largely dominated by herbaceous taxa (CH, F and H = 82% of the flora), i.e. 844 taxa. Woody or partially woody taxa (T, S and Fr) represent only 6% of the flora, i.e. 62 taxa. The number of succulent taxa (fleshy glycophytes) is negligible (6). The same remark applies to climbers (11). Crassulescent shrubs (CS) and crassulescent herbs (CH) are halophytes; they number 51; half of them are *Chenopodiaceae*.

4.3.6 *Life span (see § 4.2 for abbreviations)*

Life span	Nr of spp.	%
A	575	54.9
B	27	2.6
P	433	41.3
Sh	13	1.2
Total	1048	100

The total number of taxa is superior to 1015, the codes being not mutually exclusive.

Annual taxa constitute 55% of the flora; most are herbaceous. Biennials and short lived perennials (pauciennials) represent an insignificant proportion (4%).

4.3.7 Habitat (see § 4.2 for abbreviations)

Habitat	Nr of spp.	%
Bra	3	0.2
Can	24	1.3
Cli	38	2.1
Coa	31	1.7
Con	4	0.2
Cul	453	25.5
Des	36	2.0
Dne	12	0.7
Frw	24	1.3
Irr	32	1.8
Rge	288	16.2
Roc	32	1.8
Run	24	1.4
Sea	5	0.3
Sha	33	1.9
Snd	75	4.2
Slt	64	3.6
Ste	145	8.2
Wad	47	2.6
Wel	4	0.2
Wet	96	5.4
Wld	308	17.3
Total	1778	100

The total number of taxa is superior to 1015, the codes being not mutually exclusive.

The above table shows that the area under study is a very disturbed one, subjected to heavy anthropogenic pressure, since the field and fallow taxa (Cul) represent 26% of the flora while those from wastelands (Wld and Wet) are 404 i.e. another 23%. The total of taxa from disturbed land thus represents 49% of the flora. This is further confirmed by the analysis of the features related to ecology and utilization as shown below. The taxa related to water and wet soils (Bra, Can, Frw, Irr & Wet) are 179, i.e. 10% of the flora.

4.3.8 Utilization (see § 4.2 for abbreviations)

Features	Nr of spp.	%
Cro	6	0.5
Fen	6	0.5
Fsh	10	0.9
Fue	31	2.7
Gra	410	36.3
Han	8	0.7
Med	138	12.2
Non	473	41.8
Orn	18	1.6
Sbi	5	0.4
Unk	5	0.4
Veg	17	1.5
Wbr	4	0.4
Total	1131	100

The total number of taxa is superior to 1015, the codes being not mutually exclusive.

A total of 478 taxa (42% of the flora) have no use or an unknown utilization; this confirms the data from subsection 4.3.7 on habitat; a substantial number of taxa, however, is utilized for grazing (410, i.e. 36% of the flora). A third trait has a significant number: the medicinal and herbal taxa (138, i.e. 12% of the flora).

4.3.9 Grazing value (see § 4.2 for abbreviations)

Parameter	Nr of spp.	Subtotal	%	Subtotal
T	47	581	4.6	57.3
0	534		52.7	
0-1	2	337	0.2	33.2
1	126		12.4	
1-2	44		4.3	
2	132		13.0	
2-3	33		3.3	
3	54	96	5.3	9.5
3-4	13		1.3	
4	13		1.3	
4-5	4		0.4	
5	12		1.2	
Total	1014		100	100

The above table shows that 96 taxa in the flora (9.5%) have a fair to good grazing value. Unfortunately most of those have become rare and none of them is dominant. It shows, however, that the flora has still, in principle, a good potential for range regeneration, if the land

were appropriately managed. The depletion is not yet beyond repair. The number of taxa with a low grazing value is 337 (33%) while those of no grazing value or toxic (poisonous) to stock are 581 (57%); this confirms, once more, the previous diagnosis: heavy anthropozoic pressure on disturbed and depleted land and vegetation.

4.3.10 Remarks and edaphic biological and ecological classification (see § 4.2 for abbreviations)

Traits	Nr of spp.	%
Cha	55	4.4
End	18	1.4
Ere	42	3.4
Fod	14	1.1
Gly	13	1.0
Hal	118	9.5
Hyd	33	2.6
Hyg	104	8.3
Nat	10	0.8
Nit	56	4.5
Pel	24	1.9
Phr	16	1.3
Psa	177	14.2
Sil	46	3.7
SO ₄	23	1.8
Sum	21	1.7
Tac	39	3.1
Wee	439	35.2
Total	1248	100

The total number of taxa is superior to 1015, the codes being not mutually exclusive.

Again, the taxa from disturbed land are in large numbers (Wee + Nit = 495 i.e. 40% of the flora).

Psammophytes are in sizeable numbers (14% of the flora) and often make up the best grazing; hygrophytes and hydrophytes make up 11% of the flora and halophytes 10%.

Chasmophytes are not in large numbers but they have a important role as cliffs and rocks constitute a refuge where species that have become rare can find shelter and relative protection due to a relative site inaccessibility to man and stock. Many cliffs, runnels and wadis constitute sites of high biodiversity, where 50 to 100 ± rare species can be found over an area of 1000 m². Among those one may include:

<i>Allium mareoticum</i>	<i>Micromeria nervosa</i>
<i>Anabasis oropediorum</i>	<i>Moricandia nitens</i>
<i>Artemisia herba-alba</i>	<i>Oryzopsis miliacea</i>
<i>Capparis rupestris</i>	<i>Periploca angustifolia</i>
<i>Convolvulus oleifolius</i>	<i>Phagnalon rupestre</i>
<i>Dactylis hispanica</i>	<i>Phlomis fruticosa</i>
<i>Dichanthium annulatum</i>	<i>Polygonum equisetiforme</i>
<i>Echiochilon fruticosum</i>	<i>Prasium majus</i>
<i>Euphorbia dendroides</i>	<i>Rhamnus oleoides</i>
<i>Euphorbia bivonae</i>	<i>Rhus tripartita</i>
<i>Euphorbia hierosolymitana</i>	<i>Silene fruticosa</i>
<i>Fagonia cretica</i>	<i>Silene vulgaris</i>
<i>Globularia arabica</i>	<i>Thymus capitatus</i>
<i>Helianthemum sessiliflorum</i>	<i>Varthemia candicans</i>
<i>Helianthemum sphaerocalyx</i>	<i>Verbascum letourneuxii</i>
<i>Hyparrhenia hirta</i>	<i>Ziziphus lotus</i>
<i>Lotus creticus</i>	

As mentioned above the large number of hydrophytes and hygrophytes reflects the presence in the eastern part of Marmarica of large irrigated zones, west of Alexandria.

Biodiversity has been herein evaluated on a 6 degrees scale according to the number of species recorded, in spring-time, over a standard sampling area of 2500 m² (1/4 hectare or 0.6 feddan):

1: 0-10 - Depleted / Desertized	4: 40-60 - Medium
2: 10-20 - Poor / Low	5: 60-80 - Good
3: 20-40 - Fair	6: > 80 - Outstanding

5. Conclusions

Marmarica is a botanically rich province, considering its climatic aridity, since it harbours almost 50% of the Egyptian flora and 21% of the Libyan flora, similar in that to Sinaï (same size: 960 taxa). The 1015 Marmarican taxa represent the equivalent of 37% of the Saharan flora on only 2.6% of the surface area. Sinaï, however, due to its high mountains, has a higher rate of biodiversity, with 31 endemic taxa vs 18 for Marmarica. The same rationale applies to the comparison between Libyan, Marmarica and Cyrenaïca. The latter, due to its semi-arid and locally sub-humid bioclimate, its forest vegetation and its geographic isolation is an «island of endemism» with some 100 endemic taxa and many micro-endemics, for an overall flora of ca. 900 taxa, i.e. 11% vs less than 3% for Egypt, ca. 1.8% for Marmarica as a whole, 4% for the Libyan part and 1.8% for the Egyptian part.

The Marmarica flora belongs to the Ibero-Maghribian sub-province of the Mediterraneo-steppic province within the Mediterranean phytogeographic region; some Irano-Turanian and Saharo-Arabian elements are also present due to the vicinity of these two phytogeographic entities but they represent less than 10% of the flora each, while the tropical element is almost absent. But Egyptian Marmarica, likewise the Nile valley and delta, harbours a large number of cosmopolitan and pluriregional taxa tied to irrigation. Conversely Libyan Marmarica where irrigation is scant does not possess this cosmopolitan flora. These facts were carefully analysed in detail in my opus of 1995 on the bioclimatology and phytogeography of the arid steppes N of the Sahara and its more recent synthesis in English language (LE HOUÉROU, 2001).

An important characteristic of the Marmarica flora and vegetation is their high degree of anthropization. Most pristine steppes (e.g. of *Artemisia herba alba*) are reduced to a few scattered plots of a few feddans (1 feddan = 1 acre = 4200 m²). The only one of any size I could find had about 20 ha at ca. 12 km E of Sidi Barrani along the highway to Marsa-Matruh, it was only made possible by the initiative of a «conservative» settled farmer. According to the FAO (1970) range map «Shih» steppes still covered some 220000 hectares W of El Alamein in the late 1960's mostly between El Alamein and Fuka (187 500 ha) with two nuclei S and E of Sidi Barrani (12 500 ha) and SW of Negheila (22 500 ha) (FAO, 1970; VAN DER VEEN & al., 1968).

Species that are sought for by man and livestock find (temporary?) refuge in cliffs, runnels, wadi cañons and rocks. If some action is to be taken for the restoration of biodiversity it should be a total protection (exclosure) of these refuges.

Many other conclusions could be drawn from Table 3 through the systematic study of correlations between a number of parameters such habitat and utilization, grazing value versus families and genera, etc. I leave this studies to the potential users of the present work. I have proposed a concrete programme of conservation to protect the relict species, as a potential source of drought-tolerant and potentially economic germplasm material for arid-land revegetation (LE HOUÉROU, 1998e). As feared, nothing happened after four years, in spite of the fact that the resources were available, from a IBRD loan, to carry out such a programme.

ACKNOWLEDGEMENTS

The author is indebted to Prof. Loutfy Boulos, Cairo, Dr. Edouard Le Floc'h, Montpellier, and three anonymous reviewers for pertinent comments on the draft. He is also thankful to Mr Pascal Maier and Ms Anne-Marie Rosier from the library of Montpellier II University for their useful contribution to setting up some of the bibliographic references.

REFERENCES

The present bibliography is meant to cover the various aspects of vegetation and rangeland sciences in Egyptian and Libyan Marmarica. It includes some 170 references; it is believed to be fairly exhaustive, covering both formal and informal scientific publications, including the so-called «grey literature». Not all of them are mentioned in the text, tables or figures.

- ABDEL-KADER, F. (1980). *Soil erosion*. REMDENE n° 1, 4: 4/1-4/27. University of Alexandria.
- ABDEL-KADER, F. (1981). *Semi-detailed soil survey, El Omayed pilot area*. REMDENE n° 2, 3(3): 8-33. University of Alexandria.
- ABDEL-RAHMAN, A. A., M. A. AYYAD & M. M. EL-MONAYARI (1966). Hydroecology of the sand dune habitat at Burg-el-Arab. *Bull. Fac. Sci. Cairo Univ.* 40: 29-54.
- ADEL-RAHMAN, A. A. & M. A. HAMMOUDA (1960). Vegetation development in fenced areas at Ras-el-Hekma in four years. *U.A.R. J. Bot.* 3: 1-11.
- ABDEL-RAZIK, M. (1976). *A study on the vegetation composition, productivity and phenology in a Mediterranean desert ecosystem at Omayed (Egypt)*. MSc thesis. University of Alexandria. 106 p.
- ABDEL-RAZIK, M., M. ABDEL-AZIZ & M. A. AYYAD (1984). Multivariate analysis of vegetational variation in different habitats of Omayed (Egypt). *Vegetatio* 57: 167-175.
- ABDEL-RAZIK, M., M. A. AYYAD & S. HENEIDY (1988a). Phytomass and mineral composition in range biomass of a Mediterranean arid ecosystem (Egypt). *Acta Oecol., Oecol. Pl.* 9: 359-370.
- ABDEL-RAZIK, M., M. A. AYYAD & S. HENEIDY (1988b). Preference of grazing mammals for forage species and their nutritive value in a Mediterranean desert ecosystem (Egypt). *J. Arid Environm.* 15: 297-305.
- AHMED, A. M. & M. M. MOUNIR (1982). *Regional studies on the natural resources of the NW coastal zone, Egypt*. US National Science Foundation. Oklahoma State University and the UNEP Remote Sensing Centre. Academy of Scientific Research and Technology, Cairo. 195 p.
- ALI, S. I. & S. M. H. JAFRI (eds.) (1976-1977). *Flora of Libya*. Al Faateh University, Tripoli.

- ASLAN, M. H. (1959). *Adaptability and palatability of some forage plants in the western desert of Egypt*. MSc thesis. University of Cairo.
- ATTA, B. A. M. (1953). *Ecological and sociological studies on the plant communities of Mariut*. MSc thesis. University of Alexandria. 130 p.
- AYYAD, M. A. (1957). *An ecological study of Ras el Hikma*. MSc thesis. University of Cairo. 281 p.
- AYYAD, M. A. (1971a). An edaphic study of habitats at Ras-el-Hekma. *Desert Inst. Bull.* 19: 245-259.
- AYYAD, M. A. (1971b). A study of solar radiation on sloping surfaces at Alexandria. *U.A.R. J. Bot.* 14: 15: 65-73.
- AYYAD, M. A. (1973). Vegetation and environment of the western Mediterranean coastal land of Egypt. I The habitat of sand dunes. *J. Ecol.* 61: 509-523.
- AYYAD, M. A. (1976). Vegetation and environment of the western Mediterranean coastal land of Egypt. IV The habitat of non-saline depressions. *J. Ecol.* 64: 713-722.
- AYYAD, M. A. (1978). A preliminary assessment of the effect of protection on the vegetation of the Mediterranean desert ecosystem. *Taeckholmia* 9: 85-101.
- AYYAD, M. A. (1979). *Phytomass, phenology and necromass*. SAMDENE n° 5, 1: 1/1-1/60. University of Alexandria.
- AYYAD, M. A. (1981). Soil-vegetation-atmosphere interactions. In: GOODALL, D. W. & R. M. PERRY (eds.), *Arid Land Ecosystems: Structure, Functioning and Management* 2: 9-31. Cambridge University Press.
- AYYAD, M. A. (1983a). Some aspects of land transformation in the western Mediterranean desert of Egypt. *Advances Space Res.* 2(8): 19-29.
- AYYAD, M. A., M. ABDEL-RAZIK & S. EL-DARIER (1985). Variations in nutrient contents in olive agroecosystems in the western desert of Egypt. *Proc. Egypt. Bot. Soc.* 4: 1277-1297.
- AYYAD, M. A., M. ABDEL-RAZIK & N. GHALI (1983). On the phenology of desert species of western Mediterranean coastal region of Egypt. *Int. J. Ecol. Environm. Sci.* 9: 169-183.
- AYYAD, M. A. & M. Y. AMMAR (1973). Relationship between local physiographic variations and the distribution of common Mediterranean desert species. *Vegetatio* 27: 163-176.
- AYYAD, M. A. & M. Y. AMMAR (1974). Vegetation and environment of the western Mediterranean coastal land of Egypt. II The habitat of inland ridges. *J. Ecol.* 62: 439-456.
- AYYAD, M. A. & M. A. EL BAYYOUMI (1979). On the phytosociology of sand dunes of the western desert of Egypt. In: SINGH, J. S. & B. GOPAL (eds.), *Glimpses of Ecology*: 219-237. International Scientific Publications, Jaipur.
- AYYAD, M. A. & R. M. EL-GHAREEB (1972). Microvariations in edaphic factors and species distribution in a Mediterranean salt desert. *Oikos* 23: 125-131.
- AYYAD, M. A. & R. M. EL-GHAREEB (1973). Correlation of the distribution of common halophilous species at Burg-el-Arab (Egypt) with salinity and level of the water table. *Egypt. J. Bot.* 16: 379-386.
- AYYAD, M. A. & R. M. EL-GHAREEB (1982). Salt marshes vegetation of the western Mediterranean desert of Egypt. *Vegetatio* 49: 3-19.
- AYYAD, M. A. & A. A. EL-GHONEIMY (1976). Phytosociological and environmental gradients in a sector of the western desert of Egypt. *Vegetatio* 31: 93-102.
- AYYAD, M. A. & H. F. EL-KADI (1981). *Consumption and behaviour of grazing animals*. REMDENE n° 2, 1: 2/1-2/31. University of Alexandria.
- AYYAD, M. A. & H. F. EL-KADI (1982). Effect of protection and controlled grazing on the vegetation of a Mediterranean desert ecosystem in northern Egypt. *Vegetatio* 49: 129-139.
- AYYAD, M. A. & S. I. GHABBOUR (1977). Systems analysis of Mediterranean desert ecosystems of northern Egypt. *Environm. Conservation* 4: 91-101.
- AYYAD, M. A. & S. H. HILMY (1974). The distribution of *Asphodelus microcarpus* and associated species on the western Mediterranean coast of Egypt. *Ecology* 55: 511-524.
- AYYAD, M. A. & S. KAMAL (1980). *Analysis of vegetation*. REMDENE n° 1, 1: 3/1-3/30. University of Alexandria.
- AYYAD, M. A. & S. KAMAL (1988). Ground surveys and remote sensing of vegetation and land-use in Mariut (Egypt). In: EL-RAEY, M. (ed.), *Symposium of Environmental Sciences*: 134-154. University of Alexandria.
- AYYAD, M.A. & E. LE FLOCH (eds) (1983). *An ecological assessment of renewable resources for agricultural development in western Mediterranean coastal region of Egypt*. CNRS, ORSTOM, REMDENE. Montpellier, Paris, Alexandria. 104 p.

- AYYAD, M.A., K. H. SHALTOUT, N. N. GHALY, H. F. EL-KADI & S. A. OMAR (1979). *Phytomass, phenology and necromass*. SAMDENE n° 5, 1: 1/1-1/8. University of Alexandria.
- AYYAD, M. A. & H. VAN KEULEN (eds) (1987). *The 'Mariut' project. Final report submitted to Directorate General for International Cooperation (DGIS)*, part 2: 61-92, part 3. CABO-report, Wageningen. 150 p.
- BATANOUNY, K. H. (1968). On the autoecology of *Pithyranthos tortuosus* (Desf.) Benth. & Hook. *Bull. Fac. Sci. Cairo Univ.* 42: 35-45.
- BATANOUNY, K. H. & S. ABU EL-SOUOD (1972). Ecological and phytosociological study of a sector in the Libyan desert. *Vegetatio* 25: 335-356.
- BATANOUNY, K. H. & M. A. F. ZAKI (1969). Root development of two common species in different habitats in the Mediterranean subregion in Egypt. *Acta Bot. Acad. Sci. Hung.* 15: 217-226.
- BATANOUNY, K. H. & M. A. F. ZAKI (1973). Range potentialities of a sector in the Mediterranean coastal region in Egypt. *Vegetatio* 27: 115-130.
- BOULOS, L. (1977). A check list of the Libyan flora. 1: Introduction and Adiantaceae-Orchidaceae. *Publ. Cairo Univ. Herb.* 7-8: 115-141.
- BOULOS, L. (1979). A check list of the Libyan flora. 2: Salicaceae-Neuradaceae. *Candollea* 34: 21-48.
- BOULOS, L. (1983). *Medicinal Plants of North Africa*. Référence Publications, Inc.
- BOULOS, L. (1995). *Flora of Egypt. Checklist*. Al Hadara Publishing.
- BOULOS, L. (1999). *Flora of Egypt* 1. Al Hadara Publishing.
- BOULOS, L. (2000). *Flora of Egypt* 2. Al Hadara Publishing.
- BOULOS, L. & M. EL HADIDI (1984). *The Weed Flora of Egypt*. American University in Cairo Press.
- BOULOS, L. & C. JEFFREY (1979). A check list of the Libyan flora. 3: Compositae. *Candollea*: 34: 307-332
- CONANT, L. C. & G. H. GOUDARZI (1964). *Geologic map of the Kingdom of Libya*. Map I – 350 A, Scale 1: 2,000,000. Miscellaneous Geologic Investigations US Geological Survey, Department of the Interior.
- DRAZ, O. (1954). Some desert plants and their uses in animal feeding (*Kochia indica* and *Prosopis juliflora*). *Publ. Inst. Désert Egypte* 2.
- DURAND, E. & G. BARRATTE (1910). *Florae Libycae Prodromus*. Genève.
- EL-BAYYOUMY, M. A. (1976). *Ecological Phytosociological studies on the sand dune vegetation at Mariut area*. MSc thesis. University of Tanta. 70 p.
- EL-GHAREEB, R. M. (1975). *A study of the vegetation-environmental complex of saline and marshy habitats on the North Western coast of Egypt*. PhD thesis. University of Alexandria. 203 p.
- EL-GHAREEB, R. M. & K. SHALTOUT (1978). *Standing Crop biomass*. SAMDENE n° 4, 1: 4/1-4/74. University of Alexandria.
- EL-GHONEMY, A. A. (1973). Phytosociological and Ecological studies of the maritime sand dunes communities in Egypt. I Zonation of vegetation and soil along a dune side. *Bull. Inst. Désert Egypte* 23(2): 463-473.
- EL-GHONEMY, A. A. & A. S. EL-DIN (1978). *Root systems of perennials*. SAMDENE n° 4, 1: 8/1-8/18. University of Alexandria.
- EL-GHONEMY, A. A. & E. M. GEWAIFEL (1976). The phytosociological groups and their habitats at Sidi Barrani. *Egypt. J. Bot.* 19: 77-87.
- EL-GHONEMY, A. A. & H. F. EL-KADI (1978). *Range Management*. SAMDENE n° 4, 1: 10/1-10/20. University of Alexandria.
- EL-GHONEMY, A. A., K. SHALTOUT, W. VALENTINE & A. WALLACE (1977). Distribution pattern of *Thymelaea hirsuta* (L.) Endl. and associated species along the Mediterranean coast of Egypt. *Bot. Gaz.* 138: 479-489.
- EL-GHONEMY, A. A. & T. M. TADROS (1970). Socio-ecological studies of the natural plant communities along a transect between Alexandria and Cairo. *Bull. Fac. Sci. Cairo Univ.* 10: 329-407.
- EL-HADIDI, M. N. & M. A. AYYAD (1974). Floristic and ecological features of Wadi Habis (Egypt). In: GUINOCHET, M., G. GUITTONNEAU, P. OZENDA, P. QUEZEL & C. SAUVAGE (eds.), *La Flore du Bassin Méditerranéen: essai de systématique synthétique*: 247-258. Editions du CNRS.
- EL-KADY, H. F. (1980). *Effect of grazing pressure on certain ecological parameters of some fodder plants of the Mediterranean coast of Egypt*. MSc thesis. University of Tanta. 95 p.
- EL-KADY, H. F. (1987). *A study of range ecosystems of the western Mediterranean coastal desert of Egypt*. PhD thesis. Technical University of Berlin.

- EL-NAGA, M. A. (1980). *Nutritive value of the plants at Omayed*. REMDENE n° 1, 1: 5/1-5/4. University of Alexandria.
- EL-NAGA, M. A. (1981). *Nutrition of domestic animals*. REMDENE n° 2, 1: 3/1-3/8. University of Alexandria.
- EL-NAGA, M. A. (1982). *Nutrition of domestic animals*. REMDENE n° 3, 1: 4/1-4/26. University of Alexandria.
- EL-SHARKAWI, H. (1961). *Ecological and sociological studies of the plant communities of the Fuka-Ras el Hikma area western desert of Egypt*. MSc thesis. University of Alexandria.
- EL-SHARKAWI, H. (1977). *Water flow and productivity in the ecosystem*. SAMDENE n° 3, 1: 4/1-4/11. University of Alexandria.
- EL-SHARKAWI, H. (1978). *Water flow and productivity in the ecosystem*. SAMDENE n° 4, 1: 5/1-5/45. University of Alexandria.
- EL SHARKAWI, H. (1979). *Water flow and productivity*. SAMDENE n° 5, 1: 3/1-3/31. University of Alexandria.
- FAO (1970a). *Pre-investment survey of the northwest coastal region comprehensive account on the project*. Technical Report n° 1. ESE: SF/UAR 49, FAO, Rome. 109 p.
- FAO (1970b). *Pre-investment survey of the northwest coastal region, United Arab Republic*. Technical Report n° 3. ESE: SF/UAR 49, FAO, Rome. 335 p.
- FANTOLI, A. (1952) *Le piogge della Libia*. Arti Grafici Editrice.
- FATHY, A. M. & A. M. MEHANNA (1980). Water studies. In: AYYAD, M. A. (ed.), *Regional Environmental Management of Mediterranean Desert Ecosystems of Northern Egypt (SAMDENE)*. Progress report n° 1. University of Alexandria. 41 p.
- FLORET, C. (1981). The effects of protection on steppic vegetation in the Mediterranean arid zone of southern Tunisia. *Vegetatio*: 46.117-129.
- FLORET, C. & E. LE FLOCH (1973). *Production, sensibilité et évolution de la végétation et du milieu en Tunisie pré-saharienne*. Document n° 71. CEPE/CNRS, Montpellier. 45 p.
- FLORET, C., E. LE FLOCH & R. PONTANIER (1983). Phytomasse et production en Tunisie présaharienne. *Acta Oecol., Oecol. Pl.* 4 (18): 133-152.
- FLORET, C., E. LE FLOCH, R. PONTANIER & F. ROMANE (1978). Simulation of the impact of different levels of human pressure on the grazing lands in Southern Tunisia. In: Hyder, D. N. (ed.) *Proceedings of the 1st International Rangeland Congress*: 52-54. 1978 August 14-18. Society for Range Management, Denver.
- FLORET, C., E. LE FLOCH, F. ROMANE & R. PONTANIER (1981). Dynamique de systèmes écologiques de la zone aride. *Acta Oecol., Oecol. Appl.* 2: 195-214.
- GALAL, E. S. F., N. G. SELIGMAN, H. MANSOUR, A. M. ABOUL NAGA, A. M. EL-SERAFY & H. K. ALMADDY (1990). *Multiple criterium analysis of lamb production systems in the North-Western coastal zone in Egypt*. Cooperative Arid Lands Agriculture Research Program (CALAR), San Diego State University Foundation. 46 p.
- GHABBOUR, S. I. (1976). *Environmental and development of coastal desert*. International Seminar on Development and Management of Resources of coastal Areas, Berlin. 27 p.
- GHABBOUR, S. I. (1983). *Pre-proposal report for agricultural development in the NW coastal zone of Egypt: integrating barley cultivation and sheep herding*. Institute of Africa Development, University of Cairo. 49 p.
- GIRGIS, W. A. (1970). Phytosociological studies on the vegetation of Mariut area project. *Egypt. J. Bot.* 13: 235-254.
- GIRGIS, W. A. (1973). Phytosociological studies on the vegetation of Ras-el-Hikma-Mersa Matruh coastal plain. *Egypt. J. Bot.* 16: 393-409.
- HAMMOUDA, S.A.K. (1988). *A study of vegetation and land-use in the western Mediterranean desert of Egypt*. PhD thesis. University of Alexandria. 205 p.
- HERMANS, M. (1982). *An autoecological study of Mediterranean desert species at El Omayed, Egypt. III Gymnocarpos decandrum, Rumex pictus, Echiochilon fruticosum & Plantago albicans*. Student Report. University of Nijmegen.
- IBRAHIM, K. M. (1968). Some important native browse and forb forage plant of northern UAR. Their ecology in relation to range management. *Pakistan J. Forest.* 18: 155-168.
- IMAM, M. (1978). *Seed studies*. SAMDENE n° 2, 1: 6/1-6/66. University of Alexandria.
- IMAM, M. (1979). Seed and seedling studies. In: GHABBOUR, S. I. & M. KASSAS (eds.), *Analysis and Management of Mediterranean desert ecosystems*: 110-113. Proceedings of the International Workshop of SAMDENE Project. University of Alexandria.
- JAFRI, S. M. H. & A. A. EL GADI (eds) (1977-1989). *Flora of Libya*. Al Faateh University, Tripoli.
- KASSAS, M. (1971). Dynamic of desert vegetation. In: H. ODABASI & S. E. ULUG (eds.), *Environmental problems and their international implications*: 27-46. Colorado Association University Press.

- KASSAS, M. (1972). A brief history of land-use in Mareotis region, Egypt. *Minerva Biol.* 1: 167-174.
- KASSAS, M. (1979). Mareotis: past, present and future. In: GHABBOUR, S. I. & M. KASSAS (eds.), *Analysis and Management of Mediterranean desert ecosystems*: 24-28. Proceedings of the International Workshop of SAMDENE Project. University of Alexandria.
- KASSAS, M. & W. A. GIRGIS (1964). Habitat and plant communities in the Egyptian Desert.: V The limestone plateau. *J. Ecol.* 52: 107-119.
- KASSAS, M. & M. IMAM (1954). Habitat and plant communities in the Egyptian desert. III The wadi bed ecosystem. *J. Ecol.* 42: 424-441.
- KEITH, H. G. (1965). *A preliminary check list of the Libyan flora*. Ministry of Agriculture, Tripoli.
- LE HOUÉROU, H. N. (1959). *Recherches écologiques et floristiques sur la végétation de la Tunisie Méridionale*. PhD thesis. 2 vol. Institut de Recherches Sahariennes, Algier. 510 p.
- LE HOUÉROU, H. N. (1962). *Les pâturages naturels de la Tunisie aride et désertique*. Institut des Sciences Economiques Appliquées, Paris-Tunis. 106 p.
- LE HOUÉROU, H. N. (1965). *Improvement of natural pasture and fodder resources. Report to the Government of Libya*. EPTA report n° 1979. Plant Production & Protection Division, FAO. 51 p.
- LE HOUÉROU, H. N. (1969). La végétation de la Tunisie steppique. *Ann. Inst. Natl. Rech. Agron. Tunisie* 42(5): 1-624.
- LE HOUÉROU, H. N. (1975). Les pâturages naturels de l'Afrique du Nord: typologie, production, productivité et développement. In: *Inventaire et cartographie des pâturages tropicaux africains*: 41-55. Actes du Colloque de Bamako (Mali), 3-8 mars 1975. Centre International pour l'Élevage en Afrique (CIPEA), Addis Abeba.
- LE HOUÉROU, H. N. (1976). Contribution à une bibliographie écologique des régions arides de l'Afrique et de l'Asie du Sud-Ouest (végétation, pâturages, élevage, agriculture, nomadisme et désertisation). In: MONOD, T. (ed.), *La désertification au sud du Sahara*: 169-212. Actes du colloque tenu à Nouakchott, 17-19 décembre 1973. NEA, Dakar-Abidjan.
- LE HOUÉROU, H. N. (1984). Outline of the bioclimatology of Libya. *Bull. Soc. Bot. France, Actual. Bot.* 131: 157-178.
- LE HOUÉROU, H. N. (1989). Classification éoclimatique des zones arides (s.l.) de l'Afrique du Nord. *Ecol. Medit.* 15(3/4): 95-143.
- LE HOUÉROU, H. N. (1992a). Relations entre la variabilité des précipitations et celles de la production primaire et secondaire en zone aride. In: LE FLOC'H, E., M. GROUZIS, A. CORNET & J. C. BILLE (eds.), *L'aridité, une contrainte au développement*: 193-216. ORSTOM, Paris.
- LE HOUÉROU, H. N. (1992b). Rangeland management in Northern Africa and the Near East: evolution, trends and development outlook. In: GASTON, A., M. KERNICK & H. N. LE HOUÉROU (eds.), *Proceedings of the 4th International Rangeland Congress 1*: 544-553. CIRAD, Montpellier.
- LE HOUÉROU, H. N. (1992c). Feeding shrubs to sheep in the Mediterranean arid zone. In: GASTON, A., M. KERNICK & H. N. LE HOUÉROU (eds.), *Proceedings of the 4th International Rangeland Congress 2*: 639-644. CIRAD-UCIST, Montpellier.
- LE HOUÉROU, H. N. (1992d). The role of saltbushes (*Atriplex* spp.) in arid land rehabilitation in the Mediterranean Basin: a review. *Agroforest. Systems* 18: 107-148.
- LE HOUÉROU, H. N. (1995). *Bioclimatologie et biogéographie des steppes arides du Nord de l'Afrique - Diversité biologique, développement durable et désertisation*. CIHEAM, Montpellier.
- LE HOUÉROU, H.N. (1997a). *Germplasm for Arid Land Rehabilitation: NW Egypt and Jordan*. Facilitation Unit, ICARDA, Cairo. 91 p.
- LE HOUÉROU, H. N. (1997b). Biogeography of the arid steppeland North of the Sahara. In: BARAKAT, H. N. & A. K. HEGAZY (eds.), *Reviews in Ecology: Desert Conservation and Development*: 207-228. University of Cairo.
- LE HOUÉROU, H. N. (1998a). Restoration and rehabilitation of arid and semi-arid Mediterranean-type vegetation and ecosystems in Western Asia and Northern Africa (WANA): a review. *Arid Soils Res. Rehabil.* 14: 3-14.
- LE HOUÉROU, H. N. (1998b). *Fodder shrubs development in the NW coast of Egypt. Report to the Matrouh Resource Management Project*. Facilitation Unit, ICARDA, Cairo. 23 p.
- LE HOUÉROU, H. N. (1998c). *Fodder production from multipurpose trees and shrubs in the eastern slopes of the West Bank. Report to the Palestinian Authority*. Facilitation Unit, ICARDA, Cairo. 24 p.
- LE HOUÉROU, H. N. (1998d). *Outline of a strategy for rangeland and feed resources development in the Arabian Peninsula. Report to the Arabian Peninsula Regional Programme (APRP)*. ICARDA, Dubai. 15 p.

- LE HOUÉROU, H. N. (1998e). *Biodiversity: native plant Resource Conservation in Marmarica. Report to the Matrouh Resource Management Project*. ICARDA, Cairo. 13 p.
- LE HOUÉROU, H. N. (2001). Biogeography of the arid steppeland north of the Sahara. *J. Arid Environm.* 48: 103-128
- LE FLOC'H, E. (1981). *Mapping of land occupation*. REMDENE n° 2, 1: 5/1-5/2. University of Alexandria.
- LE FLOC'H, E. (1983). *Contribution à une étude ethnobotanique de la flore tunisienne*. Imprimerie Officielle de la République Tunisienne.
- LONG, G. A. (1955). The study of natural vegetation as a basis for pasture improvement in the western desert of Egypt. *Bull. Inst. Désert Egypte* 5(2): 18-45.
- LONG, G. A. (1955). *Grazing problems in the western desert of Egypt*. Plant Production & Protection Division, 55/4/2/2497, FAO. 37 p.
- LONG, G. A. & M. A. AYYAD (1956). *Phytoecological map of Ras el Hekma*. Plant Production & Protection Division, 56/9/7130, FAO. 38 p.
- LONG, G. A. (1978). *Premières propositions pour un programme de cartographie des ressources renouvelables et des terres concernant la zone cotière du désert occidental Egyptien*. Ecothèque Méditerranéenne, CEPE/CNRS, Montpellier. 21 p.
- LONG, G. A. (1979). Mapping of renewable resources for land development and land-use decision with special references to the coastal western desert of Egypt. In: GHABBOUR, S. I. & M. KASSAS (eds.), *Analysis and Management of Mediterranean desert ecosystems*: 37-60. Proceedings of the International Workshop of SAMDENE Project. University of Alexandria.
- MAGAZZINI, A. (1963). *Medie climatici mensili ed annuali per la Libia*. Meteorological Service Ministry of Communications, Tripoli. 36p.
- MIGAHID, A. M. (1958). *Plants as ground water indicators in desert areas*. Training course on hydrology of arid and semi-arid regions. Desert Research Institute, UNESCO, Cairo. 13 p.
- MIGAHID, A. M. & M. A. AYYAD (1959a). An ecological study of Ras el Hikma district. I The climatic environment of the vegetation. *Bull. Inst. Désert Egypte* 9(2): 1-45.
- MIGAHID, A. M. & M. A. AYYAD (1959b). An ecological study of Ras el Hikma district. II The factor of soil water content. *Bull. Inst. Désert Egypte* 9(2): 46-73.
- MIGAHID, A. M. & M. A. AYYAD (1959c). An ecological study of Ras el Hikma district. III Plant habitats and communities. *Bull. Inst. Désert Egypte* 9(2): 74-98.
- MIGAHID, A. M., A. A. ABD EL RAHMAN, A. M. EL-SHAFEI & M. A. HAMMOUDA (1955). Types of habitat and vegetation at Ras el Hikma. *Bull. Inst. Désert Egypte* 5(2): 107-190.
- MIGAHID, A. M., K. H. BATANOUNY & M. A. F. ZAKI (1971). Phytosociological and ecological study of a sector in the Mediterranean coastal region of Egypt. *Vegetatio* 23: 113-134.
- MIGAHID, A. M. & M. N. EL-SHOUBAGUI (1958). The ecological amplitude of the desert fodder grass «*Panicum turgidum*». II. Edaphic conditions. *Bull. Inst. Désert Egypte* 8(2): 21-65.
- MIGAHID, A. M., H. M EL-SHARKAWI, K. H. BATANOUNY & A. F. SHALABY (1963). Phytosociological and ecological study of Matkila sector of Sidi-Barrani. I. Sociology of the communities. *Feddes Repert.* 84: 747-760.
- PAMPANINI, R. (1931). *Prodromo della flora Cirenaica*. Forli.
- PAMPANINI, R. (1936). Aggiunte e correzioni al «prodromo delle Flora cirenaica». *Arch. Bot. (Forli)* 12: 17-53.
- PENMAN, H. L. (1948). Natural evaporation from open water, bare soil and grass. *Proc. Roy. Soc. London, Ser. B, Math. Phys. Sci.* 193: 120-145.
- REGNER, H. J. (1997). *Rainfall distribution in the North-South extension at Marsa-Matruh*. El Qasr Rural Development Project GTZ. Marsa Matruh and Eschborn. 16 p.
- SALEM, B. B. M. (1990). *Remote sensing and land-use in the North Western desert of Egypt*. PhD thesis. University of Alexandria.
- SHALTOUT, K. H. (1983). *An ecological study on Thymelaea hirsuta (L.) Endl. in Egypt*. PhD thesis. University of Tanta.
- SHALTOUT, K. H. (1987). Pattern, phenology and sex ratio of Egyptian Thymelaea hirsuta populations. *Vegetatio* 72: 67-73.
- SHALTOUT, K. H. & M. A. AYYAD (1988). Structure and standing crop of Egyptian Thymelaea hirsuta populations. *Vegetatio* 74: 137-142.
- SHALTOUT, K. H. & R. EL-GHAREEB (1984). Effet of protection on the phytomass and primary production of ecosystems of the western Mediterranean desert of Egypt. I Ecosystem of non-saline depressions. *Bull. Fac. Sci. Alexandria Univ.* 25: 109-131.

- SHOLTEN, M. E., M. W. A. C. HUKKELHOVEN, M. A. AYYAD & M. J. A. WERGER (1981). Vegetation analysis in the coastal dune ecosystem of the western Egyptian desert. *Folia Geobot. Phytotax.* 16: 293-308.
- TÄCKHOLM, V. (1974). *Student's flora of Egypt*. Ed. 2. Cairo University.
- TÄCKHOLM, V. & L. BOULOS (1972). Supplementary notes to student's flora of Egypt. Second edition. *Publ. Cairo Univ. Herb.* 5.
- TADROS, T. M. (1953). A phytosociological study of halophilous communities from Mareotis. *Vegetatio* 4: 103-124.
- TADROS, T. M. (1956). An ecological survey of the semi-arid coastal strip of the eastern desert of Egypt. *Bull. Inst. Désert Egypte* 6(2): 28-56.
- TADROS, T. M. (1958). *Plant communities a ground water indicators and the water economy of some desert communities*. Training course on hydrogeology of arid and semi-arid regions, Desert Research Center, El-Matariya & UNESCO, Cairo. 6 p.
- TADROS, T. M. & B. A. M. ATTA (1958a). Further contributions to the study of sociology and ecology of the halophilous plant communities from Mareotis. *Vegetatio* 8: 137-160.
- TADROS, T.M. & B. A. M. ATTA (1958b). Plant communities of barley fields and uncultivated desert areas of Mareotis. *Vegetatio* 8: 161-175.
- TADROS, T.M. & H. M. EL SHARKAWI (1960a). Phytosociological and ecological studies on the vegetation of Fuka-Ras El Hekma area. I Sociology of the communities. *Bull. Inst. Désert Egypte* 10(1): 37-60.
- TADROS, T. M. & H. M. EL-SHARKAWI (1960b). Phytosociological and ecological studies on the vegetation of Fuka-Ras El Hekma area. II Consistency and homogeneity of the open desert communities. *Bull. Inst Désert Egypte* 10(1): 61-83.
- TADROS, T.M. & H. M. EL-SHARKAWI (1960c). Phytosociological and ecological studies on the vegetation of Fuka-Ras el Hekma area. III A vegetation map of the area. *Bull. Inst. Désert Egypte* 10(1): 85-93.
- VAN DUIVENBODEN, N. (1985). *Animal Husbandry in the North-Western Coastal Zone of Egypt*. Theoretical Production Ecology. CABO, Wageningen. 52 p.
- VAN DUIVENBODEN, N. (1987). *Evaluation of Animal Husbandry in the North-Western Coastal Zone of Egypt. (Literature Review and Simulation)*. CABO Publishing n° 72, Wageningen. 204 p.
- VAN DUIVENBODEN, N. (1988). *The Effect of Grazing on the Growth of Subshrubs and the Water Balance in the North Western Coastal Zone of Egypt. (Measurement and Simulation)*. Theoretical Production Ecology. CABO, Wageningen. 191 p.
- VAN DUIVENBODEN, N. (1989). Contributions of various feed components to feed availability in integrated barley/livestock systems in the north-western coastal zone of Egypt: a simulation study. *J. Arid Environm.* 16: 217-228.
- VAN KEULEN, H. & G. W. J. VAN DE VEN (1988). Application of interactive multiple goal linear programming techniques for analysis and planning of regional agricultural development: a case study for the Mariut region (Egypt). In: J. M. BOUSSARD (ed.), *Agriculture: Socio-Economic Factors in Land Evaluation*: 36-57. Proceedings of a conference held in Brussels 7-9 oct. 1987. OPOCE, Luxembourg.
- VAN DER VEEN, J. P. H., Z. ABU GUENDIA & A. NASSEF, A. (1968). *Report on range carrying capacity survey, NW Coastal Zone*. UAR. EGDDO/UNSF/FAO Pre-investment survey of NW coastal region project. UAR, FAO, Rome, Ministry of Agriculture, Cairo. 22 p. & 1 map 1:500000.
- WALTER H. & H. LIETH (1960). *Klimadiagramm Weltatlas*. Veb Gustav Fisher Verlag.
- ZAHRAN, M. A. (1986). Forage potentialities of *Kochia indica* and *K. prostrata* in arid land, with particular reference to Saudi Arabia. *Arab Gulf J. Sci. Res.* 4: 53-68.
- ZAHRAN, M.A., B. K. MUHAMMAD & A. A. EL-DINGAWI (1992). Establishment of *Kochia* forage halophytes in the salt affected lands of the Arab countries. *J. Environm. Sci. Mansoura Univ.* 4: 93-113.
- ZAHRAN, M.A. & A. J. WILLIS (1992). *The vegetation of Egypt*. Chapman & Hall.

Table 3. – Plant biodiversity table (see pp. 282-307)

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Adiantum	capillus-venereis	Adiantaceae	Kozbaritel Bir	FC	I-VI	F	P	Can-Wad	Orn	0	Hyg
Aizoon	canariense	Aizoaceae	Ghassool	CC	I-III	CH	A	Ste-Wid	Non	0	Nit
Aizoon	hispanicum	Aizoaceae	Ghassool	FR	I-II	CH	A	Ste-Wid	Non	0	Nit
Mesembryanthemum	crystallinum	Aizoaceae	Ghassool	CC	I-III	CH	A	Wid	Non	0	Hal
Mesembryanthemum	nodiflorum	Aizoaceae	Ghassool	CC	I-III	CH	A	Rge-Wid	Non	0	Nit
Opophytum	forskallii	Aizoaceae	Chassool	CC	I-III	CH	A	St-Wid	Non	0	Hal
Alisma	gramineum	Alismaceae	-	FR	I-II	H	P	Can-Frw	Non	0	Hyd
Damasonium	alisma	Alismaceae	-	FR	I-II	H	P	Can-Frw	Non	0	Hyd
Amaranthus	caudatus	Amaranthaceae	Fisa al Kalb	FC	I-II	H	A	Wid	Non	0	Wee
Amaranthus	cruentus	Amaranthaceae	Ruaaf	C	I-II	H	A	Wid	Non	0	Wee
Amaranthus	graecizans	Amaranthaceae	-	CC	I-II	H	A	Wid	Non	0	Wee
Amaranthus	retroflexus	Amaranthaceae	-	R	I-II	H	A	Wid	Non	0	Wee
Amaranthus	silvestris	Amaranthaceae	-	C	I-II	H	A	Wid	Non	0	Wee
Amaranthus	tricolor	Amaranthaceae	-	R	I-II	H	A	Wid	Non	0	Wee
Amaranthus	viridis	Amaranthaceae	Amarantoun	CC	I-II	H	A	Wid	Non	0	Wee
Cornulaca	monacantha	Amaranthaceae	Haad	FC	III-IV	CS	P	Des-Snd	Gra-Med	3	Gly-Psa
Narcissus	tazetta	Amaryllidaceae	Nargis	R	I-II	H	P	Wet-Wid	Med-Orn	T	Hyg
Panocratum	arabicum	Amaryllidaceae	Sawsan	R	I-II	H	P	Coa-Snd	Orn	T	Wee
Panocratum	maritimum	Amaryllidaceae	Sawsan	CC	I	H	P	Coa-Snd	Orn	T	Hal
Pistacia	atlantica	Anacardiaceae	Batoum	RR	I-III	T	P	Des-Wad	Fue-Gra-Med	2	Phr
Pistacia	lentiscus	Anacardiaceae	Za'roor, Batoum Derw	RR	I-III	HS	P	Cl-Wad	Fue-Gra-Med	2	Cha
Rhus	tripartita	Anacardiaceae	Areen, Gdari, Ern	RR	I-III	HS	P	Cl-Run-Wad	Fen-Gra-Med	2	Cha
Nerium	oleander	Apocynaceae	Defia	R	I-II	HS	P	Run-Wad	Med-Orn	T	Nat-Phr
Arisarum	vulgare	Araceae	Netirish	C	I-II	H	P	Rge-Wid	Med	T	Wee
Eminium	speculatum	Araceae	Erqueta	C	I-II	H	P	Wet	Non	0	Wee
Asclepias	fruticosa	Asclepiadaceae	Gheil	R	I-II	HS	P	Roc	Non	T	Cha
Calotropis	procera	Asclepiadaceae	Asclepiad.Krom Berry	FR	III-VI	HS	P	Irr	Non	T	Wee
Caralluma	europaea	Asclepiadaceae	Dahmous	R	I-III	Su	P	Rge-Roc	Non	0	Cha
Cynanchum	acutum	Asclepiadaceae	Ollaiq, Muadded	FC	I-III	H	P	Irr	Non	T	Wee
Pergularia	tomentosa	Asclepiadaceae	-	FC	I-III	Fr	P	Ste	Med	T	Wee
Periploca	angustifolia	Asclepiadaceae	Haliaba	FR	I-III	HS	P	Cl-Roc-Run	Fsh-Gra	4	Cha
Leontice	leontopetalum	Berberidaceae	Rakaf	R	I-II	H	P	Cul-Wid	Orn	0	Wee
Alkanna	lehmannii	Boraginaceae	Hinna el Ghoul	CC	I-II	H	P	Cul-Rge	Gra-Med	1	Psa
Alkanna	orientalis	Boraginaceae	Libbeid	R	I-III	H	P	Cl-Run	Gra	2	Cha
Anchusa	aegyptiaca	Boraginaceae	Kahla	CC	I-III	H	A	Cul-Rge	Gra	1	Psa
Anchusa	hispidia	Boraginaceae	Kahla	C	I-II	H	A	Cul-Rge	Gra	2	Psa
Anchusa	humilis	Boraginaceae	Kahla	C	I-II	H	A	Cul-Rge	Gra	1	Psa
Anchusa	milleri	Boraginaceae	Kahla	FC	I-II	H	A	Cul-Rge	Gra	1	Psa
Anchusa	undulata	Boraginaceae	Kahla	R	I-III	H	A-B	Cul-Rge	Gra	1	Psa
Arnebia	decumbens	Boraginaceae	Attan	FC	I-III	H	A	Rge-Snd-Ste	Gra	3	Psa-Tac

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Arnebia	linearifolia	Boraginaceae	Kahla	FC	I-II	H	A	Rge-Snd	Gra	2	Psa
Borago	officinalis	Boraginaceae	-	FR	I-II	H	A	Cul	Non	1	Psa
Buglossoides	incrassatum	Boraginaceae	-	R	I-III	H	A	Cul-Rge	Non	0	Wee
Buglossoides	tenuiflora	Boraginaceae	-	FR	I-III	H	A	Cul-Rge	Non	0	Wee
Echium	angustifolium	Boraginaceae	-	CC	I-II	H	P	Cul-Rge-Wid	Non	0	Wee
Echium	glomeratum	Boraginaceae	-	R	I-II	H	P	Wid	Non	0	Wee
Echium	plantagineum	Boraginaceae	Lisan el Thour	RR	I-II	H	A	Cul-Wid	Gra-Med	1	Wee
Echium	rauwolfi	Boraginaceae	Hinna el Ghoul	CC	I-II	H	A-B	Cul-Rge-Ste	Gra	1	Wee
Echium	rubrum	Boraginaceae	-	C	I-II	H	A	Wid	Non	0	Wee
Echium	sabulicola	Boraginaceae	-	R	I-II	H	A	Wid	Non	0	Wee
Echium	sericeum	Boraginaceae	-	C	I-III	H	P	Rge-Snd-Ste-Wid	Non	0	Wee
Echlochilon	fruticosum	Boraginaceae	Okroosh	FC	I-II	DS	P	Rge-Snd-Ste	Gra	4	Psa
Heliotropium	arbatense	Boraginaceae	Atana	R	III-IV	Fr	P	Des-Wad	Gra	2	Ere
Heliotropium	curassavicum	Boraginaceae	-	R	I-II	H	P	Sit	Non	1	Hal
Heliotropium	dignum	Boraginaceae	Rogh, S'gaa	CC	III-VI	Fr	P	Des-Rge-Ste	Gra-Med	2	Ere
Heliotropium	europaeum	Boraginaceae	Aqrabana	FC	I-II	H	A	Cul	Non	0	Wee
Heliotropium	hirsutissimum	Boraginaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Heliotropium	supinum	Boraginaceae	Koddeih, Zurreqa	C	I-III	H	A	Cul	Non	0	Psa
Lappula	spinocarpos	Boraginaceae	Qaliet	C	I-III	H	A	Rge-Ste	Gra	2	Psa-Tac
Motkiopsis	cliata	Boraginaceae	Halama	FR	III-VI	Fr	P	Des-Rge-Ste	Gra-Med	3	Psa
Nonea	melanocarpa	Boraginaceae	-	FR	I-II	H	A	Cul-Rge	Gra	1	Wee
Nonea	vivianii	Boraginaceae	-	FR	I-II	H	A	Rge-Snd-Ste	Gra	2	Psa
Ogastemma	pusillum	Boraginaceae	-	R	III-VI	H	A	Rge-Ste	Gra	2-3	Psa-Tac
Trichodesma	africanum	Boraginaceae	Hameem	FC	III-VI	H	A-B	Des-Wad	Gra	1-2	Psa
Campanula	erinus	Campanulaceae	-	RR	I-II	H	A	Irr-Wid	Non	0	Wee
Capparis	rupestris	Capparaceae	Kabbar	CC	I-III	TS	P	Cil-Roc-Wad	Med-Orn	0	Cha
Capparis	spinosa	Capparaceae	Kabbar	C	III-VI	TS	P	Cil-Roc-Wad	Med-Orn	0	Ere
Arenaria	serpyllifolia	Caryophyllaceae	Labbaikh	FR	I-II	H	A	Cul	Gra	0	Wee
Gymnocarpos	decander	Caryophyllaceae	Jarad Garad	CC	I-IV	DS	P	Sha-Ste	Gra	4	Ere
Herniaria	cinerea	Caryophyllaceae	-	FC	I-II	H	A	Rge-Sha	Med	0	Wee
Herniaria	cyrenaica	Caryophyllaceae	-	RR	I-II	H	P	Rge-Sha	Gra	1	Ere
Herniaria	fontanesii	Caryophyllaceae	-	RR	I-III	H	P	Rge-Sha	Gra	1	Ere
Herniaria	hemistemon	Caryophyllaceae	-	CC	I-III	H	P	Rge-Sha	Gra	1	Ere
Herniaria	hirsuta	Caryophyllaceae	Mouker, Makir	FC	I-III	H	A	Cul	Med	2	Wee
Loefflingia	hispanica	Caryophyllaceae	-	FC	I-III	H	A	Rge-Snd-Ste	Gra	1	Tac
Minuartia	hybrida	Caryophyllaceae	-	RR	I-II	H	A	Cil-Roc-Run	Gra	1-2	Cha-Psa
Minuartia	mediterranea	Caryophyllaceae	-	RR	I-II	H	A	Cil-Roc-Run	Gra	1-2	Cha
Paronychia	arabica	Caryophyllaceae	Ramram	C	I-III	H	A-B	Snd-Ste	Gra-Med	1-2	Psa
Paronychia	argentea	Caryophyllaceae	Bessat el Ard, Tai el Areb	FC	I-III	H	P	Rge-Ste	Gra	2	Psa
Paronychia	capitata	Caryophyllaceae	-	R	I-III	H	P	Snd-Ste	Gra-Med	1-2	Psa

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Petrorhagia	ilyrica	Caryophyllaceae	Sammah	FR	I-II	Fr	P	Cli-Run	Gra	2	Cha
Polycarphaea	robairaea	Caryophyllaceae	Mokoor	R	III-VI	H	A-Sh	Des-Rge	Gra	2	Ere-Tac
Polycarpon	succulentum	Caryophyllaceae	-	CC	I-II	Su	A	Cul-Snd	Non	0	Gly-Hal-Wee
Polycarpon	tetraphyllum	Caryophyllaceae	-	C	I-II	Su	A	Cul-Snd	Non	0	Gly-Hal-Tac-Wee
Pteranthus	dochotomus	Caryophyllaceae	-	FC	I-IV	CH	A	Sit	Non	0	SO4
Rhodoalsine	geniculata	Caryophyllaceae	Heriba	FR	I-II	H	A	Cli-Roc-Run	Non	1-2	Cha-Psa
Silene	aegyptiaca	Caryophyllaceae	-	RR	I-III	H	A	Cul	Non	0	Wee
Silene	apetala	Caryophyllaceae	Deheina	FC	I-II	H	A	Cul	Non	0	Wee
Silene	behen	Caryophyllaceae	Deheina	R	I-II	H	A	Cul	Non	1	Wee
Silene	biappendiculata	Caryophyllaceae	Deheina	C	I-II	H	A	Cul	Non	0	End-Wee
Silene	colorata	Caryophyllaceae	Deheina	C	I-III	H	A	Cul-Snd-Ste	Non	0	Wee
Silene	fruticosa	Caryophyllaceae	-	RR	I-II	Fr	P	Cli-Roc	Gra	1-2	Cha-End
Silene	nocturna	Caryophyllaceae	Deheina, Abul Nagaf	FC	I-II	H	A	Cul	Non	0	Wee
Silene	rubella	Caryophyllaceae	Naashash	C	I-II	H	A	Cul-Rge	Gra	0	Wee
Silene	succulenta	Caryophyllaceae	Atar el Bhar	C	I	Su	P	Coa-Snd	Non	0	Psa
Silene	tridentata	Caryophyllaceae	Deheina	R	I-II	H	A	Cul-Rge	Non	0	Wee
Silene	villosa	Caryophyllaceae	Attan	R	II-IV	H	A	Con-Snd	Gra	1-2	Psa
Silene	vivianii	Caryophyllaceae	Deheina	FC	I-II	H	A	Cul	Non	0	Wee
Silene	vulgaris	Caryophyllaceae	-	R	I-II	H	P	Cli	Gra	1-2	Cha
Spergula	fallax	Caryophyllaceae	Rihay	C	I-II	H	A	Cul	Non	0	Gly-Hal-Wee
Spergularia	bocconeii	Caryophyllaceae	-	R	I-II	H	A	Cul	Non	0	Hal-Wee
Spergularia	diandra	Caryophyllaceae	-	CC	I-II	H	A	Cul-Wid	Non	0	Gly-Hal-Wee
Spergularia	marina	Caryophyllaceae	Abu Ghulam	CC	I-III	CH	P	Sit	Gra	3-4	Hal
Spergularia	maritima	Caryophyllaceae	Mesht	R	I-III	CH	P	Sit	Gra-Med	3-4	Nit-Wee
Spergularia	rubra	Caryophyllaceae	Esbba Hamra	R	I-II	H	A	Cul	Med-Non	0	Hal-Wee
Stellaria	media	Caryophyllaceae	Qizzaza, Bughlan	R	I-II	CH	A	Cul-Wid	Med	0	Nit
Stellaria	pallida	Caryophyllaceae	Qizzaza	CC	I-II	CH	A	Cul-Wid	Non	0	Wee
Vaccaria	pyramidata	Caryophyllaceae	Fool el Arab	C	I-II	H	A	Cul	Med	0	Wee
Ceratophyllum	demersum	Ceratophyllaceae	Horeish	C	I-II	H	P	Frw	Non	0	Hyd
Anabasis	articulata	Chenopodiaceae	Belbel	CC	II-IV	CS	P	Roc-Sha-Ste	Gra	1	Gly
Anabasis	oropediorum	Chenopodiaceae	A'jrem	CC	I-II	DS	P	Roc-Sha-Ste	Gra	3	Gly-Sil
Arthrocnemum	macrostachyum	Chenopodiaceae	Shinaan	C	I-III	CS	P	Sit	Gra	1	Hal
Atriplex	coriacea	Chenopodiaceae	Ratam	R	I-II	CS	P	Sit-Wet	Gra	2	Hal
Atriplex	glauca	Chenopodiaceae	Qetteifa	FR	I-III	TS	P	Rge-Ste	Fsh-Gra	3	Hal
Atriplex	halimus	Chenopodiaceae	Qattaf	CC	I-IV	CS	P	Rge-Sit	Fsh-Gra-Med	3	Hal
Atriplex	inflata	Chenopodiaceae	-	FR	I-III	CH	A	Wid	Non	0	Hal-Nit-Wee
Atriplex	leucoclada	Chenopodiaceae	Roghil	R	I-II	HS	A	Rge	Fsh-Gra	2-3	Hal
Atriplex	littoralis	Chenopodiaceae	-	R	I-II	H	A	Cul-Wid	Non	0	Hal-Nit
Atriplex	mollis	Chenopodiaceae	Quattaf	R	I-II	TS	Sh	Rge-Ste	Fsh-Gra	3	Hal
Atriplex	portulacoides	Chenopodiaceae	Qattaf bahri	C	I	TS	P	Coa-Sit	Gra	2	Hal

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Atriplex	rossa	Chenopodiaceae	-	R	I-II	H	A	Cul-Wid	Non	0	Hal-Nit-Wee
Atriplex	semibaccata	Chenopodiaceae	Qattaf	FR	I-II	TS	P	Sit-Ste	Gra	3	Hal-Nat
Atriplex	spongiosa	Chenopodiaceae	-	R	I-II	CH	A-B	Wid	Non	0	Hal-Nit-Wee
Bassia	arabica	Chenopodiaceae	Thelata	C	I-II	DS	P	Rge-Sit-Ste	Gra	0	Hal
Bassia	indica	Chenopodiaceae	-	FC	I-II	H	A-Sh	Rge-Wid	Gra	2-3	Nat-Wee
Bassia	muricata	Chenopodiaceae	Ghobbeira	CC	I-IV	H	A	Snd-Ste	Non	2-3	Gly
Beta	vulgaris	Chenopodiaceae	Salq, Dirs el Kalb	FC	I	H	A	Sit	Gra	2	Hal
Blitum	virgatum	Chenopodiaceae	-	R	I-II	H	A	Cul	Non	0	Nit
Chenopodium	album	Chenopodiaceae	Rokaab al Jamal	CC	I-III	H	A	Cul-Wid	Non	0	Nit-Wee
Chenopodium	ambrosoides	Chenopodiaceae	Nifna, Natna	CC	I-II	H	A	Wid	Med	0	Nit-Wee
Chenopodium	giganteum	Chenopodiaceae	-	RR	I-II	H	A	Wid	Non	0	Nit-Wee
Chenopodium	glaucum	Chenopodiaceae	-	R	I-III	H	A	Cul-Wid	Non	0	Nit-Wee
Chenopodium	murale	Chenopodiaceae	Lisaan et Teir, Abu Ifen	CC	I-II	H	A	Wid	Non	0	Nit-Wee
Halocnemum	strobilaceum	Chenopodiaceae	Sabad	C	I-III	CS	P	Sit	Non	0	Hal
Halopeplis	amplexicaulis	Chenopodiaceae	-	R	I-II	CH	A	Sit	Non	0	Hal
Halopeplis	perfoliata	Chenopodiaceae	-	R	I-II	CS	P	Sit	Non	0	Hal
Haloxylon	salicornicum	Chenopodiaceae	Belbel	CC	II-IV	CS	P	Rge-Snd-Ste	Fue-Gra	1	Gly
Haloxylon	scoparium	Chenopodiaceae	Remeth	CC	I-IV	DS	P	Rge-Ste	Fue-Gra-Med	1	Gly-Sil
Noaea	mucronata	Chenopodiaceae	Athir	CC	I-IV	DS	P	Roc-Sha-Ste	Fue-Gra	2	Gly
Salicornia	europaea	Chenopodiaceae	Hamd	C	I-II	CH	A	Sit	Non	0	Hal
Salicornia	fruticosa	Chenopodiaceae	Hamd	C	I-II	CS	P	Sit	Gra	1	Hal
Salicornia	perennis	Chenopodiaceae	Hamd	C	I-II	CS	P	Sit	Gra	1	Hal
Salsola	imbricata	Chenopodiaceae	Woqeid	CC	III-VI	CS	P	Des-Rge	Gra	0-1	Hal-Pel
Salsola	inermis	Chenopodiaceae	-	FC	I-II	CH	A	Roc-Sha	Non	0	Hal-Pel
Salsola	kali	Chenopodiaceae	Eshmaan	CC	I	CH	A	Coa-Dne	Non	0	Hal-Psa
Salsola	longifolia	Chenopodiaceae	Heddeid	FC	I-III	CS	P	Sit-Ste	Gra	2-3	Hal
Salsola	oppositifolia	Chenopodiaceae	-	R	I-II	CS	P	Ste-Wid	Non	0	Nit
Salsola	schweinfurthii	Chenopodiaceae	-	RR	II-IV	CS	P	Rge-Sit-Ste	Gra	1	Hal
Salsola	tetragona	Chenopodiaceae	-	RR	I-IV	CS	P	Sit	Fue-Gra	1	Hal-Phr
Salsola	tetrandra	Chenopodiaceae	Damaan	R	I-II	CS	P	Sit	Gra	2	Hal
Salsola	villosa	Chenopodiaceae	Khareet	CC	II-III	CS	P	Rge-Sha-Ste	Gra	2-3	Hal
Salsola	volkensi	Chenopodiaceae	Caremba	C	I-II	CH	A	Coa-Dne-Wad	Non	0	Hal-Psa-Wee
Suaeda	aegyptiaca	Chenopodiaceae	Soweid	CC	I-II	CH	A	Sit	Non	0	Hal
Suaeda	altissima	Chenopodiaceae	Soweid	RR	I-III	CH	A	Sit	Non	0	Hal
Suaeda	palaestina	Chenopodiaceae	Soweid	RR	I-III	CS	P	Sit	Gra	1	Hal
Suaeda	pruinosa	Chenopodiaceae	Soweid	R	I-III	CS	P	Sit	Gra	1	Hal
Suaeda	splendens	Chenopodiaceae	Soweid	RR	I-II	CH	A	Sit	Non	0	Hal
Suaeda	vera	Chenopodiaceae	Soweid	C	I-III	CS	P	Sit	Gra	1	Hal
Suaeda	vermiculata	Chenopodiaceae	Soweid	C	I-III	CS	P	Sit	Gra	1	Hal
Traganum	nudatum	Chenopodiaceae	Dhamran	FC	II-IV	CS	P	Rge-Ste	Gra	2-3	Hal-S04

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Fumana	arabica	Cistaceae	-	C	I-II	Fr	P	Rge-Sha	Gra	2	Cha
Fumana	thymifolia	Cistaceae	-	R	I-II	Fr	P	Rge-Sha	Gra	2	Cha
Helianthemum	aegyptiacum	Cistaceae	Koshein	R	I-II	H	A	Rge	Gra	2	Psa
Helianthemum	ciliatum	Cistaceae	Koshein	C	I-II	Fr	P	Rge-Sha	Gra	2-3	Cha
Helianthemum	kahiricum	Cistaceae	Koshein	CC	I-III	Fr	P	Rge-Sha-Ste	Gra	3	Cha
Helianthemum	ledifolium	Cistaceae	Koshein	R	I-II	H	A	Rge-Sha-Ste	Gra	2	Psa-Tac
Helianthemum	lippii	Cistaceae	Qadib el Raal	CC	I-III	Fr	P	Rge-Snd-Ste	Gra	3-4	Psa
Helianthemum	salicifolium	Cistaceae	Koshein	R	I-II	H	A	Cul-Rge	Gra	2	Psa
Helianthemum	sessiliflorum	Cistaceae	Qadib el Raal	FR	I-II	Fr	P	Rge-Snd-Ste	Gra	3-4	Psa
Helianthemum	sphaerocalyx	Cistaceae	Koshein	R	I-II	Fr	P	Rge-Ste	Gra	3	Cha-End
Helianthemum	stipulaceum	Cistaceae	Koshein	FC	I-II	Fr	P	Rge-Sha	Gra	2-3	Cha
Helianthemum	vesicarium	Cistaceae	Koshein	FC	I-II	Fr	P	Rge-Sha-Ste	Gra	2-3	Cha
Cleome	amblyocarpa	Cleomaceae	Zeita, Magnuna	CC	I-III	H	A	Cul-Ste-Wid	Med	T	Psa-Wee
Gynandropsis	gynandra	Cleomaceae	Abu Qara, Abu Qarn	R	I-III	H	A	Wid	Non	0	Nat-Nit-Wee
Achillea	fragrantissima	Compositae	Ge Soom	RR	III-VI	H	P	Des-Wid	Gra-Med	1	Ere
Achillea	santolina	Compositae	Bishreen, Chalhata	R	I-II	H	P	Cul-Wid	Med	0	Sil
Amberboa	crupinoides	Compositae	-	RR	I-II	H	A	Cul-Wid	Non	0	Sil
Amberboa	lippii	Compositae	Kerzaran	CC	I-III	H	A	Cul-Ste-Wid	Non	0	Sil
Ambrosia	maritima	Compositae	Damaseisa	R	I	H	B	Coa-Snd	Med	0	Psa
Anacyclus	alexandrinus	Compositae	Rigel Ghorab	CC	I-III	H	A	Rge-Ste-Wid	Gra	2	Wee
Anthemis	chia	Compositae	Arbeyaan	RR	I-II	H	A	Cul-Wid	Non	0	Wee
Anthemis	indurata	Compositae	Arbeyana	R	I-II	H	A	Cul-Wid	Non	0	Wee
Anthemis	melampodina	Compositae	Firaakomm Ali	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Anthemis	microsperma	Compositae	Arbeuaan	C	I-II	H	A	Cul-Wid	Non	0	Wee
Anthemis	pseudo-cotula	Compositae	Basoon, Iribayan	CC	I-II	H	A	Cul-Wid	Med	0	Wee
Anvillea	garcini	Compositae	Nogod	R	III-VI	Fr	P	Des-Wad	Non	0	Ere
Artemisia	herba-alba	Compositae	Sheeh	C	I-III	H	P	Rge-Ste	Fsh-Gra-Med	3	Sil
Artemisia	judaica	Compositae	Sheeb	FC	III-VI	H	P	Wad	Gra-Med	1	Ere
Artemisia	monosperma	Compositae	Aader	FC	III-VI	H	P	Des-Rge-Ste	Gra	0	Ere-Psa
Artemisia	scoparia	Compositae	-	RR	I-II	Fr	Sh	Wid	Non	0	Wee
Aster	squamatus	Compositae	-	FC	I-II	H	B	Wid	Non	0	Hal-Hyg-Sum
Asteriscus	pygmaeus	Compositae	Musmar el Ard	CC	I-III	H	A	Rge-Wid	Non	0	Ere-Sil
Asteriscus	spinus	Compositae	-	CC	I-III	H	A	Wid	Non	0	Nit
Atractylis	cancelata	Compositae	-	FR	I-II	H	A	Rge-Wid	Non	0	Wee
Atractylis	carduus	Compositae	Akaash	CC	I-III	H	P	Rge-Wid	Non	0	Wee
Atractylis	prolifera	Compositae	-	CC	I-III	H	A	Cul-Rge-Ste	Non	0	Wee
Bidens	pilosa	Compositae	Sada	FC	I-II	H	A	Wet-Wid	Non	0	Hyg-Wee
Calendula	arvensis	Compositae	A0safra, Ayn el Bagar	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Carduncellus	ericephalus	Compositae	Kharsroof	C	II-IV	H	P	Cul-Wad	Non	0	Ere
Carduncellus	mareoticus	Compositae	Sarr	CC	I-V	H	P	Cul-Ste-Wad-Wid	Non	0-1	End-Sil-Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Carduus	getulus	Compositae	Horshof	CC	I-III	H	A	Cul-Wid	Non	0	Wee
Carduus	pynoccephalus	Compositae	Lisaanelkelb	C	I-II	H	A	Cul-Wid	Non	0	Wee
Carlina	involutrata	Compositae	Moraar	CC	I-III	H	P	Rge-Ste-Wid	Non	0	Wee
Carthamus	glaucus	Compositae	-	FR	I-II	H	A	Cul-Wid	Non	0	Wee
Carthamus	lanatus	Compositae	Shook Antar	CC	I-III	H	A	Cul-Wid	Non	0	Wee
Carthamus	tenuis	Compositae	Qoos	FC	I-II	H	A	Cul-Wid	Med	0	Wee
Centaurea	aegialophila	Compositae	Akash	CC	I	H	P	Coa-Snd	Med	0	Hal-Psa
Centaurea	alexandrina	Compositae	Moraar	CC	I-III	H	A	Cul-Wid	Med	0	Wee
Centaurea	calcitrapa	Compositae	Moraar, Sh,k	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Centaurea	dimorpha	Compositae	Moraar	C	I-II	H	P	Cul-Ste-Wid	Non	0	Psa
Centaurea	furfuracea	Compositae	-	FC	II-IV	H	A	Cul-Rge-Ste	Non	0	Psa
Centaurea	glomerata	Compositae	Yamaar, Arjojaq,	C	I-II	H	A	Cul-Wid	Med	0	Wee
Centaurea	melitensis	Compositae	Moraar	FR	I-II	H	A	Cul-Wid	Non	0	Sil
Centaurea	pullata	Compositae	Moraar	C	I-II	H	B	Cul	Non	0	Wee
Centaurea	pumilio	Compositae	Moraar	FR	I-II	H	P	Cul-Wid	Non	0	Wee
Centaurea	sositalialis	Compositae	Moraar	RR	I-II	H	A	Cul-Wid	Non	0	Wee
Ceruaea	pratensis	Compositae	Garawaan	R	I-II	H	A	Cul-Wid	Non	0	Wee
Chlamidophora	tridentata	Compositae	-	R	II-IV	H	A	Rge-Ste	Gra	2	Ere
Chrysanthemum	coronarium	Compositae	Oqhtwan, Reza'o	CC	I-II	H	A	Cul-Wid	Med	0	Wee
Cichorium	endivia subsp. Pumilum	Compositae	Shikoria, Sires	FC	I-II	H	A-B	Cul-Wid	Gra-Med	2-3	Per-Sil
Conyza	bonariensis	Compositae	Hashishat el Gabal	C	I-II	H	A	Cul-Wid	Med	0	Wee
Cotula	anthemoides	Compositae	Ribyaan	C	I-II	H	A	Cul-Wet	Med	0	Hyg
Cotula	cinerea	Compositae	Sakaraan, Roboua,	CC	III-IV	H	P	Des-Rge	Gra-Med	1	Ere
Crepis	bulbosa	Compositae	-	FR	I	H	P	Snd-Wid	Gra	1-2	Psa
Crepis	clausomis	Compositae	-	RR	I-II	H	P	Cul-Rge	Gra	1	Wee
Crepis	libyca	Compositae	-	R	I-II	H	P	Cul-Rge-Ste	Gra	2	Tac-Wee
Crepis	micrantha	Compositae	-	FC	I-II	H	A	Cul-Rge	Gra	2	Wee
Crupina	crupinastrum	Compositae	-	RR	I-II	H	A	Cul	Non	0	Sil
Cynara	cornigera	Compositae	Karshoof, Khorshef	FC	I-II	H	P	Cul-Rge-Wid	Med-Veg	0	Per-Wee
Echinops	galatensis	Compositae	Kasheer	C	I-III	H	P	Cul-Rge	Non	0	Wee
Echinops	spinossimus	Compositae	Kaddad	CC	I-III	H	P	Cul-Rge	Med	0	Wee
Echinops	spinosis	Compositae	Kasheer, Sh,k el Gamel	CC	I-III	H	P	Rge-Sha	Med	0	Wee
Echinops	taeckholmianus	Compositae	Korshof	R	I-II	H	P	Rge-Wid	Non	0	End
Eclipta	alba	Compositae	Sada, Suwwed	R	I	H	A	Wet-Wid	Med	0	Hyg-Wee
Ethulia	conyzoides	Compositae	Hashishel Faras	C	I-II	H	A	Cul	Non	0	Wee
Filago	contracta	Compositae	-	FC	I-III	H	A	Cul-Rge	Non	2	Wee
Filago	desertorum	Compositae	-	CC	I-III	H	A	Cul-Wid	Non	0	Sil-Tac
Filago	mareotica	Compositae	-	C	I-II	H	A	Sit	Non	0	Hal
Flaveria	contrayerba	Compositae	Sada	R	I-II	H	A	Wet-Wid	Non	0	Hyg-Sum
Garhadiolus	angulosus	Compositae	-	CC	I-II	H	A	Cul-Wid	Gra	2	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Geropogon	hybridus	Compositae	Thanal al Faras	FC	I-II	H	B	Cul	Gra	2	Wee
Gnaphalium	crispatum	Compositae	-	R	I-II	H	A	Wld	Non	0	Wee
Gnaphalium	luteo-album	Compositae	Saboon Afre et	FC	I-II	H	A	Wet	Non	0	Hyg
Gnaphalium	polycaulon	Compositae	-	R	I-II	H	A	Wld	Non	0	Wee
Grangea	maderaspantana	Compositae	-	R	I-II	H	A	Cul-Wld	Non	0	Wee
Gymnarrhena	micrantha	Compositae	Kirsheif	FR	III-IV	H	A	Rge-Sha-Ste	Non	0	Cha-Tac
Hedynois	rhagadioloides	Compositae	-	CC	I-II	H	A	Cul-Wld	Gra	2	Wee
Helichrysum	conglobatum	Compositae	-	RR	I-II	H	P	Cli-Snd	Non	0	Cha-Psa
Helichrysum	orientale	Compositae	-	C	I-II	H	P	Cli-Roc	Non	0	Cha
Hyoseris	radiata	Compositae	-	CC	I-II	H	A	Cul-Wld	Gra	2	Wee
Hyoseris	scabra	Compositae	-	RR	I-II	H	A	Cul-Wld	Gra	2	Wee
Ifloga	labillardieri	Compositae	Krayshif el Gadi	CC	I-IV	H	A	Rge	Gra	2	Ere-Psa
Ifloga	spicata	Compositae	Krayshif el Gadi	CC	I-IV	H	A	Rge-Ste	Gra	0	Ere-Psa
Inula	crithmoides	Compositae	Zeita	C	I	H	P	Coa-Roc	Non	0	Cha
Inula	viscosa	Compositae	Magramam	R	I-II	H	P	Wet-Wld	Med	0	Nit-Pel
Koelpinia	linearis	Compositae	-	C	II-IV	H	A	Rge-Snd-Ste	Gra	3	Psa-Tac
Lactuca	saigna	Compositae	Hawa, Khass el Bagar	R	I-II	H	A-B	Cul	Med	0	Wee
Lasiopogon	muscooides	Compositae	Kreishit el Gadge	FR	I-III	H	A	Wld	Non	0	Nit-Wee
Launaea	angustifolia	Compositae	Noqod	C	III-IV	H	A-B	Rge-Ste	Gra	2	S04
Launaea	capitata	Compositae	Hoadaah	CC	III-IV	H	A	Rge-Snd	Gra	3	Tac
Launaea	nudicaulis	Compositae	Howaa	CC	I-III	H	A	Cul-Rge-Wld	Non	0	Wee
Launaea	resedifolia	Compositae	-	FC	I-III	H	B-P-Sh	Rge-Snd-Ste	Gra	3-4	Psa
Launaea	tenuloba	Compositae	Silen	FR	III-IV	H	A-B	Rge-Ste	Gra	3	Psa
Leontodon	hispidulus	Compositae	-	FC	I-II	H	A	Cul-Wld	Gra	2	Wee
Leontodon	simplex	Compositae	Hawthan	FR	I-II	H	A	Cul-Wld	Non	0	Wee
Leontodon	tuberosus	Compositae	Hawthan	FR	I-II	H	P	Cul-Wld	Gra	2	Wee
Mantisalca	salmantica	Compositae	-	RR	I-II	H	A	Cul-Wld	Non	0	Sil-Wee
Matricaria	aurea	Compositae	Qommeydi	FR	I-II	H	A	Wet	Non	0	Hyg
Matricaria	recutita	Compositae	Baboonig	R	I-II	H	A	Wet	Med-Veg	0	Hyg-Nat
Nauplius	graveolens	Compositae	Rabd	FR	III-VI	H	P	Des-Wad	Non	0	Ere
Notobasis	syriaca	Compositae	Kirsheif	FR	I-II	H	A	Cul-Wld	Non	0	Wee
Onopordon	alexandrinum	Compositae	Stood el Hanash	CC	I-III	H	B	Cul-Wld	Non	0	Psa
Ormenis	mixta	Compositae	-	RR	I-II	H	A	Wet-Wld	Non	0	Hyg
Otanthus	maritimus	Compositae	Shiba	R	I	H	P	Coa-Snd	Med	0	Hal
Phagnalon	barbeyanum	Compositae	Khanaanit	R	I-II	H	P	Cli	Gra	1-2	Cha
Phagnalon	rupestre	Compositae	Taan el Arneb	FC	I-II	H	P	Cli-Ste	Gra	1-2	Cha
Picris	asplenoides	Compositae	Halawaa	CC	I-II	H	A	Cul-Wld	Gra	2	Wee
Picris	cyanocarpa	Compositae	-	RR	I-II	H	A	Cul-Wld	Gra	2	Tac
Picris	sprengeriana	Compositae	Moraar	FC	I-II	H	A	Cul-Wld	Gra	1	Wee
Pluchea	dioscoridis	Compositae	Barr,f	R	I-II	H	P	Wld	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Pulicaria	arabica	Compositae	Abou AOn Safra	FC	I-III	H	A	Wet-Wid	Non	0	Hyg
Pulicaria	crispa	Compositae	Ghobbeira, Aarfeg	FR	III-VI	H	P	Des-Wad	Med	0	Ere
Reichardia	picroides	Compositae	Morrir	FR	I-II	H	P	Cul-Rge	Gra	3	Sil
Reichardia	tingitana	Compositae	Galaween	FC	I-III	H	P	Cul-Rge-Ste	Gra	3	Sil
Rhagadiolus	stellatus	Compositae	-	C	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Rhanterium	suaveolens	Compositae	A'arfeg	R	II-IV	S	P	Rge	Gra	3	Psa
Scolymus	hispanicus	Compositae	-	FR	I-II	H	A-Sh	Cul-Wid	Non	0	Wee
Scorzonera	undulata	Compositae	Dabbah	CC	I-III	H	P	Rge-Ste	Gra-Veg	3	Wee
Senecio	coronopifolius	Compositae	«Moraar; Q, rres»	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Senecio	glaucus	Compositae	Moraar	C	I-II	H	A	Cul-Wid	Non	0	Wee
Senecio	vulgaris	Compositae	Moraar	C	I-II	H	A	Cul-Wid	Non	0	Wee
Silybum	marianum	Compositae	Shook Simmari, Si el Gamel	FR	I-II	H	P	Cul-Wid	Med	0	Nit
Sonchus	asper	Compositae	-	FC	I-II	H	A	Cul-Wid	Non	0	Wee
Sonchus	macrocarpus	Compositae	-	R	I-II	H	A	Cul-Wid	Non	0	Wee
Sonchus	oleraceus	Compositae	Go'odeid, Galawen	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Sonchus	tenerimus	Compositae	-	R	I-II	H	A	Cul-Wid	Non	0	Wee
Spheranthus	suaveolens	Compositae	Habaqbaq	R	I-III	H	P	Rge	Non	0	Wee
Taraxacum	minimum	Compositae	-	RR	I-II	H	P	Cul-Wid	Gra	2	Wee
Urospermum	picroides	Compositae	Galawein, Salis	FC	I-II	H	A	Cul-Wid	Gra	1-2	Pel-Wee
Varthemia	condicans	Compositae	-	FC	I-II	H	P	Cil-Roc-Ste	Med	0	Cha-Pel
Xanthium	spinosum	Compositae	Shobbeit	FR	I-II	H	A	Wid	Non	0	Wee
Xanthium	strumarium	Compositae	Karakh el Bhar, Shabka	R	I-II	H	A	Wid	Non	0	Wee
Xanthium	silvatica	Compositae	-	RR	I-II	Cl-H	P	Cul	Non	0	Wee
Calystegia	althaeoides	Convulvaceae	Maaddah, Luwwaya	CC	I-II	Cl-H	P	Cul	Med	0	Psa-Wee
Convolvulus	arvensis	Convulvaceae	Araaryeh, Ulleq, Lebena	CC	I-II	H	P	Cul	Gra	1	Sil-Wee
Convolvulus	fatmensis	Convulvaceae	-	FR	I-III	H	A	Rge-Wid	Gra	1	Sil
Convolvulus	humilis	Convulvaceae	-	RR	I-II	Cl-H	A	Cul	Non	0	Sil-Wee
Convolvulus	lanatus	Convulvaceae	Bayaad	CC	I-III	Fr	P	Con-Snd	Non	3-4	Psa
Convolvulus	lineatus	Convulvaceae	-	RR	I-II	H	P	Cul	Gra	2	Wee
Convolvulus	oleifolius	Convulvaceae	-	FR	I-II	Fr	P	Cil-Run-Ste	Non	3	Cha
Convolvulus	pilosellifolius	Convulvaceae	-	FR	I-II	H	P	Cul	Gra	2	Wee
Convolvulus	siculus	Convulvaceae	-	FC	I-II	Cl-H	A	Cul	Non	0	Sil-Wee
Convolvulus	stachydifolius	Convulvaceae	-	RR	I-II	H	P	Rge	Gra	2	Psa
Convolvulus	supinus	Convulvaceae	-	FR	I-II	Cl-H	P	Cul	Non	0	Wee
Convolvulus	tricolor	Convulvaceae	-	FR	I-II	Cl-H	P	Cul	Non	0	Wee
Cressa	cretica	Convulvaceae	Molleih, Mellina	CC	I-III	H	A	Sit	Non	0	Hal
Cuscuta	campestris	Convulvaceae	-	RR	I-II	Par	P	Cul-Rge	Non	0	Wee
Cuscuta	epithymum	Convulvaceae	-	FC	I-II	Par	P	Cul-Rge	Non	0	Wee
Cuscuta	europaea	Convulvaceae	-	C	I-II	Par	P	Cul-Rge	Non	0	Wee
Cuscuta	palaestina	Convulvaceae	-	RR	I-II	Par	P	Cul-Rge	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Cuscuta	pedicellata	Convolvulaceae	Hamul	CC	I-II	Par	P	Cul-Rge	Non	0	Wee
Cuscuta	planiflora	Convolvulaceae	Harret al Zaatar	CC	I-II	Par	P	Cul-Rge	Med	0	Wee
Ipomaea	cairica	Convolvulaceae	-	FC	I-II	Ch-H	P	Irr	Non	0	Sum
Ipomaea	imperati	Convolvulaceae	-	R	I-II	Ch-H	P	Irr	Non	0	Sum
Crassula	alata	Crassulaceae	-	RR	II-III	Su	A	Wld	Non	0	Nit
Umbilicus	horizontalis	Crassulaceae	Widna	R	I-II	Su	A	Wld	Non	0	Nit
Alyssum	desertorum	Cruciferae	-	RR	II-III	H	A	Rge	Gra	1-2	Wee
Alyssum	simplex	Cruciferae	-	RR	II-III	H	A	Cul-Rge	Gra	1-2	Wee
Anastatica	hierochuntica	Cruciferae	Keff Maryam, Hidd M.	CC	III-VI	H	A	Des-Ste	Med	0	Ere-Pel-Tac
Biscutella	depressa	Cruciferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Biscutella	didyma	Cruciferae	Sofra	R	I-II	H	A	Cul-Wid	Non	0	Wee
Brassica	nigra	Cruciferae	Libsaan, Khardal Abiad	CC	I-II	H	A	Irr	Med	0	Wee
Brassica	rapa	Cruciferae	-	C	I-II	H	B	Irr	Non	0	Wee
Brassica	tournefortii	Cruciferae	Shiltam	CC	I-IV	H	A	Cul-Snd	Non	0	Psa-Wee
Cakile	maritima	Cruciferae	Figl el Gamel	CC	I	CH	A	Coa-Snd	Non	0	Psa
Camelina	hispida	Cruciferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Camelina	rumelica	Cruciferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Capsella	bursa-pastoris	Cruciferae	Kes el RaO	CC	I-II	H	A	Wld	Med	0	Nit
Carrichtera	annua	Cruciferae	Qineybra	CC	I-III	H	A	Cul-Rge-Ste-Wid	Non	0	Sil-Wee
Conringia	orientalis	Cruciferae	-	RR	II-III	H	A	Cul	Non	0	Wee
Coronopus	squamatus	Cruciferae	Rashaad, Harra	C	I-II	H	A-B	Wet	Non	0	Hyg-Pel
Descurainia	sophia	Cruciferae	-	RR	I-II	H	A	Wld	Non	0	Wee
Didesmus	bipinnatus	Cruciferae	-	C	II-IV	H	A	Cul-Rge	Gra	1	Wee
Diptotaxis	erucoides	Cruciferae	-	R	I-II	H	A-B	Cul-Wid	Non	2	Wee
Diptotaxis	simplex	Cruciferae	-	C	I-II	H	A	Cul	Non	0	Wee
Diptotaxis	viminea	Cruciferae	-	R	I-II	H	A	Cul-Wid	Non	0	Wee
Enarthrocarpus	lyratus	Cruciferae	Shiltam	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Enarthrocarpus	pterocephalus	Cruciferae	-	R	I-III	H	A	Cul-Wid	Non	0	Wee
Enarthrocarpus	strangulatus	Cruciferae	-	C	I-II	H	A	Cul-Wid	Non	0	Wee
Eremobium	aegyptiacum	Cruciferae	Dakhayaan	CC	I-III	H	P	Snd-Ste	Gra	1	Psa
Eruca	sativa	Cruciferae	Gergeer, Gargir	R	I-III	H	A	Cul-Ste-Wid	Gra-Med	2-3	Wee
Erucaria	crassifolia	Cruciferae	Kromb sahra	R	III-VI	H	A	Rge	Gra	2	SO4
Erucaria	hispanica	Cruciferae	Rawq	C	I-II	H	A	Cul	Non	0	Wee
Erucaria	microcarpa	Cruciferae	-	R	II-III	H	A	Cul-Rge	Gra	1	Wee
Erucaria	pinnata	Cruciferae	Saleek	C	I-IV	H	A	Cul-Rge	Gra	1	Wee
Erysimum	repandum	Cruciferae	-	RR	I-II	H	A	Cul-Rge	Non	0	Wee
Farsetia	aegyptiaca	Cruciferae	Garba	CC	II-VI	H	A	Rge	Gra	1-2	Ere
Farsetia	longisiliqua	Cruciferae	Hook	R	III-VI	H	P	Rge	Gra	1-2	Ere
Isatis	lusitanica	Cruciferae	Sofeira	R	I-II	H	A	Cul	Non	0	Wee
Isatis	microcarpa	Cruciferae	-	R	I-II	H	A	Cul	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Lepidium	aucheri	Cruciferae	-	RR	I-II	H	A	Cul-Wid	Non	0	End-Wee
Lepidium	draba	Cruciferae	-	R	I-II	H	A	Cul-Wid	Non	0	Wee
Lepidium	latifolium	Cruciferae	Rashad	R	I-II	H	P	Cul-Wid	Non	0	Wee
Lobularia	arabica	Cruciferae	Dahyaan	C	I-III	H	A	Cul-Snd	Non	0	Psa
Lobularia	libyca	Cruciferae	Khorn el Ibra	C	I-III	H	A	Cul-Snd	Gra	1	Psa
Lobularia	maritima	Cruciferae	Aguerma	FR	I	H	A	Coa-Dne	Med	0	Psa
Malcomia	nana	Cruciferae	-	RR	I-II	H	A	Rge	Gra	2-3	Psa
Malcomia	pygmaea	Cruciferae	Shigara	R	I-II	H	A	Rge-Wld	Gra	2-3	Psa
Matthiola	bicornis	Cruciferae	-	RR	I-III	H	A	Cul	Gra	2	Psa
Matthiola	fruticulosa	Cruciferae	Mantoor	RR	I-II	H	P	Sha-Snd-Ste	Gra	3	Psa
Matthiola	livida	Cruciferae	Shigara	CC	III-VI	H	A	Des	Gra	2	Ere
Matthiola	longipetala	Cruciferae	-	CC	I-III	H	A	Cul-Rge-Ste	Gra	2	Psa
Matthiola	pumilio	Cruciferae	-	CC	I-II	H	A	Cul	Gra	2	Psa
Moricandia	nitens	Cruciferae	H'mim	FC	I-II	Fr	P	Cul-Rge	Fsh-Gra	3-4	Sl-S04
Nasturtiopsis	coronopifolia	Cruciferae	-	R	I-III	H	A	Rge-Wld	Med	0	Sl-S04
Nasturtium	officinale	Cruciferae	Qurrat el Ayn	C	I-II	H	P	Can-Frw	Med-Veg	2-3	Hyd
Neoturularia	torulosa	Cruciferae	-	R	I-II	H	A	Rge-Wld	Non	0	Wee
Ochthodium	aegyptiacum	Cruciferae	-	R	I-II	H	A	Cul	Non	0	Wee
Pseuderucaria	clavata	Cruciferae	-	R	II-III	CH	A	Rge-Ste	Gra	2	S04
Raphanus	raphanistrum	Cruciferae	Figl, Mechtehiya	RR	I-II	H	A	Irr	Non	0	Wee
Rapistrum	rugosum	Cruciferae	-	CC	I-III	H	A	Cul-Wld	Non	0	Wee
Rorippa	palustris	Cruciferae	-	C	I-II	H	A	Can-Frw	Non	0	Hyg
Schouwia	thebaica	Cruciferae	Mahad	R	III-VI	H	A	Des-Wad	Gra	3	Ere
Sinapis	alba	Cruciferae	Khardal	R	I-II	H	A	Irr	Med	0	Wee
Sinapis	arvensis	Cruciferae	Khardal	CC	I-II	H	A	Irr	Med	0	Wee
Sinapis	pubescens	Cruciferae	Khardal	C	I-II	H	A	Cul	Med	0	Wee
Sisymbrium	irio	Cruciferae	Shilyat, Figl el Gamal	CC	I-II	H	A	Cul-Wld	Non	0	Wee
Zilla	spinosa subsp. biparmata	Cruciferae	Zilla, Shebrom	C	III-VI	DS	P	Des-Wad	Gra-Med	3	End-Ere
Bryonia	cretica	Cucurbitaceae	Tatwa, Fashira	R	I-II	Cl	P	Cul-Snd	Med	0	Psa
Citrullus	colocynthis	Cucurbitaceae	Handal	CC	I-IV	Cl	P	Rge-Snd-Ste	Med	0	Psa
Juniperus	phoenicea	Cupressaceae	A'arar	R	I-II	HS-T	P	Roc	Fue-Gra-Med	1	FoRe
Cymodocea	nodosa	Cymodoceaceae	Qoshir	C	I	H	P	Sea	Non	0	Hyd
Cynomorium	coccineum	Cynomoriaceae	Zab et Ard	FC	I-III	Par	P	Slt	Med	0	Hal
Carex	divisa	Cyperaceae	Sard	C	I-II	H	P	Cul-Wet	Non	0	Hyg
Carex	extensa	Cyperaceae	-	R	I-II	H	P	Cul-Wet	Non	0	Hyg
Cyperus	alopecuroides	Cyperaceae	Koosh, Samar	CC	I-II	H	P	Wet	Non	0	Hyg
Cyperus	capitatus	Cyperaceae	S'eed	C	I-II	H	P	Wet	Non	0	Hyg
Cyperus	conglomeratus	Cyperaceae	Oshb	C	II-VI	H	P	Des-Snd	Gra	3	Ere-Psa
Cyperus	difformis	Cyperaceae	Ageera	C	I-II	H	A	Wet	Non	0	Hyg
Cyperus	digitatus	Cyperaceae	Qorreih	C	I-II	H	P	Wet	Non	0	Hyg

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Cyperus	esulentus	Cyperaceae	Habb el Azeez	R	I-II	H	P	Irr-Wet	Med-Veg	0	Hyg
Cyperus	fuscus	Cyperaceae	S'ed	R	I-II	H	P	Wet	Non	0	Hyg
Cyperus	laevigatus	Cyperaceae	Dees, S'eed, Burbet	CC	I-II	H	P	Wet	Non	0	Hyg
Cyperus	rotundus	Cyperaceae	S'ed el Homaar	CC	I-II	H	P	Irr-Wet	Med	0	Hyg
Cyperus	schimperianus	Cyperaceae	S'eed	R	I-II	H	P	Wet	Non	0	Hyg
Eleocharis	geniculata	Cyperaceae	-	R	I-II	H	A-Sh	Wet	Non	0	Hyg
Eleocharis	palustris	Cyperaceae	-	C	I-II	H	P	Wet	Non	0	Hyg
Eleocharis	parvula	Cyperaceae	-	RR	I-II	H	P	Wet	Non	0	Hyg
Fimbristylis	bisumbellata	Cyperaceae	Forrysh, Shaara	CC	I-II	H	A	Wet	Non	0	Hyg
Fimbristylis	pubescens	Cyperaceae	Shaara	R	I-II	H	P	Wet	Non	0	Hyg
Schoenus	nigricans	Cyperaceae	-	R	I-II	H	P	Sit-Wet	Non	0	Hal-Hyg
Scirpus	articulatus	Cyperaceae	Smaar	RR	I-II	H	A	Wet	Non	0	Hyg
Scirpus	holoschenus	Cyperaceae	Dees	R	I-II	H	P	Sit-Wet	Non	0	Hal-Hyg
Scirpus	inclinatus	Cyperaceae	Samaar	R	I-II	H	P	Wet	Non	0	Hyg
Scirpus	litoralis	Cyperaceae	Khabb	CC	I-II	H	P	Wet	Non	0	Hyg
Scirpus	maritimus	Cyperaceae	Smaar	CC	I-II	H	P	Sit-Wet	Non	0	Hal-Hyg
Scirpus	praelongus	Cyperaceae	Dees	C	I-II	H	P	Wet	Non	0	Hyg
Scirpus	supinus	Cyperaceae	Khabb	R	I-II	H	A	Wet	Non	0	Hyg
Cephalaria	syriaca	Dipsacaceae	Siwaan	R	I-II	H	A	Rge-Wld	Gra	1	Wee
Pteroccephalus	plumosus	Dipsacaceae	Sofeira	FR	I-III	H	A	Rge	Gra	1	Tac-Wee
Scabiosa	eremophila	Dipsacaceae	Kaahel Ghazal	R	I-III	H	A	Rge	Gra	1	Wee
Scabiosa	rhizantha	Dipsacaceae	Siwaan	R	I-II	H	A	Rge	Gra	1	Wee
Ephedra	alata	Ephedraceae	Ald, Alendra	R	III-IV	HS	P	Des-Snd	Gra-Med	1-2	Ere
Ephedra	aphylla	Ephedraceae	Algarn, Maad	FC	I-II	HS	P	Cil-Run-Wad	Gra-Med	3	Cha
Chrozophora	tinctoria	Euphorbiaceae	Ghobbeira	FC	I-III	H	A	Wld	Non	T	Sum
Euphorbia	arguta	Euphorbiaceae	Libbein	C	II-III	H	A	Rge-Wld	Non	T	Wee
Euphorbia	bivonae	Euphorbiaceae	Libbein	R	I-II	S	P	Roc	Non	T	Cha
Euphorbia	bivonae	Euphorbiaceae	Libbein	FR	I-II	DS	P	Cil	Non	T	Cha
Euphorbia	chamaepeplus	Euphorbiaceae	Bena	RR	I-II	H	A	Cul-Wld	Non	T	Wee
Euphorbia	dendroides	Euphorbiaceae	Libbein	R	I-II	DS	P	Cil-Roc	Non	T	Cha
Euphorbia	falcata	Euphorbiaceae	Libbein	RR	I-II	H	A	Cul-Wld	Non	T	Wee
Euphorbia	forskallii	Euphorbiaceae	Shaarel Agouz	C	I-II	H	A	Cul-Wld	Non	T	Wee
Euphorbia	helioscopia	Euphorbiaceae	Libbein, Saada	C	I-II	H	A	Cul-Wld	Med	T	Nit-Wee
Euphorbia	hirta	Euphorbiaceae	Libbein	C	I-II	H	A	Wld	Non	T	Wee
Euphorbia	mauritanica	Euphorbiaceae	Libbein	R	I-II	H	P	Wld	Non	T	Wee
Euphorbia	paralias	Euphorbiaceae	Shagaret el Hanash	C	I	Fr	P	Coa-Snd	Non	T	Psa
Euphorbia	parvula	Euphorbiaceae	Libbein	RR	I-II	H	A	Wld	Non	T	Wee
Euphorbia	peplis	Euphorbiaceae	Libbein	R	I-II	H	A	Cul-Wld	Non	T	Wee
Euphorbia	peplus	Euphorbiaceae	Widenia	CC	I-II	H	A	Cul-Wld	Med	T	Wee
Euphorbia	prostrata	Euphorbiaceae	Libbein	C	I-II	H	A	Cul-Wld	Non	T	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Euphorbia	pterocecca	Euphorbiaceae	Libbein	RR	I-II	H	A	Cul-Wid	Non	T	Wee
Euphorbia	punctata	Euphorbiaceae	Libbein	RR	I-II	H	A	Cul-Wid	Non	T	Wee
Euphorbia	retusa	Euphorbiaceae	No'manya	C	III-VI	Fr	P	Des-Rge-Ste	Non	T	Ere
Euphorbia	terraccina	Euphorbiaceae	Libbein	C	I-III	Fr-H	P	Cul-Ste-Wid	Non	T	Wee
Haplophyllum	tuberculatum	Euphorbiaceae	Shag, el Kelb, F, el	CC	I-III	Fr	P	Cul-Wid	Med	T	Sum
Mercurialis	annua	Euphorbiaceae	Halboob	R	I-II	H	A-B	Cil-Run-Wid	Med	T	Nit
Phyllanthus	rotundifolius	Euphorbiaceae	Tarlanoi hindi	FC	I-II	H	A	Cul-Wid	Non	0	Wee
Ricinus	communis	Euphorbiaceae	Kreroua, Kherwa	FR	I-II	TS	P	Snd-Wad-Wid	Med	T	Nat-Psa
Frankenia	hirsuta	Frankeniaceae	Molleih	R	I-IV	H	P	Sit	Non	0	Hal
Frankenia	pulverulenta	Frankeniaceae	Molleih	C	I-IV	H	A	Sit	Non	0	Hal
Frankenia	revoluta	Frankeniaceae	-	RR	III-VI	Fr	P	Sit	Non	0	Hal
Fumaria	bracteosa	Fumariaceae	-	C	I-II	H	A	Cul	Non	0	Wee
Fumaria	capreolata	Fumariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Fumaria	densiflora	Fumariaceae	Zeita	CC	I-II	H	A	Cul	Non	0	Wee
Fumaria	gaillardotii	Fumariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Fumaria	judaica	Fumariaceae	-	R	I-II	H	A	Cul	Med	0	Wee
Fumaria	microstachys	Fumariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Fumaria	officinalis	Fumariaceae	-	RR	I-II	H	A	Cul	Med	0	Wee
Fumaria	parviflora	Fumariaceae	Shahatrag, H. al Sabyan	CC	I-II	H	A	Cul	Med	0	Wee
Hypecoum	aegyptiacum	Fumariaceae	-	C	I-II	H	A	Des-Rge	Gra	0	Psa-Tac
Hypecoum	aequilobum	Fumariaceae	-	R	I-III	H	A	Cul-Rge	Gra	0	Psa-Tac
Hypecoum	littorale	Fumariaceae	-	RR	I	H	A	Coa-Snd	Non	0	Psa
Hypecoum	pendulum	Fumariaceae	Sateek	RR	I-III	H	A	Cul-Rge-Ste	Non	2	Psa-Tac
Hypecoum	procumbens	Fumariaceae	-	RR	I-II	H	A	Cul-Rge	Non	0	Psa-Tac
Centaureum	maritimum	Gentianaceae	-	RR	I-II	H	A	Sit	Non	0	Hal-Psa
Centaureum	pulchellum	Gentianaceae	Kantarion	CC	I-III	H	A	Sit	Med	0	Hal-Hyg
Centaureum	spicatum	Gentianaceae	Menash el Dibban	C	I-II	H	A	Sit	Med	0	Hal-Hyg
Erodium	arborescens	Geraniaceae	Dahma	R	I-III	CH	P	Rge-Ste-Wid	Non	0	S04
Erodium	chium	Geraniaceae	Ghazlan	R	I-III	H	A	Cul-Rge	Gra	2	Wee
Erodium	cicutium	Geraniaceae	Abu Mosfa	R	I-II	H	A	Cul-Rge	Gra	2	Wee
Erodium	cicutarium	Geraniaceae	Dahmyea abu Ghazal	FC	I-II	H	A	Cul-Rge	Gra-Med	2	Wee
Erodium	crassifolium	Geraniaceae	-	C	I-II	H	P	Cul-Rge-Ste	Gra	1	Nit
Erodium	glaucoophyllum	Geraniaceae	Timmeir	CC	I-III	CH	P	Rge-Ste-Wid	Non	0	Sil-S04
Erodium	gruinum	Geraniaceae	Abu Mosfa	R	I-II	H	A	Cul-Rge	Gra	2	Wee
Erodium	laciniatum	Geraniaceae	Moryhaal	CC	I-III	H	A	Cul-Rge	Gra	2-3	Psa
Erodium	malacoides	Geraniaceae	Oqqeil	CC	I-II	H	A	Cul-Rge-Ste	Gra	2-3	Sil
Erodium	moschatum	Geraniaceae	Meseika	R	I-II	H	A	Cul-Rge	Gra	2	Wee
Erodium	neuradifolium	Geraniaceae	-	C	I-II	H	A	Cul-Rge	Gra	2-3	Psa
Erodium	oxyrhynchum	Geraniaceae	-	C	I-II	H	P	Cul-Rge-Ste	Gra	1	Wee
Geranium	dissectum	Geraniaceae	-	R	I-II	H	A	Cul	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Geranium	molle	Geraniaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Globularia	alypum	Globulariaceae	Zrega	FR	I-II	H	P	Cil-Roc-Ste	Gra-Med	2	Cha
Globularia	arabica	Globulariaceae	Zrega	FC	I-II	H	P	Cil-Roc-Ste	Gra-Med	2	Cha
Aegilops	bicornis	Gramineae	Shaer el Ides	CC	I-II	H	A	Cul-Rge-Wid	Gra	1	Wee
Aegilops	geniculata	Gramineae	-	FR	I-II	H	A	Cul-Rge-Wid	Gra	1	Wee
Aegilops	kotschy	Gramineae	Shaer el Fahr	C	I-II	H	A	Cul-Rge-Wid	Gra	1	Wee
Aegilops	longissima	Gramineae	-	R	I-II	H	A	Cul-Rge-Wid	Gra	1	Wee
Aegilops	pergrina	Gramineae	-	RR	I-II	H	A	Cul-Rge-Wid	Gra	1	Wee
Aegilops	ventricosa	Gramineae	Shaer el Fahr	RR	I-II	H	A	Cul-Rge-Wid	Gra	2	Wee
Aeluropus	lagopoides	Gramineae	Akreesh	CC	I-III	H	P	Sit	Gra	1	Hal
Agropyron	crisatum	Gramineae	-	R	I-II	H	P	Cul	Gra	3	Wee
Alopecurus	myosuroides	Gramineae	-	R	I-II	H	P	Wet	Gra	2	Hyg
Ammochloa	palaestina	Gramineae	-	FC	I-II	H	A	Cul-Rge	Non	0	Hal
Ammophila	arenaria	Gramineae	Gazzoof	CC	I	H	P	Coa-Snd	Non	0	Hal-Psa
Aristida	funiculata	Gramineae	Qaw	R	II-IV	H	A	Rge-Snd	Gra	2	Psa
Arundo	donax	Gramineae	Ghaab, B,s	CC	I-III	H	P	Wet	Han-Wbr	2	Hyg
Avena	barbata	Gramineae	Khafoor	CC	I-III	H	A	Cul-Wid	Gra	2	Wee
Avena	fatua	Gramineae	Khafoor, Zummayr	FC	I-II	H	A	Cul-Wid	Gra	2	Wee
Avena	sterilis	Gramineae	Khafoor	FC	I-II	H	A	Cul-Wid	Gra	2	Wee
Brachiaria	mutica	Gramineae	Moddeid	R	I-II	H	P	Irr	Gra	5	Fod-Hyg-Sum
Brachiaria	repens	Gramineae	Nissela	R	I-II	H	P	Wet	Gra	2-3	Hyg-Sum
Brachypodium	distachyum	Gramineae	Shaer Barri	CC	I-III	H	A	Rge-Ste-Wid	Non	1	Wee
Briza	maxima	Gramineae	Sahliah	R	I-II	H	A	-	Gra	1-2	Wee
Briza	minor	Gramineae	Sahliah	RR	I-II	H	A	Wid	Gra	1-2	Wee
Bromus	aegyptiacus	Gramineae	Khafoor	-	I-II	H	A	Cul-Wid	Gra	2	Wee
Bromus	diandrus	Gramineae	Saf Soof	R	I-II	H	A	Cul-Wid	Gra	2	Wee
Bromus	fasciculatus	Gramineae	-	CC	I-III	H	A	Cul-Rge-Wid	Gra	2	Wee
Bromus	japonicus	Gramineae	Saf Soof	RR	I-II	H	A	Cul-Wid	Gra	2	Wee
Bromus	madritensis	Gramineae	-	R	I-II	H	A	Cul-Wid	Gra	2	Wee
Bromus	rubens	Gramineae	-	C	I-III	H	A	Cul-Rge-Wid	Gra	2	Wee
Bromus	scoparius	Gramineae	Sabal el Fahr	C	I-II	H	A	Cul-Wid	Gra	2	Wee
Bromus	tectorum	Gramineae	Sabal abu Hossein	R	I-II	H	A	Cul-Wid	Gra	2	Wee
Castalia	tuberculata	Gramineae	-	R	I-II	H	A	Rge	Gra	2	Psa
Catapodium	rigidum	Gramineae	-	RR	I-II	H	A	Cul-Wid	Gra	1	Psa
Cenchrus	ciliaris	Gramineae	Hada	RR	I-III	H	P	Cil-Run	Gra	5	Fod-Psa
Centropodia	forskallii	Gramineae	Migel en Naga	FR	II-IV	H	P	Rge-Snd	Gra	2	Ere-Psa
Corynephorus	divaricatus	Gramineae	-	FR	I-II	H	A	Coa-Snd	Non	0	Psa
Crithopsis	delleana	Gramineae	-	RR	I-II	H	A	Rge-Ste-Wid	Gra	1	Wee
Crypsis	aculeata	Gramineae	-	R	I-II	H	A	Wet	Non	0	Hyg
Crypsis	alopecuroides	Gramineae	-	R	I-II	H	A	Wet	Non	0	Hyg

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Crypsis	schoenoides	Gramineae	-	R	I-II	H	A	Wet	Non	0	Hyg
Cutandia	dichotoma	Gramineae	Sammah	CC	I-III	H	A	Con-Snd-Ste	Gra	2	Psa-Tac
Cutandia	maritima	Gramineae	Sammah	RR	I	H	A	Coa-Wid	Gra	1	Psa
Cutandia	memphitica	Gramineae	Sammah	CC	I-III	H	A	Coa-Snd-Ste	Gra	2	Psa-Tac
Cynodon	dactylon	Gramineae	Nigel, Negem	CC	I-III	H	P	Cul	Gra-Med	5	Fod-Psa-Wee
Cynosurus	coloratus	Gramineae	Sahlah	R	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Cynosurus	echinatus	Gramineae	Sahlah	C	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Dactylis	hispanica	Gramineae	-	C	I-II	H	A	Cli-Run	Gra	4	Cha
Dactyloctenium	aegyptium	Gramineae	Na'eem el Saleeb	CC	I-II	H	A	Cul-Wid	Gra	2	Psa-Sum
Desmaziera	philistaea subsp. rhofisiana	Gramineae	Khafoor	RR	I-II	H	A	Rge-Snd	Gra	2	Psa
Dichanthium	annulatum	Gramineae	Abu Qoseiba, Himmara	RR	I-II	H	P	Cli-Run-Wad	Gra	3	Sum
Digitaria	sanguinalis	Gramineae	Dafira	R	I-II	H	A	Cul-Wet	Gra	1	Hyg-Sum
Dinebra	retroflexa	Gramineae	Deneeb, Nigi al Nimr	C	I-II	H	A	Irr	Gra	1	Hyg-Wee
Echinochloa	colona	Gramineae	Moddeid, Abu Rukba	CC	I-II	H	A	Irr	Gra	2	Hyg-Sum
Echinochloa	crus-galli	Gramineae	Deneiba	CC	I-II	H	A	Irr	Gra	2	Sum-Wee
Echinochloa	stagnina	Gramineae	Omshaat	RR	I-II	H	P	Can	Gra	4-5	Hyd-Sum
Elymus	elongatus	Gramineae	Gazzoof	CC	I	H	P	Coa-Snd	Gra	0	Psa
Elymus	farctus	Gramineae	Gazzoof	CC	I	H	P	Coa-Snd	Non	0	Psa
Elytrigia	cristata	Gramineae	-	FR	I	H	P	Coa-Dne	Gra	1	Psa
Elytrigia	juncea	Gramineae	-	FR	I	H	P	Coa-Dne	Gra	1	Psa
Eragrostis	aegyptiaca	Gramineae	Qaw	C	I-II	H	A	Cul-Snd	Gra	1	Psa-Wee
Eragrostis	barrelieri	Gramineae	-	R	I-II	H	A	Cul-Snd	Gra	1	Psa-Wee
Eragrostis	cilianensis	Gramineae	Ti raab	C	I-II	H	A	Wid	Gra	1	Wee
Eragrostis	pilosa	Gramineae	Heelagoog	C	I-II	H	A	Wet	Gra	1	Hyg-Wee
Eragrostis	phleoides	Gramineae	-	R	I-II	H	A	Wid	Non	0	Wee
Gastridium	altissima	Gramineae	Rodeybe	R	I-II	H	P	Can-Wet	Gra	2	Hyd-Hyg
Hordeum	marinum	Gramineae	Reish Hosny	CC	I-II	H	A	Sit	Gra	1	Hal
Hordeum	murinum	Gramineae	Reesh	CC	I-III	H	A	Cul-Wid	Gra	2	Nit
Hordeum	spontaneum	Gramineae	-	RR	I-III	H	A	Cul-Wad	Gra	2	Wee
Hyparrhenia	hirta	Gramineae	Saf Soof	FR	I-II	H	P	Run-Ste-Wad	Gra	3	Sum
Imperata	cylindrica	Gramineae	Halfa, Del al Qutt	CC	I-II	H	P	Cul-Wet	Gra-Med	1	Hyg-Wee
Lagurus	ovatus	Gramineae	-	C	I-II	H	A	Snd-Wid	Non	0	Psa
Lamarkia	aurea	Gramineae	Sahlah	FC	I-II	H	A	Wid	Non	0	Nit
Lasiurus	scindicus	Gramineae	Hada	RR	III-VI	H	P	Des-Dne-Ste	Gra	3	Ere-Fod-Psa
Leersia	hexandra	Gramineae	Sholleik	R	I-II	H	P	Wet	Gra	3	Hyg-Hyg
Leptochloa	fusca	Gramineae	Sey foon	FC	I-II	H	P	Wet	Gra	2	Hyg-Pel
Lolium	multiflorum	Gramineae	Simbil, Samma	FC	I-II	H	A-B	Cul-Irr	Gra	5	Wee
Lolium	perenne	Gramineae	Gazoon, Hashish al Faras	R	I-II	H	P-Sh	Cul-Irr	Gra-Med	5	Wee
Lolium	rigidum	Gramineae	Sahlah	C	I-II	H	A	Cul-Rge-Ste-Wid	Gra	3	Psa
Lolium	temulentum	Gramineae	Zawaan	R	I-II	H	A-B-Sh	Cul-Irr	Gra-Med	4	Fod-Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Lygeum	spartum	Gramineae	Halfa Mahbouba	CC	I-III	H	P	Rge-Sha-Ste	Han	1	S04
Melica	minuta	Gramineae	-	RR	I-II	H	P	Rge	Gra	3	Psa
Oryza	sativa	Gramineae	Rooz	R	I-II	H	P	Can-Frw	Cro-Gra	3	Hyd
Oryzopsis	miliacea	Gramineae	Himaar/Hemri	FC	I-II	H	P	Cul-Run	Gra	4	Cha
Panicum	antidotale	Gramineae	-	RR	I-III	H	P	Wad	Gra	4-5	Foc-Psa
Panicum	coloratum	Gramineae	Qasaba	FC	I-II	H	P	Irr	Gra	4	Foc-Sum-Wee
Panicum	repens	Gramineae	Nigel, Zummar	FR	I-II	H	P	Cul-Wet	Gra	2-3	Hyg
Panicum	turgidum	Gramineae	Abu Rokba	C	III-VI	HS	P	Des-Ste-Wad	Gra-Med-Sbi	2	Ere
Parapholis	filiformis	Gramineae	-	FC	I-II	H	A	Sit	Non	0	Hal
Parapholis	incurva	Gramineae	-	R	I-II	H	A	Sit	Non	0	Psa
Parapholis	marginata	Gramineae	-	R	I-II	H	A	Sit	Non	0	Hal
Paspalidium	geminatum	Gramineae	Sayfoon, Nisela	C	I-II	H	P	Wet	Gra	2	Hyg-Sum
Paspalidium	obtusifolium	Gramineae	Moddeid	R	I-II	H	P	Wet	Gra	2	Hyg-Sum
Paspalum	distichum	Gramineae	Mudded	R	I-II	H	P	Wld	Gra	4	Foc-Hyg-Sum
Pennisetum	divisum	Gramineae	Tommar	RR	III-VI	HS	P	Des-Ste-Wad	Gra	2	Ere-Psa
Pennisetum	setaceum	Gramineae	Sabat	R	II-IV	H	P	Run-Ste-Wad	Gra	1	Cha-Psa
Phalaris	canariensis	Gramineae	-	C	I-II	H	A	Cul	Gra	3	Sil
Phalaris	minor	Gramineae	Shaer el Fahr	CC	I-III	H	A	Cul-Rge	Gra	2	Sil
Phalaris	paradoxa	Gramineae	-	CC	I-III	H	A	Cul-Rge	Gra	1	Pel
Phragmites	australis	Gramineae	Qasab Hagana, Ghab	CC	I-III	H	P	Sit	Han-Med-Wbr	1	Hal-Hyg
Poa	annua	Gramineae	-	FC	I-II	H	A	Cul-Wld	Gra	1	Nit-Wee
Poa	infirmia	Gramineae	Sahlah	FC	I-II	H	A	Cul	Gra	1-2	Nit-Wee
Polygonum	monspeliensis	Gramineae	Deil el Dott	C	I-II	H	A	Sit	Non	0	Hal
Polygonum	viridis	Gramineae	Deir el Fahr	C	I-II	H	P	Sit	Non	0	Hal
Rostraria	cristata	Gramineae	-	C	I-II	H	A	Cul-Rge	Gra	2	Sil
Rostraria	hispidia	Gramineae	-	CC	I-III	H	A	Cul-Rge-Ste	Gra	2	Psa
Rostraria	pumila	Gramineae	-	CC	I-III	H	A	Cul-Rge	Gra	2	Psa
Saccharum	spontaneum	Gramineae	Boos	C	I-II	H	P	Cul-Wet	Gra-Sbi-Wbr	2	Hyg
Schismus	arabicus	Gramineae	Baima	CC	I-II	H	A	Rge-Snd-Ste	Gra	3	Psa
Schismus	barbatus	Gramineae	Baima	CC	I-II	H	A	Rge-Snd-Ste	Gra	3	Psa
Setaria	verticillata	Gramineae	Yadaab, Qamb al Far	CC	I-II	H	A	Cul-Wld	Gra	1	Sum-Wee
Setaria	verticilliformis	Gramineae	Del el Far	R	I-II	H	A	Cul-Wld	Gra	1	Sum-Wee
Sorghum	halapense	Gramineae	Djarrapp	R	I-II	H	P	Cul-Wet	Gra	3	Hyg-Wee
Sorghum	virgatum	Gramineae	Garawa	R	I-II	H	A	Wet	Non	0	Hyg-Wee
Sphenopus	divaricatus	Gramineae	-	CC	I-II	H	A	Sit	Non	0	Hal
Sporobolus	pungens	Gramineae	Shakaam	C	I	H	P	Coa-Snd	Gra	0	Hal
Sporobolus	spicatus	Gramineae	Sammah	C	I-II	H	P	Sit	Gra	2	Hal
Stipa	capensis	Gramineae	Behima/Sabat	CC	I-III	H	A	Rge-Sha-Ste	Gra	1	Sil
Stipa	lagascae	Gramineae	Gawther	FR	I-III	H	P	Rge-Snd-Ste	Gra	3-4	Psa
Stipa	parviflora	Gramineae	Safssoof	CC	I-III	H	P	Roc-Sha-Ste	Gra	3	Cha

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Stipagrostis	ciliata	Gramineae	Assahn	FC	I-III	H	P	Rge-Ste	Gra	2	Sil-SO4
Stipagrostis	lanata	Gramineae	Shaf Shoof	R	II-III	H	P	Dne-Rge	Gra	2	Psa
Stipagrostis	obtusata	Gramineae	Saf Soof	FR	II-IV	H	P	Rge-Snd-Ste	Gra	2	Psa
Stipagrostis	plumosa	Gramineae	Nawa Beida	FR	II-IV	H	P	Rge-Snd-Ste	Gra	2	Psa
Stipagrostis	scoparia	Gramineae	Sabat	FR	II-IV	H	P	Dne-Rge	Gra-Med-Sbi	1	Psa
Taeniatherum	caput-medusae	Gramineae	Salsoof	RR	I-II	H	A	Cul-Wid	Gra	1	Wee
Triplachne	nitens	Gramineae	-	R	I-II	H	A	Coa-Dne-Sit	Non	0	Hal-Psa
Trisetaria	macrochaeta	Gramineae	-	FR	I-II	H	A	Cul-Rge	Gra	2	Wee
Vulpia	brevis	Gramineae	Sahliah	R	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Vulpia	bromoides	Gramineae	Sahliah	RR	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Vulpia	fasciculata	Gramineae	Sahliah	R	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Vulpia	myuros	Gramineae	Sahliah	RR	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Vulpia	peccinella	Gramineae	Sahliah	R	I-II	H	A	Cul-Wid	Gra	1-2	Wee
Halophila	stipulacea	Hydrocharitaceae	-	RR	I-II	H	P	Can-Frw	Non	0	Hyd
Najas	graminea	Hydrocharitaceae	-	R	I-II	H	A	Can-Frw	Non	0	Hyd
Najas	marina	Hydrocharitaceae	Horreish	FC	I-II	H	A	Bra-Can	Non	0	Hyd
Najas	minor	Hydrocharitaceae	-	R	I-II	H	A	Can-Frw	Non	0	Hyd
Ottelia	alismoides	Hydrocharitaceae	Widna	FR	I-II	H	A	Can-Frw	Non	0	Hyd
Gladiolus	italicus	Iridaceae	Seil el Ghorab	R	I-III	H	P	Cul-Wid	Orn	0	Wee
Gymandris	monophylla	Iridaceae	Kheita	C	I-III	H	P	Rge-Wid	Non	0	Wee
Gymandris	syrinchium	Iridaceae	Kheita	CC	I-III	H	P	Rge-Ste-Wid	Non	0	Wee
Juncus	acutus	Juncaceae	Samaar/mor	CC	I-III	H	P	Sit-Wet	Han-Med	0	Hal-Hyg
Juncus	bufonius	Juncaceae	Shaar el Kird	C	I-II	H	A	Wet-Wid	Non	0	Hyg
Juncus	fontanesii	Juncaceae	-	R	I-II	H	P	Wet-Wid	Non	0	Hyg
Juncus	hybridus	Juncaceae	-	R	I-II	H	A	Wet-Wid	Non	0	Hyg
Juncus	rigidus	Juncaceae	Samaar	CC	I-III	H	P	Sit-Wet	Non	0	Hal-Hyg
Juncus	subulatus	Juncaceae	Haleyn	C	I-II	H	P	Wet-Wid	Han	0	Hyg
Ajuga	iva	Labiatae	Shandagoara	R	I-II	H	P	Cil-Run	Med	0	Cha
Ballota	damascena	Labiatae	Asaghan	FC	I-II	H	P	Wid	Non	0	Nit
Ballota	pseudo-dictamnus	Labiatae	-	FR	I-II	Fr	P	Wid	Non	0	Wee
Lamium	amplexicaule	Labiatae	Taqiyt el Ghorah	CC	I-II	H	A	Cul	Non	0	Wee
Marrubium	alysson	Labiatae	Frsiyoun, Marriout	CC	I-III	H	P	Wid	Non	0	Nit
Marrubium	vulgare	Labiatae	Robeia	C	I-II	H	P	Wid	Med	0	Nit
Phlomis	fioccosa	Labiatae	Zeheira	FC	I-II	HS	P	Cil-Run	Non	0	Cha
Prasium	majus	Labiatae	Shoofal	FC	I-II	Fr	P	Cil-Run	Non	1	Cha
Salvia	aegyptiaca	Labiatae	Raahl	FC	I-III	Fr	P	Sna-Ste	Gra	1	Sil
Salvia	lanigera	Labiatae	Nameia	FC	I-III	H	P	Rge-Ste	Gra	3	Sil
Salvia	spinosa	Labiatae	Marymya	R	I-II	H	A	Wid	Non	0	Sil
Salvia	verbenaca	Labiatae	Thalaba	FC	I-III	H	P	Rge-Ste	Gra	3	Psa
Satureja	nervosa	Labiatae	-	FC	I-II	Fr	P	Cil-Roc	Gra	2	Cha

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Teucrium	brevifolium	Labiatae	-	FR	I-III	Fr	P	Rge	Gra	1	Cha-End
Teucrium	pollum	Labiatae	Ga'da	CC	I-III	Fr	P	Rge-Sha-Ste	Med	0	Cha
Thymus	capitatus	Labiatae	Zaater	R	I-II	DS	P	Cil-Roc-Ste	Med-Veg	1	Cha
Acacia	ehrenbergiana	Leguminosae	Seyal	RR	III-VI	HS-T	P	Des-Wad	Gra-Med	1	Ere-Phr
Acacia	raddiana	Leguminosae	Talh	R	III-VI	HS-T	P	Des-Wad	Gra-Med	3	Ere-Phr
Alhagi	graecorum	Leguminosae	Agool, Aq, I	FC	III-VI	Fr	P	Sit-Wld	Gra-Med	1	Hal-Phr
Anagyris	foetida	Leguminosae	Garood, Karroub el Khinzir	RR	I-II	HS	P	Cul-Rge	Med	T	Sil
Anthyllis	vulneraria	Leguminosae	-	RR	I-II	H	A-P	Cul-Rge-Ste	Gra-Med	2-3	Wee
Argyrobobium	uniflorum	Leguminosae	-	FR	I-VI	DS	P	Rge-Snd-Ste	Gra	4	Psa
Astragalus	asterias	Leguminosae	Abu Egeifa	R	I-II	H	A	Cul-Rge	Gra	1-2	Psa
Astragalus	boeoticus	Leguminosae	Mehallaq	C	I-II	H	A	Cul-Wld	Non	0	Wee
Astragalus	caprinus	Leguminosae	-	C	I-III	H	P	Rge-Ste-Wld	Non	0	Psa
Astragalus	fruticosus	Leguminosae	-	C	I-III	Fr	P	Rge	Non	0	Ere
Astragalus	hamosus	Leguminosae	Qorein	C	I-II	H	A	Cul-Wld	Non	0	Sil-Wee
Astragalus	hauarensis	Leguminosae	Danel Farah	R	III-VI	H	A	Rge-Snd	Gra	3-4	Psa-Tac
Astragalus	hispidulus	Leguminosae	-	R	III-IV	H	A	Des-Rge	Gra	2	Psa-Tac
Astragalus	mareoticus	Leguminosae	-	R	I-III	H	A	Rge	Gra	2	Psa
Astragalus	spinosus	Leguminosae	-	CC	III-VI	DS	P	Rge-Wad	Gra	1	Ere
Astragalus	tribuloides	Leguminosae	Bel et Gamel	CC	I-II	H	A	Cul-Rge	Non	0	Psa-Tac-Wee
Astragalus	trigonus	Leguminosae	-	FC	III-VI	DS	P	Des-Wad	Gra	1	Ere
Ebenus	armitagei	Leguminosae	-	RR	I	Fr	P	Rge-Roc	Gra	2	Cha-End
Hedysarum	coronarium	Leguminosae	-	RR	I-II	H	A-B	Irr	Gra	5	Fod-Pel
Hedysarum	spinosissimum	Leguminosae	-	R	I-III	H	A	Rge-Snd-Ste	Gra	3	Psa
Hippocrepis	areolata	Leguminosae	-	CC	I-II	H	A	Rge-Ste	Gra	3	Psa
Hippocrepis	cyclocarpa	Leguminosae	-	CC	I-II	H	A	Cul-Rge	Gra	2	Sil
Hippocrepis	unisiliquosa	Leguminosae	Omm Danara	RR	I-III	H	A	Rge-Ste	Gra	3	Psa
Hymenocarpus	circinnatus	Leguminosae	-	R	I-II	H	A	Cul-Rge-Ste	Gra	1	Wee
Lathyrus	aphaca	Leguminosae	Hamaam al Burg	C	I-II	H	A	Cul	Gra	3	Wee
Lathyrus	hirsutus	Leguminosae	Hada, Duhreg	C	I-II	H	A	Cul	Gra	2	Wee
Lathyrus	marmoratus	Leguminosae	-	CC	I-II	H	A	Cul	Gra	2	Wee
Lathyrus	pseudocicera	Leguminosae	-	FC	I-II	H	A	Cul	Gra	2	Wee
Lathyrus	sativus	Leguminosae	-	C	I-II	H	A	Cul	Gra	2	Wee
Lathyrus	setifolius	Leguminosae	-	RR	I-II	H	A	Cul	Gra	2	Wee
Lotus	corniculatus	Leguminosae	Rigl al Asfur	RR	I-II	H	P	Cul-Wet	Gra	5	Hyg
Lotus	creticus	Leguminosae	Eshb	C	I-II	H	P	Rge-Snd-Ste	Gra	4	Psa
Lotus	cyrtioides	Leguminosae	Eshb	RR	I-II	H	P	Coa-Rge-Snd	Gra	3	Hal-Psa
Lotus	edulis	Leguminosae	-	R	I-II	H	P	Cul	Gra	1	Wee
Lotus	halophilus	Leguminosae	Herbith	CC	I-III	H	A	Cul-Rge-Ste	Gra	1-2	Psa
Lotus	ornithopodioides	Leguminosae	Ommelkheishat	R	I-II	H	A	Cul	Gra	1	Wee
Lotus	palustris	Leguminosae	-	RR	I-II	H	P	Cul-Wet	Gra	3	Hyg

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Lotus	pedunculatus	Leguminosae	-	RR	I-II	H	P	Cul-Wet	Gra	4	Hal-Hyg
Lotus	polyphyllus	Leguminosae	-	C	I	H	P	Coa-Dne	Gra	0	Hal-Psa
Lotus	tetragonolobus	Leguminosae	Asai el Arous	C	I-II	H	P	Cul-Wet	Gra	4	Hal-Hyg
Medicago	arabica	Leguminosae	-	R	I-II	H	A	Cul	Gra	2	Wee
Medicago	coronata	Leguminosae	-	R	I-II	H	A	Cul-Wild	Gra	2-3	Wee
Medicago	hypogea	Leguminosae	Adreis	RR	I-II	H	A	Cul-Rge-Ste	Gra	2	Wee
Medicago	intertexta	Leguminosae	Khasag	RR	I-II	H	A	Cul-Rge	Gra	2-3	Wee
Medicago	lacinata	Leguminosae	-	FC	I-IV	H	A	Cul-Rge-Ste	Gra	2-3	Psa-Tac
Medicago	littoralis	Leguminosae	-	CC	I-III	H	A	Cul-Rge-Ste	Gra	2-3	Psa
Medicago	lupulina	Leguminosae	-	R	I-II	H	A-B-Sh	Cul-Wild	Gra	2-3	Wee
Medicago	marina	Leguminosae	-	C	I	H	P	Coa-Snd	Non	0	Hal
Medicago	minima	Leguminosae	-	FR	I-IV	H	A	Rge-Ste	Gra	1	Psa
Medicago	monspeliaca	Leguminosae	-	R	I-II	H	A	Cul-Rge	Non	0	Hal
Medicago	orbicularis	Leguminosae	-	R	I-II	H	A	Cul	Gra	3	Wee
Medicago	polycreata	Leguminosae	-	R	I-II	H	A	Cul-Rge	Non	0	Hal
Medicago	polymorpha	Leguminosae	Oqqeil, Nafal	CC	I-II	H	A	Cul	Gra	3-4	Wee
Medicago	rigidula	Leguminosae	-	RR	I-II	H	A	Cul-Rge	Gra	2-3	Sil
Medicago	truncatula	Leguminosae	-	C	I-III	H	A	Cul-Rge-Ste	Gra	2-3	Psa
Medicago	tuberculata	Leguminosae	-	RR	I-II	H	A	Cul-Rge	Gra	2-3	Wee
Melilotus	elegans	Leguminosae	Handekoeq	RR	I-II	H	A	Cul-Rge	Non	0	Hal-Wee
Melilotus	indicus	Leguminosae	Handekoeq	CC	I-II	H	A	Cul-Rge	Med	0	Hal-Wee
Melilotus	sulcatus	Leguminosae	Handekoeq	R	I-II	H	A	Cul-Rge	Non	0	Hal-Wee
Onobrychis	crisia-galli	Leguminosae	Seleisa	C	I-II	H	A	Rge-Snd	Gra	2	Psa
Ononis	mitissima	Leguminosae	-	R	I-II	H	A	Cul-Wild	Non	0	Wee
Ononis	matrix subsp.atrix	Leguminosae	Littein	FR	I-II	DS	P	Rge	Gra	1	Psa
Ononis	reclinata	Leguminosae	Hotteiba	C	I-II	H	A	Cul	Non	0	Psa
Ononis	serrata	Leguminosae	-	C	I-II	H	A	Cul-Snd	Non	0	Psa
Ononis	scicula	Leguminosae	-	R	I-II	H	A	Cul-Wild	Non	0	Wee
Ononis	vaginalis	Leguminosae	-	CC	I	DS	P	Coa-Snd	Non	0	Psa
Pisum	sativum	Leguminosae	-	R	I-II	H	A	Cul	Gra	4-5	Wee
Prosopis	farcta	Leguminosae	-	R	I-II	DS	P	Cul-Irr-Wild	Fue	0	Phr-Wee
Prosopis	juliflora	Leguminosae	-	FC	I-II	HS-T	P	Wild	Fen-Fue-Fsh-Gra	2-3	Nat-Phr
Retama	monosperma	Leguminosae	R'tam	RR	I-II	HS	P	Rge-Wad	Fue-Gra-Sbi	1-2	Psa
Retama	raetam	Leguminosae	R'tam	FR	I-VI	HS	P	Rge-Wad	Fue-Gra-Med-Sbi	1-2	Psa
Scorpiurus	muricatus	Leguminosae	Thanab al Aqrab	CC	I-II	H	A	Cul	Gra	3	Pel
Securigera	securidaca	Leguminosae	-	RR	I-II	H	A	Cul	Non	0	Wee
Sesbania	sesban	Leguminosae	Seseebaan	Fc	I-II	HS	P	Irr	Wbr	-	Nat
Trifolium	alexandrinum	Leguminosae	Berseem	CC	I-II	H	A	Irr	Cro	5	Fod
Trifolium	angustifolium	Leguminosae	-	RR	I-II	H	A	Cul	Gra	1	Wee
Trifolium	argutum	Leguminosae	-	R	I-II	H	A	Cul	Gra	1	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Trifolium	campestre	Leguminosae	-	R	I-II	H	A	Cul	Gra	1	Wee
Trifolium	dasyurum	Leguminosae	-	R	I-II	H	A	Cul-Irr	Gra	1-2	Wee
Trifolium	fragiferum	Leguminosae	-	RR	I-II	H	P	Wet	Cro	5	Fod-Hal
Trifolium	nigrescens	Leguminosae	-	RR	I-II	H	A	Cul	Gra	1	Wee
Trifolium	patens	Leguminosae	-	RR	I-II	H	A	Wet	Gra	1-2	Wee
Trifolium	purpureum	Leguminosae	-	R	I-II	H	A	Cul	Cro	5	Fod
Trifolium	resupinatum	Leguminosae	Qort	CC	I-II	H	A	Wet	Cro	5	Fod-Hal
Trifolium	scabrum	Leguminosae	-	RR	I-II	H	A	Cul	Gra	1	Wee
Trifolium	stellatum	Leguminosae	-	R	I-II	H	A	Cul	Gra	1-2	Wee
Trifolium	tomentosum	Leguminosae	Kreishit	C	I-II	H	A	Rge-Wet	Gra	1-2	Wee
Trigonella	anguina	Leguminosae	-	FC	I-IV	H	A	Cul-Wid	Non	0	Hal-Tac-Wee
Trigonella	arabica	Leguminosae	Wideina	R	I-II	H	A	Cul-Snd	Gra	4	Psa
Trigonella	berlythea	Leguminosae	-	RR	I-II	H	A	Cul	Non	0	Wee
Trigonella	cylindrica	Leguminosae	-	R	I-II	H	A	Cul-Wid	Non	0	Wee
Trigonella	glabra	Leguminosae	Deraaq, Ishb al Malik	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Trigonella	laciniata	Leguminosae	Deraaq	CC	I-II	H	A	Cul-Wid	Non	0	Wee
Trigonella	maritima	Leguminosae	Deraaq	C	I-II	H	A	Cul-Wid	Non	0	Hal-Wee
Trigonella	occulta	Leguminosae	-	R	I-II	H	A	Wid	Non	0	Wee
Trigonella	stellata	Leguminosae	Gargas	C	I-II	H	A	Cul-Wid	Non	0	Hal-Wee
Trigonella	tetraphyllum	Leguminosae	-	RR	I-II	H	A	Cul-Ste-Wid	Non	0	Sil-Wee
Tripodion	ervilia	Leguminosae	-	RR	I-II	H	A	Cul	Gra	2	Wee
Vicia	lutea	Leguminosae	Bakr	C	I-II	H	A	Cul	Gra	2	Wee
Vicia	monantha	Leguminosae	Khareg	CC	I-III	H	A	Cul-Rge	Gra	3	Psa
Vicia	narbonensis	Leguminosae	Nooman Berri	C	I-II	H	A	Cul	Gra	2	Pel-Wee
Vicia	peregrina	Leguminosae	-	R	I-II	H	A	Cul	Gra	2	Wee
Vicia	sativa	Leguminosae	Eddeis	CC	I-II	H	A-B	Cul	Cro-Med	5	Fod
Vicia	villosa	Leguminosae	-	RR	I-II	H	A	Cul	Gra	3	Wee
Lemma	gibba	Lemnaceae	Ades el Mya	CC	I-II	H	P	Can-Frw-Wel	Non	0	Hyd
Pseudowolffia	hyalina	Lemnaceae	Ades el Mya	C	I-II	H	P	Can-Frw-Wel	Non	0	Hyd
Spirodelea	polyrrhiza	Lemnaceae	Ades el Mya	FC	I-II	H	P	Can-Frw-Wel	Non	0	Hyd
Allium	ampeloprasum	Liliaceae	Thoam	FR	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	aschersonianum	Liliaceae	-	R	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	barthianum	Liliaceae	-	R	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	blomfeldianum	Liliaceae	-	RR	I-II	H	P	Cul-Wid	Non	0	End-Wee
Allium	curtum	Liliaceae	-	R	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	desertorum	Liliaceae	Thoam Hanash	RR	I-II	H	P	Cul-Rge-Ste	Non	0	End-Wee
Allium	erdelli	Liliaceae	-	RR	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	mareoticum	Liliaceae	-	R	I-II	H	P	Cul-Wid	Non	0	End-Wee
Allium	myrianthum	Liliaceae	Lawas	RR	I-II	H	P	Cul-Wid	Non	0	End-Wee
Allium	pollens	Liliaceae	-	R	I-II	H	P	Cul-Wid	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Allium	roseum	Liliaceae	-	R	I-IV	H	P	Cul-Wid	Gra-Veg	3	Psa
Allium	sphaerocephalon	Liliaceae	Sennel Ghazzal	R	I-II	H	P	Cul-Wid	Non	0	Wee
Allium	trifoliatum	Liliaceae	-	RR	I-II	H	P	Cul-Wid	Non	0	Wee
Asparagus	aphyllus	Liliaceae	Aqool	FC	I-III	H	P	Rge-Run-Wid	Non	0	Cha
Asparagus	stipularis	Liliaceae	Sekkum	R	I-III	S	P	Wild	Fue	0	Fce
Asphodelus	ramosus	Liliaceae	Ansai	CC	I-III	H	P	Rge-Ste-Wid	Gra	2	Nit-Psa
Asphodelus	tenuifolius	Liliaceae	Barous	C	I-II	H	A	Rge-Ste-Wid	Non	0	Nit-Psa
Asphodelus	viscidulus	Liliaceae	Berouagh	CC	I-II	H	A	Rge-Ste-Wid	Non	0	Nit-Psa-Tac
Bellevia	eigii	Liliaceae	Bossei	RR	I-II	H	P	Cul	Non	0	Wee
Bellevia	flexuosa	Liliaceae	Bossei	C	I-II	H	P	Cul	Non	0	Wee
Bellevia	macrobotrys	Liliaceae	Bossei	RR	I-II	H	P	Cul	Non	0	Wee
Bellevia	mauritanica	Liliaceae	Bossei	RR	I-II	H	P	Cul	Non	0	Wee
Bellevia	romana	Liliaceae	Bossei	RR	I-II	H	P	Cul	Non	0	Wee
Bellevia	sessiliflora	Liliaceae	Bossei	RR	I-II	H	P	Cul	Non	0	Wee
Bellevia	trifoliata	Liliaceae	Bossei	R	I-II	H	P	Cul	Non	0	Wee
Colchicum	ritchii	Liliaceae	Okna	FR	I-II	H	P	Cul-Ste-Wid	Non	T	Sil-Wee
Dipcadi	erythraeum	Liliaceae	Hankal-ay	R	I-II	H	P	Rge-Ste-Wid	Non	0	Psa
Gagea	dyana	Liliaceae	-	R	I-II	H	P	Rge-Wid	Non	0	Wee
Gagea	fibrosa	Liliaceae	-	R	I-II	H	P	Rge-Wid	Non	0	Wee
Gagea	reticulata	Liliaceae	-	R	I-II	H	P	Rge-Wid	Non	0	Wee
Gagea	albiflorum	Liliaceae	Bossei	RR	I-II	H	P	Cul-Wid	Non	0	Wee
Muscari	bicolor	Liliaceae	Bossei	RR	I-II	H	P	Cul-Wid	Non	0	Wee
Muscari	comosum	Liliaceae	Bossei	CC	I-II	H	P	Cul-Wid	Non	0	Wee
Muscari	neglectum	Liliaceae	Bossei	R	I-III	H	P	Cul-Wid	Non	0	Wee
Muscari	parviflorum	Liliaceae	Bossei	R	I-II	H	P	Cul-Wid	Non	0	Wee
Ornithogalum	narbonense	Liliaceae	-	RR	I-II	H	P	Cul	Non	0	Wee
Ornithogalum	trichophyllum	Liliaceae	Basol Hanash	R	I-II	H	P	Cul	Non	0	Wee
Scilla	peruviana	Liliaceae	-	R	I-II	H	P	Rge-Wid	Non	T	Wee
Urginea	undulata	Liliaceae	-	C	I-II	H	P	Rge-Ste-Wid	Gra	1	Wee
Linum	decumbens	Linaceae	-	RR	I-II	H	A	Cul-Rge	Gra	1	Psa
Linum	strictum	Linaceae	Kittan	R	I-II	H	A	Cul-Rge	Gra	1	Psa
Linum	auriculata	Lythraceae	Rig el Hamma	R	I-II	H	A	Wet	Non	0	Hyg
Ammania	baccifera	Lythraceae	Rig el Hamma	C	I-II	H	A	Wet	Non	0	Hyg
Ammania	multiflora	Lythraceae	Rig el Hamma	RR	I-II	H	A	Wet	Non	0	Hyg
Ammania	senegalensis	Lythraceae	Rig el Hamma	R	I-II	H	A	Wet	Non	0	Hyg
Lythrum	hyssopifolia	Lythraceae	-	C	I-III	H	A	Wet	Non	0	Hyg
Lythrum	juncum	Lythraceae	-	R	I-II	H	P	Wet	Non	0	Hyg
Lythrum	tribractatum	Lythraceae	-	R	I-II	H	A-B	Wet	Non	0	Hyg
Alcea	acaulis	Malvaceae	Khatma	RR	I-II	H	A	Cul-Wid	Non	0	Wee
Althaea	ludwigii	Malvaceae	Khatma	FC	II-IV	H	A	Rge-Wid	Gra	2	Ere-Psa

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Hibiscus	trionum	Malvaceae	Teel Sheitani	FC	I-II	H	A	Cul	Non	0	Pel
Lavatera	cretica	Malvaceae	Khatma	R	I-II	H	A-B	Cul	Orn	0	Wee
Malva	aegyptia	Malvaceae	Raqma	FR	I-III	H	A	Rge-Wld	Gra-Veg	1-2	Psa
Malva	nicaensis	Malvaceae	Khobbiza	FC	I-II	H	A	Cul-Wld	Gra-Veg	1-2	Nit
Malva	parviflora	Malvaceae	Khobbiza	CC	I-II	H	A	Rge-Wld	Gra-Med-Veg	1-2	Nit
Malva	sylvestris	Malvaceae	Khobbiza	FC	I-II	H	A	Cul-Wld	Gra-Med-Veg	1-2	Nit
Sida	alba	Malvaceae	-	FC	I-III	H	P	Rge-Wld	Gra	1	Wee
Marsilea	aegyptiaca	Marsiliaceae	Qoreyfa	R	I-II	F	P	Can-Wel	Non	0	Hyd
Gilvus	lotoides	Molluginaceae	Ghobbeira, Mughera	R	I-III	H	A	Wld	Non	0	Hal
Neurada	procumbens	Neuradaceae	Losseiq	CC	II-VI	H	A	Rge-Snd-Ste	Non	0	Psa-Tac
Nymphaea	coerulea	Nymphaeaceae	Bashmeen azreg	C	I-II	H	P	Frw	Orn	0	Hyd
Nymphaea	lotus	Nymphaeaceae	Bashmeen abiod	C	I-II	H	P	Frw	Med-Orn	0	Hyd
Olea	europaea subsp. oleaster	Oleaceae	Zebuj	RR	I-II	TS	P	Roc	Fsh-Non	3	FoRe
Phyllirea	media	Oleaceae	-	RR	I-II	TS	P	Roc	Fue	3	FoRe
Epilobium	hirsutum	Onagraceae	Alfa, Sikh	FC	I-II	H	P	Wet	Non	0	Hyg
Ludwigia	stolonifera	Onagraceae	-	R	I-II	H	P	Wet	Non	0	Hyg
Cistanche	phelypaea	Orobanchaceae	Halook, Danum	CC	I-II	Par	P	Cul	Med	0	Hal-Wee
Cistanche	tubulosa	Orobanchaceae	-	FR	I-II	Par	P	Cul	Non	0	Hal-Wee
Cistanche	violacea	Orobanchaceae	-	FR	I-II	Par	P	Cul	Non	0	Hal-Wee
Orobanchae	cernua	Orobanchaceae	-	R	I-II	Par	A	Cul	Non	0	Wee
Orobanchae	crenata	Orobanchaceae	Haluik Mitabi	CC	I-II	Par	A	Cul	Non	0	Wee
Orobanchae	lavandulacea	Orobanchaceae	-	RR	I-III	Par	A	Cul	Non	0	Wee
Orobanchae	minor	Orobanchaceae	-	R	I-II	Par	A	Cul	Non	0	Wee
Orobanchae	ramosa	Orobanchaceae	Halook	CC	I-III	Par	P	Cul	Non	0	Wee
Orobanchae	schultzi	Orobanchaceae	-	RR	I-III	Par	A	Cul	Non	0	Wee
Oxalis	corniculata	Oxalidaceae	Hamd	FC	I-II	H	P	Cul-Wld	Non	0	Wee
Oxalis	pes-caprae	Oxalidaceae	Hamd	C	I-II	H	P	Cul-Wld	Non	0	Wee
Phoenix	dactylifera	Palmae	Naghia	CC	I-VI	H	P	Cul-Run-Wad	Han-Med-Veg	0	Phr
Glaucium	arabicum	Papaveraceae	-	R	I-II	H	P	Cul	Non	0	Wee
Glaucium	corniculatum	Papaveraceae	-	R	I-II	H	A	Cul-Wld	Non	0	Wee
Glaucium	grandiflorum	Papaveraceae	-	R	I-II	H	A	Wld	Non	0	Wee
Papaver	argemone	Papaveraceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Papaver	dubium	Papaveraceae	-	R	I-II	H	A	Cul	Med	0	Wee
Papaver	humile	Papaveraceae	-	R	I-II	H	A	Cul	Non	0	Wee
Papaver	hybridum	Papaveraceae	-	C	I-II	H	A	Cul	Non	0	Wee
Papaver	rhoeas	Papaveraceae	Ben Na'am	CC	I-II	H	A	Cul	Med	0	Wee
Roemeria	hybrida	Papaveraceae	-	R	I-II	H	A	Cul	Non	0	Tac-Wee
Potamogeton	cripus	Potamogetonaceae	Horreish	CC	I-II	H	P	Can-Frw	Orn	0	Hyd
Potamogeton	lucens	Potamogetonaceae	-	RR	I-II	H	P	Can-Frw	Orn	0	Hyd
Potamogeton	nodosus	Potamogetonaceae	-	FC	I-II	H	P	Can-Frw	Orn	0	Hyd


Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Potamogeton	pectinatus	Potamogetonaceae	-	CC	I-II	H	P	Can-Frw	Orn	0	Hyd
Plantago	afra	Plantaginaceae	Ootoona, Aslouj	R	I-II	H	A	Cul-Wild	Med	0	Psa-Tac
Plantago	albicans	Plantaginaceae	Yanam	CC	I-III	H	P	Rge-Snd-Ste	Gra	4-5	Psa
Plantago	arenaria	Plantaginaceae	-	R	I-II	H	A	Cul	Non	0	Psa
Plantago	coronopus	Plantaginaceae	Wideina	CC	I-III	H	A	Wild	Gra-Med	1	Hal
Plantago	crassifolia	Plantaginaceae	-	FR	I-II	H	P	Sit	Gra	3	Hal
Plantago	crispoides	Plantaginaceae	-	CC	I-III	H	A	Wild	Gra	2	Hal
Plantago	cylindrica	Plantaginaceae	Berkhemi	FR	I-II	H	A	Irr	Non	0	Wee
Plantago	lagopus	Plantaginaceae	Widna	FC	I-II	H	A	Irr	Non	0	Sil
Plantago	major	Plantaginaceae	Warraq, Lisaan al Hamal	R	I-II	H	A	Irr	Med	0	Hyg-Nit-Sil
Plantago	notata	Plantaginaceae	Yanam	R	I-II	H	A	Rge	Gra	2	Psa
Plantago	ovata	Plantaginaceae	Gesseima	CC	I-III	H	A	Rge	Gra	2	Sil
Plantago	phaeostoma	Plantaginaceae	-	R	I-II	H	A	Cul-Rge	Gra	1	Psa-Tac
Plantago	squarrosa	Plantaginaceae	-	C	I-II	H	A	Cul	Non	0	Sil
Plantago	weldenii	Plantaginaceae	Gesseima	R	I-III	H	A	Cul-Rge	Gra	1	Hal
Limoniastrum	guyonianum	Plumbaginaceae	Zeita	RR	I-III	HS	P	Sit	Fue	0	Hal
Limoniastrum	monopetalum	Plumbaginaceae	Zeita	C	I-III	HS	P	Sit	Fue	0	Hal
Limonium	echioides	Plumbaginaceae	-	RR	I-II	H	A	Rge-Ste	Non	0	Hal-Psa
Limonium	lobatum	Plumbaginaceae	-	CC	I-III	H	A	Rge-Sha-Ste	Non	0	Gly
Limonium	narbonense	Plumbaginaceae	-	R	I-II	H	P	Sit	Non	0	Hal
Limonium	pruinatum	Plumbaginaceae	Molleih	CC	I-III	Fr	P	Rge-Sha-Ste	Non	0	Hal-SO4
Limonium	raddianum	Plumbaginaceae	-	R	I-II	H	P	Sit	Non	0	Hal
Limonium	sinuatum	Plumbaginaceae	-	FC	I-III	H	A	Rge-Sha-Ste	Non	0	Gly
Limonium	tubiflorum	Plumbaginaceae	Bahman	C	I-III	H	P	Rge-Sha	Non	0	Cha-Hal
Calligonum	comosum	Polygonaceae	Artha	FR	III-VI	HS	P	Con-Dne	Fue-Gra	2	Psa
Emex	spinus	Polygonaceae	Dirs el Aggoz	CC	I-III	H	A	Wild	Unk	0	Nit
Fallopia	convolvulus	Polygonaceae	-	R	I-II	H	A	Cul	Non	0	Hyg-Wee
Persicaria	aviculare	Polygonaceae	Quordab	R	I-II	H	P	Wet	Non	1	Hyg
Persicaria	bellardi	Polygonaceae	Quordab	CC	I-II	H	A	Wet	Non	0	Hyg
Persicaria	limbata	Polygonaceae	Quordab	RR	I-II	H	P	Wet	Non	0	Hyg
Persicaria	maculosa	Polygonaceae	Quordab	R	I-II	H	A	Wet	Non	0	Hyg
Persicaria	maritimum	Polygonaceae	Quordab	FR	I	Fr-DS	P	Coa-Dne	Non	0	Psa
Persicaria	plebeium	Polygonaceae	Quordab	CC	I-II	H	P	Wet	Non	0	Hyg
Persicaria	salicifolia	Polygonaceae	Quordab	CC	I-II	H	P	Wet	Non	0	Hyg
Persicaria	senegalensis	Polygonaceae	Quordab	C	I-II	H	P	Wet	Non	0	Hyg
Polygonum	equisetiforme	Polygonaceae	Quordab	FC	II-III	Fr-DS	P	Rge-Ste	Gra	3-4	Psa
Rumex	cyprus	Polygonaceae	Khamsees	C	I-II	H	A	Wet	Unk	2-3	Cha
Rumex	dentatus	Polygonaceae	Hommeid	R	I-II	H	A	Irr-Wet	Unk	0	Hyg-Wee
Rumex	pictus	Polygonaceae	Hommeid	C	I-II	H	A	Wet	Unk	0	Hyg-Wee
Rumex	vesicarius	Polygonaceae	Hambeit, Hummad	FC	I-III	H	A	Rge-Ste	Gra-Med-Veg	0	Nit

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Eichhornia	crassipes	Pontederiaceae	Halassandi	CC	I-II	H	P	Can-Frw	Orn	2	Hyd-Nat-Wee
Portulaca	oleracea	Portulacaceae	Rigla, Rashad	CC	I-II	CH	A	Wld	Med-Veg	2	Nit
Posidonia	oceanica	Posidoniaceae	Shetenaara	RR	I	H	P	Sea	Non	0	Hyd
Anagallis	arvensis	Primulaceae	Ebeila, Ayn al Gamal	CC	I-III	H	A	Cul-Wld	Non	0	Wee
Asterolinon	linum-stellatum	Primulaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Coris	monspeliensis	Primulaceae	-	RR	I-III	CH	P	Rge-Ste	Non	0	S04
Samolus	valerandi	Primulaceae	Sboon Arab	R	I-II	H	P	Wet	Non	0	Hyg
Adonis	dentata	Ranunculaceae	Naab el Gemel	CC	I-III	H	A	Cul-Rge-Wld	Gra-Med	1	Psa-Tac
Amaranthus	blitoides	Ranunculaceae	-	RR	I-II	H	A	Wld	Non	0	Wee
Amaranthus	coronaria	Ranunculaceae	Zagheel	C	I-II	H	A	Cul-Wld	Non	T	Wee
Anemone	flava	Ranunculaceae	-	RR	I-II	H	A	Cul-Rge-Wld	Non	0	Wee
Consolida	bovei	Ranunculaceae	-	RR	I-II	H	A	Cul-Rge	Non	0	Psa
Delphinium	arvensis subsp. taubertii	Ranunculaceae	Kamoun Aswad	R	I-II	H	A	Cul	Non	0	Wee
Nigella	deserti	Ranunculaceae	Zorbeb	RR	I-II	H	A	Cul-Wld	Non	0	End-Wee
Nigella	asiaticus	Ranunculaceae	Shaqqeq	C	I-II	H	A	Cul-Wld	Non	0	Wee
Ranunculus	millefolius	Ranunculaceae	-	RR	I-II	H	P	Frw	Non	0	Hyd
Ranunculus	muricatus	Ranunculaceae	-	RR	I-II	H	A	Frw	Non	0	Hyg
Ranunculus	peltatus	Ranunculaceae	Zaghanta	RR	I-II	H	A	Wld	Non	0	Hyd
Ranunculus	hexagyna	Resedaceae	Denaaba	FC	II-IV	H	A	Cul-Rge	Gra	1	Tac-Wee
Cayusea	baccatus	Resedaceae	Qordi	RR	III-VI	H	P-Sh	Des-Wad	Fue-Gra	1-2	Psa
Ochradenus	linifolia	Resedaceae	Denaaban	FC	II-VI	H	A	Cul-Rge	Gra	1	Ere-Tac
Oligomeris	africana	Resedaceae	-	FR	III-VI	DS	P	Des-Wad	Gra	1-2	Ere
Randonia	alba	Resedaceae	-	R	I-II	H	A	Cul-Rge	Gra	1	Wee
Reseda	arabica	Resedaceae	Deil	CC	I-III	H	A	Cul-Rge-Site	Gra	1	Tac-Wee
Reseda	decursiva	Resedaceae	Rigl Gorab	CC	I-III	H	A-B	Cul-Rge-Site	Gra	2-3	Psa
Reseda	lutea	Resedaceae	Weyba	RR	I-III	H	A	Cul-Rge	Gra-Med	1	Wee
Reseda	luteola	Resedaceae	Baqm, Bli ha	R	I-III	H	A	Cul-Rge	Gra	1	Wee
Reseda	orientalis	Resedaceae	Deil	RR	I-III	H	A	Cul-Rge	Gra	1	Wee
Reseda	phyteuma	Resedaceae	-	RR	I-II	H	A	Cul-Rge	Gra	1	Wee
Reseda	pruinosa	Resedaceae	Denaaba	CC	I-II	H	A	Cul-Rge	Gra	1	Wee
Reseda	urnigera	Resedaceae	-	R	I-III	H	A	Cul-Rge	Gra	1	Wee
Rhamnus	oleoides	Rhamnaceae	Saboufa	RR	I-II	HS	P	Cli-Run	Fsh-Gra	3-4	Cha
Ziziphus	lotus	Rhamnaceae	Sidra	RR	I-II	HS	P	Cli-Wad	Fen-Fue-Med	2	Phr
Ziziphus	spina-christi	Rhamnaceae	Nabq, Zegzeg	RR	I-IV	T	P	Des-Wad	Fue-Med-Veg	2	Phr
Rubus	sanctus	Rosaceae	Ward Barrir	R	I-II	Cl	P	Wld	Non	0	Nit
Sanguisorba	minor	Rosaceae	Abu Graisaf	R	I-II	H	P	Cul-Rge	Gra	3-4	Wee
Callipeltis	cucullaris	Rubiaceae	Baseesa	FR	I-II	H	A	Cli-Roc	Non	0	Cha-Nit
Crucianella	aegyptiaca	Rubiaceae	-	C	I-II	H	P	Cul	Non	0	Psa
Crucianella	maritima	Rubiaceae	-	C	I	Fr	P	Coa-Snd	Non	0	Psa
Galium	canum	Rubiaceae	-	RR	I-II	H	P	Cul	Non	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Gallium	murale	Rubiaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Gallium	tricornutum	Rubiaceae	-	C	I-II	H	A	Cul	Non	0	Wee
Valantia	hispidia	Rubiaceae	-	FR	I-II	H	A	Roc	Non	0	Cha
Ruppia	cirrrosa	Ruppiaceae	Hamool	RR	I-II	H	P	Sea-Bra	Non	0	Hyd
Ruppia	maritima	Ruppiaceae	Reem	RR	I-II	H	P	Sea-Bra	Non	0	Hyd
Salix	subserrata	Salicaceae	Safsaf	R	I-II	HS-T	P	Wet	Orn	1	Hyg
Salix	tetrasperma	Salicaceae	Safsaf	R	I-II	HS-T	P	Wet	Fue	1	Hyg
Thesium	humile	Santalaceae	Habb el Kreish	FC	I-II	H-Par	A	Cul-Rge	Unk	0	Wee
Antirrhinum	orontium	Scrophulariaceae	Saysam	C	I-II	H	A	Cul-Rge	Gra	2	Psa
Kickxia	egyptiaca	Scrophulariaceae	Thal el Fahr	CC	I-III	Fr	P	Rge-Sha-Ste	Gra	2	Psa
Kickxia	elatine	Scrophulariaceae	-	C	I-II	H	A	Cul-Rge	Gra	1	Psa
Linaria	albifrons	Scrophulariaceae	-	FC	I-II	H	P	Cul	Non	0	Wee
Linaria	chalepensis	Scrophulariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Linaria	haelava	Scrophulariaceae	Halawa	C	I-II	H	A	Cul-Rge	Gra	1	Wee
Linaria	micrantha	Scrophulariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Scrophularia	canina	Scrophulariaceae	Qarsa	FR	I-III	Fr	P	Rge-Snd	Non	0	Psa
Striga	asiatica	Scrophulariaceae	Odaar	RR	I-II	Par	A	Cul-Irr	Non	0	Wee
Verbascum	letourneuxii	Scrophulariaceae	-	FC	I-II	Fr	P	Rge-Sha	Non	0	End
Veronica	persica	Scrophulariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Veronica	polita	Scrophulariaceae	-	FC	I-II	H	A	Cul	Non	0	Wee
Veronica	syriaca	Scrophulariaceae	-	RR	I-II	H	A	Cul	Non	0	Wee
Datura	metel	Solanaceae	-	R	I-II	H	A	Wid	Med	T	Nit
Datura	stramonium	Solanaceae	Semmel Fahr	R	I-II	H	A	Wid	Med	T	Nit
Hyosциamus	albus	Solanaceae	Bing	FC	I-II	H	P	Wid	Med	T	Nit
Hyosциamus	muticus	Solanaceae	Sakaraan	FC	I-VI	H	P	Wad-Wid	Med	T	Ere-Pel
Lycium	aschersoni	Solanaceae	Awsaj.	R	I-II	HS	P	Wid	Fen-Fue	0	S04
Lycium	barbarum	Solanaceae	Awsaj.	C	I-II	HS	P	Rge-Wad	Fue-Gra	1	S04
Lycium	europaeum	Solanaceae	Awsaj.	R	I-II	HS	P	Wid	Fen-Fue	1	S04
Lycium	schweinfurthii	Solanaceae	Awsaj.	R	I-II	HS	P	Wid	Fen-Fue	0	S04
Lycium	shawii	Solanaceae	Awsaj.	CC	I-III	HS	P	Rge-Wad	Fue-Gra-Med	1	S04
Nicotiana	glauca	Solanaceae	Massasa	CC	I-II	HS	P	Wid	Med	T	Nat-Nit
Solanum	nigrum	Solanaceae	Anab al dib	R	I-II	H	A	Wad	Non	T	Nit
Reaumuria	hirtella	Tamaricaceae	Molleih	CC	I-III	Fr	P	Rge-Sha-Ste	Gra	2	Hal-S04
Reaumuria	vermiculata	Tamaricaceae	Molleih	R	I-III	Fr	P	Rge-Sha-Ste	Gra	1-2	Hal-S04
Tamarix	aphylla	Tamaricaceae	Ethel	C	I-VI	T	P	Des-Wad	Fue	0	Phr
Tamarix	nilotica	Tamaricaceae	Moor	CC	I-VI	HS	P	Sit	Fue	0	Hal-Phr
Tamarix	passeriformis	Tamaricaceae	-	C	I-IV	HS	P	Wad	Fue	0	Hal-Phr
Tamarix	tetragyna	Tamaricaceae	Tarfa	C	I-VI	HS	P	Sit	Fue	0	Hal-Phr
Thymelea	hirsuta	Thymelaeaceae	Mitnan	CC	I-III	HS	P	Rge-Wid	Fue-Han-Med	T	Psa
Corchorus	olitorius	Tiliaceae	Melokhyia	CC	I-III	H	A	Irr	Gra-Veg	0	Wee

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Typha	domingensis	Typhaceae	Typhaceae	CC	I-III	H	P	Can-Frw	Han	0	Hyd-Hyg
Ammi	majus	Umbelliferae	Khilla Baladi	C	I-II	H	A	Cul	Med	0	Pel-Sil-Wee
Ammi	visnaga	Umbelliferae	Khilla sheitani	R	I-II	H	A	Wet-Wid	Med	T	Hyg-Pel
Apium	crassipes	Umbelliferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Apium	graveolens	Umbelliferae	Karafs	CC	I-II	H	A	Cul	Non	0	Wee
Apium	leptophyllum	Umbelliferae	-	FC	I-II	H	A	Cul	Non	0	Wee
Apium	nodiflorum	Umbelliferae	-	RR	I-II	H	P	Wet	Non	0	Hyg
Brachyapium	dichotomum	Umbelliferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Bupleurum	lancifolium	Umbelliferae	Halawaan	C	I-II	H	A	Cul	Non	0	Pel-Wee
Bupleurum	nanum	Umbelliferae	-	R	I-II	H	A	Cul	Non	0	Wee
Bupleurum	nodiflorum	Umbelliferae	-	R	I-II	H	A	Cul	Non	0	Pel-Wee
Bupleurum	semicompositum	Umbelliferae	-	FC	I-II	H	A	Cul-Wid	Non	0	Hal-Pel
Coriandrum	sativum	Umbelliferae	Kozbara	C	I-II	H	A	Irr	Non	0	Wee
Critihum	maritimum	Umbelliferae	-	RR	I	C-Fr	P	Roc	Med	0	Cha
Daucus	guttatus	Umbelliferae	Shamarel Gemel	R	I-II	H	A	Cul	Non	0	Wee
Daucus	litoralis	Umbelliferae	Gezzah	C	I-II	H	A	Cul-Wid	Non	0	Wee
Daucus	syrticus	Umbelliferae	Shamarel Gemel	CC	I-III	H	A	Cul-Rge	Gra	2	Psa-Tac
Deverra	tortuosa	Umbelliferae	Qezzah	CC	I-IV	DS	P	Cul-Rge	Gra-Med	2-3	Psa
Eryngium	campestre	Umbelliferae	Foq'aa	C	I-II	H	P	Cul-Rge-Ste	Med	0	Pel-Sil
Eryngium	creticum	Umbelliferae	Shaqaool	RR	I-II	H	P	Cul-Rge	Non	0	Pel-Sil-Wee
Eryngium	marmarica	Umbelliferae	Felkh	RR	I-II	H	P	Sha-Site	Non	T	End
Malabaila	suaveolens	Umbelliferae	Tel ghoodi	FC	I-II	H	P	Cul-Rge	Non	0	Wee
Petroselinum	crispum	Umbelliferae	-	RR	I-II	H	P	Wet	Non	0	Hyg
Pseudoraiya	pumila	Umbelliferae	Shamarel Gemel	C	I	H	A	Coa-Snd	Non	0	Psa-Tac
Scandix	specten-veneris	Umbelliferae	Misheita	C	I-II	H	A	Cul	Non	0	Wee
Tordylium	aegyptiacum	Umbelliferae	-	RR	I-II	H	A	Cul	Non	0	Wee
Torilis	arvensis	Umbelliferae	Shamarel Gemel	R	I-II	H	A	Cul	Non	0	Wee
Torilis	bracteosa	Umbelliferae	-	R	I-II	H	P	Cul	Non	0	Wee
Torilis	nodosa	Umbelliferae	Shamarel Gemel	CC	I-II	H	A	Cul-Wid	Non	0	Sil-Wee
Torilis	radiata	Umbelliferae	Ghazzar Frangi	R	I-II	H	A	Cul	Non	0	Wee
Torilis	tenella	Umbelliferae	Shamarel Gemel	R	I-II	H	A	Cul	Non	0	Wee
Parietaria	judaea	Urticaceae	Widein	FC	I-II	H	A	Wid	Med	0	Nit
Parietaria	lustranica	Urticaceae	Widein	R	I-II	H	A	Roc	Med	0	Nit
Urtica	pilulifera	Urticaceae	Qorreis	FC	I-II	H	A	Wid	Med	0	Nit
Urtica	urens	Urticaceae	Qorreis, Hurreq	FC	I-II	H	A	Wid	Med	0	Nit
Valerianella	discoidea	Valerianaceae	Gesseima	RR	I-II	H	A	Rge-Wid	Gra	1	Wee
Valerianella	petrovitchii	Valerianaceae	Gesseima	RR	I-II	H	A	Cul-Wid	Gra	1	Wee
Phyla	nodiflora	Verbenaceae	Libbya	FC	I-II	H	P	Wad-Wet	Gra	1	Hyg
Verbena	officialis	Verbenaceae	Rigi al Hammam	FR	I-II	H	P	Cul	Med-Veg	0	Hyg
Verbena	supina	Verbenaceae	Morreig	FC	I-II	H	A	Wad	Gra	1	Hyg

Genus	Species	Family	Vernacular name	Frequency	Area of distribution	Biological type	Life span	Habitat	Utilization	Grazing value	Remarks
Viola	scopiuroides	Violaceae	-	FR	I-II	Fr	P	Cli	Gra	1	Cha-End
Zannichellia	palustris	Zannichelliaceae	Lahm el Naga	CC	I-II	H	P	Can-Frw	Non	0	Hyd
Zostera	noltii	Zosteraceae	Hamool	RR	I	H	P	Sea	Non	0	Hyd
Fagonia	arabica	Zygophyllaceae	-	FR	III-VI	Fr	P	Des-Rge	Non	0	Psa
Fagonia	bruguieri	Zygophyllaceae	Hallawa	FR	III-VI	Fr	P	Des-Rge	Non	0	Psa-Tac
Fagonia	cretica	Zygophyllaceae	Hallawa	C	I-III	Fr	P	Cli-Roc-Run	Non	0	Cha
Fagonia	glutinosa	Zygophyllaceae	Hallawa	FC	II-IV	Fr	P	Rge-Ste	Non	0	Psa
Fagonia	retusa	Zygophyllaceae	Ghardaq	FC	I-III	TS	P	Rge-Wid	Fue-Gra	1	Phr-SO4
Nitraria	harmala	Zygophyllaceae	Harmel	CC	I-III	H	P	Rge-Ste-Wid	Non	T	Nit
Peganum	tenella	Zygophyllaceae	-	RR	I-III	H	A	Slt	Non	0	Hal
Tetradiclis	terrestris	Zygophyllaceae	Shiq Shiq	CC	I-III	H	A	Rge-Wid	Non	0	Psa-Tac
Tribulus	aegyptium	Zygophyllaceae	Belbel	R	II-IV	Fr	P	Rge-Slt	Non	0	End
Zygophyllum	album	Zygophyllaceae	Belbel	CC	I-VI	CS	P	Rge-Wid	Non	0	Hal-SO4
Zygophyllum	coccineum	Zygophyllaceae	Qarlam	FC	III-VI	CS	P	Des-Wad	Non	0	Pel
Zygophyllum	geslini	Zygophyllaceae	Belbel	C	I-VI	CS	P	Wid	Non	0	Hal-SO4



Map 1. – Map of the Marmarica.