

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève
Band: 60 (2005)
Heft: 2

Artikel: Cytological investigation on some Ranunculus-species from Crete
Autor: Balisberger, Matthias / Widmer, Alex
DOI: <https://doi.org/10.5169/seals-879276>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Cytological investigations on some *Ranunculus*-species from Crete

MATTHIAS BALTISBERGER
&
ALEX WIDMER

ABSTRACT

BALTISBERGER, M. & A. WIDMER (2005). Cytological investigations on some *Ranunculus*-species from Crete. *Candollea* 60: 335-344. In English, English and French abstracts.

The chromosome numbers are presented for seven species (20 sites) of *Ranunculus* L. from Crete, and taxonomic and phytogeographic aspects are discussed. The number of chromosomes for *Ranunculus cupreus* Boiss. & Heldr. ($2n = 16$) is recorded for the first time. A new ploidy level was found in *Ranunculus paludosus* Poir. (pentaploid with $2n = 5x = 40$), and karyotypes are given for *Ranunculus asiaticus* L., *Ranunculus creticus* L. and *Ranunculus cupreus*.

RÉSUMÉ

BALTISBERGER, M. & A. WIDMER (2005). Données cytologiques de quelques espèces du genre *Ranunculus* de Crète. *Candollea* 60: 335-344. En anglais, résumés anglais et français.

Les nombres chromosomiques de sept espèces de *Ranunculus* (20 localités) de Crète sont présentés, et des aspects taxonomiques et phytogéographiques sont commentés. Le nombre de chromosomes pour *Ranunculus cupreus* Boiss. & Heldr. ($2n = 16$) est compté pour la première fois. Un nouveau degré de ploïdie a été trouvé pour *Ranunculus paludosus* Poir. (pentaploïde avec $2n = 5x = 40$), et les caryotypes de *Ranunculus asiaticus* L., *Ranunculus creticus* L. et de *Ranunculus cupreus* sont discutés.

KEY-WORDS: *Ranunculus* – Crete – Chromosome numbers – Karyotypes

Introduction

During an excursion to Israel and Crete in March 2000, plants were collected for biosystematic and cytological analyses and were transferred alive to the greenhouse of our institute. On these plants cytological investigations have been carried out. The results of the Israeli plants have been published recently (BALTISBERGER & WIDMER, 2004).

The cytological variability (chromosome numbers, karyotypes, ploidy levels) within the genus *Ranunculus* is rather high and not fully known, and karyological features are important for the understanding of natural groups within the genus (HÖRANDL & al., 2005). Additionally, as the genus *Ranunculus* is of special interest for us (see e.g. BALTISBERGER, 1980, 1992, 1994; MÜLLER & BALTISBERGER, 1984), we collected the *Ranunculus*-species met in Crete. Where possible, plants were collected from several sites. Cytological investigations have been carried out and the results are presented here. The seven investigated species are arranged in alphabetical order. Nomenclature follows *Flora Hellenica* (STRID, 2002).

The chromosome numbers are presented and discussed, including interesting aspects of systematics and/or geographic distribution. Literature concerning chromosome numbers was checked using FEDEROV (1974), VAN LOON (1987), as well as the series "Index to plant chromosome numbers" (MOORE, 1970, 1971, 1972, 1973, 1974, 1977; GOLDBLATT, 1981, 1984, 1985, 1988; GOLDBLATT & JOHNSON, 1990, 1991, 1994, 1996, 1998, 2000, 2003). In *Ranunculus asiaticus* L., *R. creticus* L. and *R. cupreus* Boiss. & Heldr., karyotypes are discussed.

Materials and methods

The cytological investigations were carried out on root tips. These were pretreated with colchicine (0.05%) for 2 hours, then fixed in ethanol/acetic acid (3:1), and stained and squashed in lacto-propionic orcein (DYER, 1963). Five to ten metaphases were counted out of each individual to determine the chromosome number, and if possible several individuals were investigated of each species (or each site, respectively). The numbers of investigated individuals are indicated in Table 1.

Karyotypes are discussed using the terminology for chromosome morphology proposed by LEVAN & al. (1964). The chromosomes are named according to the position of the centromeres, which is expressed with the arm ratio, "long arm to short arm". The following terms are used:

– metacentric	arm ratio	1.0-1.7;
– submetacentric	arm ratio	1.7-3.0;
– subtelocentric	arm ratio	3.0-7.0;
– acrocentric	arm ratio	more than 7.0;
– telocentric	only one arm.	

Plant material

After the indications of the site, date of sampling, number of herbarium specimen (in parenthesis specimen number of cultivated plants) are listed. Specimens are deposited in Z + ZT.

1. *Ranunculus asiaticus* L.

GREECE. Crete: Meadow, Moni Kardiotissis, near Krasion, N. Iraklion, 18.IV.1996, leg. F. Klötzli (*M. Baltisberger* 13230); Open slope, W of Mires, N. Iraklion, 18.III.2000, *M. Baltisberger* & *A. Widmer* 13545 (*M. Baltisberger* 13731); Rocky slope, NNW of Ag. Galini, on the road from Ag. Galini to Spili, N. Rethimno, 19.III.2000, (*M. Baltisberger* 13740); Bushland near Lampini, WNW of Spili, N. Rethimno, 20.III.2000, (*M. Baltisberger* 13744); Rocky slope between Selia and Rodakino, W of Spili, N. Rethimno, 20.III.2000, (*M. Baltisberger* 13741).

2. *Ranunculus creticus* L.

GREECE. Crete: Rocks in the gorge of Therisso, near Omalos, SSW of Chania, N. Chania, 18.IV.1991, *M. Eggenberger* & *B. Merz* (*M. Baltisberger* 12793); N exposed, rocky slope, NNW of Ag. Galini, on the road from Ag. Galini to Spili, N. Rethimno, 19.III.2000, *M. Baltisberger* & *A. Widmer* 13555 (*M. Baltisberger* 13716); Gorge S of Ag. Ioanis, on the road from Rethimno to Selia, SSW of Rethimno, N. Rethimno, 20.III.2000, *M. Baltisberger* & *A. Widmer* 13559 (*M. Baltisberger* 13722); N exposed rocks near the branches to Roustika off the road from Atsipopoulo to Episkopi, WSW of Rethimno, N. Rethimno, 21.III.2000, *M. Baltisberger* & *A. Widmer* 13561 (*M. Baltisberger* 13726).

3. *Ranunculus cupreus* Boiss. & Heldr.

GREECE. Crete: Rocky slope between Selia and Rodakino, W of Spili, N. Rethimno, 20.III.2000, *M. Baltisberger & A. Widmer 13558 (M. Baltisberger 13721, M. Baltisberger 13811, M. Baltisberger 13892;* duplicates in herb. Kit Tan, Copenhagen).

4. *Ranunculus ficaria* L.

GREECE. Crete: Wet place in culture of *Olea europaea* L., near Lampini, WNW of Spili, N. Rethimno, 20.III.2000, *M. Baltisberger & A. Widmer 13557 (M. Baltisberger 13719);* Wet place in culture of *Olea europaea* L., between Episkopi and Kournas, WSW of Rethimno, N. Chania, 21.III.2000, *M. Baltisberger & A. Widmer 13560 (M. Baltisberger 13724).*

5. *Ranunculus marginatus* d'Urv.

GREECE. Crete: Wet ruderal place, NW of Ag. Galini, on the road from Ag. Galini to Spili, N. Rethimno, 19.III.2000, (*M. Baltisberger 13717*).

6. *Ranunculus muricatus* L.

GREECE. Crete: Wet ruderal place, NW of Ag. Galini, on the road from Ag. Galini to Spili, N. Rethimno, 19.III.2000, (*M. Baltisberger 13718*); Open wet place in culture of *Olea europaea* L., near Lampini, WNW of Spili, N. Rethimno, 20.III.2000, (*M. Baltisberger 13720*); Wet ruderal place in the gorge S of Ag. Ioanis, on the road from Rethimno to Selia, SSW of Rethimno, N. Rethimno, 20.III.2000, (*M. Baltisberger 13723*); Open wet place in culture of *Olea europaea* L., between Episkopi and Kournas, WSW of Rethimno, N. Chania, 21.III.2000, (*M. Baltisberger 13725*).

7. *Ranunculus paludosus* Poir.

GREECE. Crete: Slope near Krasion, N. Iraklion, 17.IV.1996, leg. F. Klötzli (*M. Baltisberger 13231*); Culture of *Olea europaea* L., NW of Ag. Galini, on the road from Ag. Galini to Spili, N. Rethimno, 19.III.2000, (*M. Baltisberger 13814*); Bushland near Lampini, WNW of Spili, N. Rethimno, 20.III.2000, (*M. Baltisberger 13815*).

Results and discussion

1. *Ranunculus asiaticus* ($2n = 16$; Fig. 1A)

Ranunculus asiaticus is a rather variable species growing in open and waste places and shrubby vegetations in the Mediterranean and W Irano-Turanian area. It varies in the shape of the leaves and the colour of the flowers (red, white, and yellow), and several variants have been named. All plants from all five sites have undivided or 3-fid lower leaves with broad lobes as well as white flowers, only few plants from Lampini (*M. Baltisberger 13744*) showed flowers tinged pale rose-pink. As different morphological characters may occur in the same population, all varieties seem to fall within one range of variation of *R. asiaticus* and have therefore no systematic value (STRID, 2002). But there seems to be a tendency for the geographical distribution of the flower colours, plants from Greece showing mostly white flowers and plants from more eastern parts as Turkey (DAVIS, 1965) or Israel (BALDISBERGER & WIDMER, 2004) are red flowered.

All plants investigated from all five sites were diploid with $2n = 16$ chromosomes. The chromosome number corresponds with the indications in literature. The karyotype consists of 6 metacentric, 8 submeta- to subtelocentric and 2 acrocentric chromosomes (6m 8sm/st 2a),

similar karyotypes are given by GOEPFERT (1974), TZANOUDAKIS (1986), TAK & WAFI (1996), and BALTISBERGER & WIDMER (2004). As similar karyotypes have been found in plants from both colour types no cytological difference seems to exist between white or red coloured plants from the respective areas.

2. *Ranunculus creticus* ($2n = 16$; Fig. 1B)

Ranunculus creticus occurs in the S Aegean area and grows in shady places on rocky ledges and in gorges. It is a distinct, beautiful and often rather tall species. As it grows in similar habitats and looks like the even larger *R. cortusifolius* Willd. from the Macaronesian islands it was suggested that these two species are closely related (GOEPFERT, 1976). But molecular data suggest (PAUN & al., 2005) that the two species are not as closely related. They join the same clade but they are each nested within a group of more or less the same phytogeographical region (W- respectively E-Mediterranean). The similar habit is probably an independent adaptation to the shaded habitat. Nevertheless, based on data of chloroplast DNA, JOHANNSON (1998) found a strongly supported monophyletic group with both *R. cortusifolius* and *R. creticus* as well as some other species (e.g. *R. asiaticus*).

All plants investigated from all four sites had 16 chromosomes and were thus diploid. This corresponds with GOEPFERT (1974), MONTMOLLIN (1986), and TZANOUDAKIS (1986). The karyotype consists of 6 metacentric, 8 submeta- to subtelocentric and 2 acrocentric chromosomes, the latter with satellites (6m 8sm/st 2a_{SAT}). Similar karyotypes are given by GOEPFERT (1974) and TZANOUDAKIS (1986). These findings for *R. creticus* correspond with the karyotype data for other *Ranunculus*-species, e.g. *R. asiaticus* (see above) and *R. cortusifolius* (GOEPFERT, 1974; DALGAARD, 1985; BALTISBERGER & al., 1990). These three species are not closely related (PAUN & al., 2005) but belong to a monophyletic group (JOHANNSON, 1998).

3. *Ranunculus cupreus* ($2n = 16$; Fig. 1C)

Ranunculus cupreus is an endemic species of Crete growing in crevices and ledges on limestone, often with phrygana. It is a small species (up to 15 cm) characterized by its sparsely pubescent leaves 3-partite to base with rather deeply 3-partite lobes and the honey-leaves yellow above but copper-coloured beneath. However, the plants on which the cytological investigations were based have basal leaves which are completely glabrous and more than normally lobed (see Fig. 2), and they lack the reddish tinge of the outside of the honey-leaves (fide Kit Tan). *Ranunculus cupreus* looks rather similar to *R. millefoliatus* Vahl differing in size, less divided leaves, and the copper-coloured honey-leaves. The close relationship of these two species is supported by molecular data (PAUN & al., 2005).

The chromosome number of *R. cupreus* was not known up to now. With 16 chromosomes the plants proved to be diploid. The same number is given for the related *R. millefoliatus* (GARBARI & al., 1973; POPOVA, 1973; GOEPFERT, 1974; MARCHI & VISONA, 1982; SHOPOVA & SEKOVSKI, 1982). The karyotype of *R. cupreus* consists of 3 pairs of metacentric, 3 pairs of meta- to submetacentric, 1 pair of subtelocentric and 1 pair of acrocentric chromosomes, the subtelo- and acrocentric chromosomes bearing satellites (6m 6m/sm 2st_{SAT} 2a_{SAT}). Similar karyotypes are given for *R. millefoliatus* by GOEPFERT (1974), MARCHI & VISONA (1982) and SHOPOVA & SEKOVSKI (1982).

4. *Ranunculus ficaria* ($2n = 32$)

Ranunculus ficaria is widespread on wet and shady places in Europe, Asia, and N Africa and introduced in northern America. It is very variable and taxonomically difficult, and several ploidy levels (viz. $2x$, $3x$, $4x$) as well as aneuploid individuals (HESS & al., 1977) are known. Based on morphological characters some previous authors (e.g. JANCHEN, 1949) suggested to treat *R. ficaria*

and its allied species as a separate genus *Ficaria* Schaeff. but in most floras it remained within *Ranunculus*. Nevertheless molecular data support the separation of *Ficaria* as a distinct genus (HÖRANDL & al., 2005; PAUN & al., 2005).

Plants from the Mediterranean region have been referred to *Ranunculus ficaria* subsp. *ficariiformis* (F. W. Schultz) Rouy & Foucaud. Greek plants normally lack bulbils in the leaf axils which is typical for this subspecies; they therefore have been recently described as a distinct tetraploid subspecies, *R. ficaria* subsp. *chrysocephalus* P. D. Sell (SELL, 1991), and this seems to be the only taxon of the group occurring in S Greece and the Aegean region (STRID, 2002). Plants of both sites from Crete proved to be tetraploid with $2n = 4x = 32$ chromosomes confirming the indication by SELL (1991).

5. *Ranunculus marginatus* ($2n = 16$)

Ranunculus marginatus grows in open vegetations on wet places, in abandoned fields and on roadsides. Its distribution is said to cover SE Europe. Two varieties are distinguished (TUTIN & AKEROYD, 1993). Typical *R. marginatus* with smooth achenes is diploid with $2n = 2x = 16$ chromosomes (STRID & FRANZEN, 1981; LENTINI & al., 1988; BALTISBERGER & al., 1993) whereas *R. marginatus* var. *trachycarpus* (Fisch. & C. A. Mey.) Azn. with tuberculate achenes is tetraploid (LANGLET, 1927; GREGSON, 1965; GOEPFERT, 1974; SLAVIK & al., 1993; BALTISBERGER & BALTISBERGER, 1995). As many annuals, *R. marginatus* varies greatly in size, branching mode, and number of flowers. But it also varies in taxonomically relevant morphological characters of the achenes such as size, broadness of the margin, length of the beak, and surface (smooth, variously tuberculate till even muricate). This is also the case with the plants from Crete showing achenes with the surface from completely smooth to densely tuberculate (Fig. 3). All plants investigated were diploid with $2n = 2x = 16$, even these plants with tuberculate achenes. So the correlation between ploidy level and variety seems to be nonexistent (at least for the material from Crete).

Ranunculus marginatus often looks similar to the annual, widespread and also very variable *R. sardous* Crantz, and the two taxa are therefore often difficult to distinguish. *R. sardous* is diploid or tetraploid, and the surface of the achenes is also variable as in *R. marginatus*. At least at present no geographical pattern based on morphology and/or cytology can be detected within this complex, and it therefore was proposed to treat the respective taxa as a variable complex under the name of *R. sardous* (STRID, 2002). But molecular data indicate that the two species are not (HÖRANDL & al., 2005) or at least not closely related (PAUN & al., 2005).

6. *Ranunculus muricatus* ($2n = 48$)

Ranunculus muricatus is a widespread annual in the Mediterranean area and introduced in other regions with similar climate. It grows in open vegetation on wet places, in abandoned fields and disturbed grounds. The species is polyploid, and three levels are known viz. tetraploid with 32 (VAN LOON & DE JONG, 1978), hexaploid with 48 (e.g. DIOSDADO & PASTOR, 1996), and octoploid with 64 chromosomes (FUJISHIMA, 1986). Based on ploidy levels no geographical pattern seems to exist. All investigated plants from all four sites from Crete proved to be hexaploid with $2n = 6x = 48$ chromosomes which is the most common ploidy level within *R. muricatus*.

7. *Ranunculus paludosus* ($2n = 32, 40$; Fig. 1D)

Ranunculus paludosus grows in dry or seasonally wet meadows, in abandoned fields and open places in bushland. It is a variable and widespread species in the Mediterranean area. Two ploidy levels are known viz. diploid with $2n = 2x = 16$ and tetraploid with $2n = 4x = 32$. The investigated plants from Krasion proved to be tetraploid, confirming the ploidy level more frequently indicated in literature. But all plants from the other two sites were pentaploid with $2n = 5x = 40$ which was unknown within this species up to now (Fig. 1D).

The pentaploid plants could originate by combining a reduced gamete of a diploid with a unreduced gamete of a tetraploid plant. But as diploid *R. paludosus* is not known from Crete the origin of the pentaploid plants remains unclear. Pentaploid plants show an uneven ploidy level. Complete pairing of chromosomes in meiosis is therefore not possible. The result is an unregular meiosis, and those plants are probably not fertile. But as all plants produce subterranean stolons (up to ten stolons per plant!) which grow very fast and build remarkable rosettes of leaves within some weeks (at least in the greenhouse) the propagation of these pentaploid plants seems to be no problem. This was probably also the case in nature because both sites were rather densely covered with numerous rosettes of this species. A similar strategy was described for plants of *R. dissectus* subsp. *sibthorpii* Davis with an uneven ploidy level (here triploid) from Ulu Dag in Turkey (BALTSBERGER & TAN, 1991). This taxon usually is diploid and without stolons but a population of triploid plants was found producing subterranean stolons.

ACKNOWLEDGEMENTS

We thank Prof. Dr. F. Klötzli (Zürich) and Dr. B. Merz (Geneva) for the *Ranunculi* sampled in Crete, and Dr. K. Tan (Copenhagen) for the support with *Ranunculus cupreus*. We are very grateful to A. Skoumalova (Prague) for a draft sketch of Fig. 2. We also thank an anonymous reviewer for critical comments on the manuscript.

REFERENCES

- BALTSBERGER, M. (1980). Die Artengruppe des *Ranunculus polyanthemos* L. in Europa. *Ber. Schweiz. Bot. Ges.* 90: 143-188.
- BALTSBERGER, M. (1992). Botanische Notizen und zytologische Untersuchungen an einigen Pflanzen (insbesondere aus den Gattungen *Ranunculus* und *Achillea*) aus dem albanisch-jugoslawischen Grenzgebiet (Korab, Sar Planina). *Ber. Geobot. Inst. E. T. H. Stiftung Rübel* 58: 192-211.
- BALTSBERGER, M. (1994). *Ranunculus cacuminis* and *R. crenatus*, representatives of the *R. alpestris*-group (Ranunculaceae) on the Balkan Peninsula. *Pl. Syst. Evol.* 190: 231-244.
- BALTSBERGER, M. & E. BALTSBERGER (1995). Cytological data of Albanian plants. *Candollea* 50: 457-493.
- BALTSBERGER, M., W. HUBER & B. MERZ (1990). Zytologische Untersuchungen an einigen Pflanzen von den Kanarischen Inseln. *Ber. Geobot. Inst. E. T. H. Stiftung Rübel* 56: 142-149.
- BALTSBERGER, M., A. MULLAJ & V. TARTARI (1993). Mediterranean chromosome number reports 3 (185-203). *Fl. Medit.* 3: 348-353.
- BALTSBERGER, M. & K. TAN (1991). Investigations in *Ranunculus dissectus* subsp. *sibthorpii* (Ranunculaceae). *Willdenowia* 20: 67-72.
- BALTSBERGER, M. & A. WIDMER (2004). Cytological data of some plants species from Israel. *Israel J. Plant Sci.* 52: 171-176.
- DALGAARD, V. (1985). Chromosome studies in flowering plants from Madeira. *Willdenowia* 15: 137-156.
- DAVIS, P. H. (1965). *Ranunculus* L. In: DAVIS, P. H. (ed.), *Fl. Turkey* 1: 146-197.
- DIOSDADO, J. C. & J. E. PASTOR (1996). Consideraciones citotaxonomicas del genero *Ranunculus* L. (Ranunculaceae) en la Peninsula Iberica. *Anales Jard. Bot. Madrid* 54: 166-178.
- DYER, A. F. (1963). The use of lacto-propionic orcein in rapid squash methods for chromosome preparations. *Stain Technol.* 38: 85-90.
- FEDEROV, A. (1974). *Chromosome numbers of flowering plants*. Koeltz.
- FUJISHIMA, H. (1986). A new chromosome number of *Ranunculus muricatus* L. *Kromosomo II* 43-44: 1367-1371.
- GARBARI, F., N. TORNADORE & E. PECORI (1973). Numeri cromosomici per la Flora Italiana: 151-161. *Inform. Bot. Ital.* 5: 161-169.
- GOEPFERT, D. (1974). Karyotypes and DNA content in species of *Ranunculus* L. and related genera. *Bot. Not.* 127: 464-489.
- GOEPFERT, D. (1976). Phylogenetic studies on *Ranunculus cortusifolius* Willd. (Ranunculaceae), a Macaronesian endemic species. *Bot. J. Linn. Soc.* 72: 161-170.
- GOLDBLATT, P. (1981). Index to plant chromosome numbers 1975-1978. *Monogr. Syst. Bot. Missouri Bot. Gard.* 5: 1-533.

- GOLDBLATT, P. (1984). Index to plant chromosome numbers 1979-1981. *Monogr. Syst. Bot. Missouri Bot. Gard.* 8: 1-427.
- GOLDBLATT, P. (1985). Index to plant chromosome numbers 1982-1983. *Monogr. Syst. Bot. Missouri Bot. Gard.* 13: 1-224.
- GOLDBLATT, P. (1988). Index to plant chromosome numbers 1984-1985. *Monogr. Syst. Bot. Missouri Bot. Gard.* 23: 1-264.
- GOLDBLATT, P. & D. E. JOHNSON (1990). Index to plant chromosome numbers 1986-1987. *Monogr. Syst. Bot. Missouri Bot. Gard.* 30: 1-243.
- GOLDBLATT, P. & D. E. JOHNSON (1991). Index to plant chromosome numbers 1988-1989. *Monogr. Syst. Bot. Missouri Bot. Gard.* 40: 1-238.
- GOLDBLATT, P. & D. E. JOHNSON (1994). Index to plant chromosome numbers 1990-1991. *Monogr. Syst. Bot. Missouri Bot. Gard.* 51: 1-267.
- GOLDBLATT, P. & D. E. JOHNSON (1996). Index to plant chromosome numbers 1992-1993. *Monogr. Syst. Bot. Missouri Bot. Gard.* 58: 1-276.
- GOLDBLATT, P. & D. E. JOHNSON (1998). Index to plant chromosome numbers 1994-1995. *Monogr. Syst. Bot. Missouri Bot. Gard.* 69: 1-208.
- GOLDBLATT, P. & D. E. JOHNSON (2000). Index to plant chromosome numbers 1996-1997. *Monogr. Syst. Bot. Missouri Bot. Gard.* 81: 1-188.
- GOLDBLATT, P. & D. E. JOHNSON (2003). Index to plant chromosome numbers 1998-2000. *Monogr. Syst. Bot. Missouri Bot. Gard.* 94: 1-297.
- GREGSON, N. M. (1965). *Chromosome morphology and cytogenetics in the genus Ranunculus L.* PhD thesis. University of Liverpool.
- HESS, H. E., E. LANDOLT & R. HIRZEL (1977). *Fl. Schweiz* ed. 2, 2. Birkhäuser Verlag.
- HÖRANDL, E., O. PAUN, J. T. JOHANSSON, C. LEHNEBACH, T. ARMSTRONG, L. CHEN & P. LOCKHART (2005). Phylogenetic relationships and evolutionary traits in *Ranunculus* s.l. (Ranunculaceae) inferred from ITS sequence analysis. *Molec. Phylogen. Evol.* 36: 305-327.
- JANCHEN, E. (1949). Die systematische Gliederung der Ranunculaceen und Berberidaceen. *Denkschr. Österr. Akad. Wiss., Math.-Naturwiss. Kl.* 108: 1-82.
- JOHANSSON, J. T. (1998). Chloroplast DNA restriction site mapping and the phylogeny of *Ranunculus* (Ranunculaceae). *Pl. Syst. Evol.* 213: 1-19.
- LANGLET, O. F. I. (1927). Beiträge zur Zytologie der Ranunculaceen. *Svensk Bot. Tidskr.* 21: 1-17.
- LENTINI, F., S. ROMANO & F. M. RAIMONDO (1988). Numeri cromosomici per la Flora Italiana: 1185-1196. *Inform. Bot. Ital.* 20: 637-646.
- LEVAN, A., K. FREDGA & A. A. SANDBERG (1964). Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201-220.
- MARCHI, P. & L. VISONA (1982). Numeri cromosomici per la Flora Italiana: 889-905. *Inform. Bot. Ital.* 14: 248-258.
- MONTMOLLIN, B. DE (1986). Etude cytotaxonomique de la flore de la Crète. III. Nombres chromosomiques. *Candollea* 41: 431-439.
- MOORE, R. J. (1970). Index to plant chromosome numbers 1968. *Regnum Veg.* 68: 1-115.
- MOORE, R. J. (1971). Index to plant chromosome numbers 1969. *Regnum Veg.* 77: 1-112.
- MOORE, R. J. (1972). Index to plant chromosome numbers 1970. *Regnum Veg.* 90: 1-134.
- MOORE, R. J. (1973). Index to plant chromosome numbers 1967-1971. *Regnum Veg.* 90: 1-539.
- MOORE, R. J. (1974). Index to plant chromosome numbers 1972. *Regnum Veg.* 91: 1-108.
- MOORE, R. J. (1977). Index to plant chromosome numbers 1973/74. *Regnum Veg.* 96: 1-257.
- MÜLLER, M. & M. BALTISBERGER (1984). Cytotaxonomische Untersuchungen in der Artengruppe des *Ranunculus alpestris* (Ranunculaceae). *Pl. Syst. Evol.* 145: 269-289.
- PAUN, O., C. LEHNEBACH, J. T. JOHANSSON, P. LOCKHART & E. HÖRANDL (2005). Phylogenetic relationships and biogeography of *Ranunculus* and allied genera (Ranunculaceae) in the Mediterranean region and in the European Alpine System. *Taxon* 54, in press.
- POPOVA, M. (1973). Cytotaxonomic investigation in genus *Ranunculus L.* *Izv. Bot. Inst. (Sofia)* 24: 233-239.
- SELL, P. D. (1991). A new subspecies of *Ranunculus ficaria L.* *Bot. J. Linn. Soc.* 106: 117-118.
- SHOPOVA, M. & Z. SEKOVSKI (1982). *Chromosome atlas of some Macedonian angiosperms.* Macedonian Academy of Sciences and Arts.

- SLAVIK, B., V. JAROLIMOVA & J. CHRTEK (1993). Chromosome counts of some plants from Cyprus. *Candollea* 48: 221-230.
- STRID, A. (2002). *Ranunculus* L. In: STRID, A. & K. TAN (ed.), *Fl. Hellen.* 2: 38-69.
- STRID, A. & R. FRANZEN (1981). Chromosome numbers. In: LÖVE, A. (ed.), Chromosome number reports LXXIII. *Taxon* 30: 829-842.
- TAK, M. A. & B. A. WAFAI (1996). Somatic chromosome structure and nucleolar organization in *Anemone coronaria* L., *Ranunculus asiaticus* L. and *Eranthis hyemalis* Salisb. (Ranunculaceae). *Phytomorphology* 46: 377-385.
- TUTIN, T. G. & J. R. AKEROYD (1993). *Ranunculus* L. In: TUTIN, T. G. & al. (ed.), *Fl. Eur.* 1, ed. 2: 269-286.
- TZANOUDAKIS, D. (1986). Chromosome studies in the Greek flora. I. Karyotypes of some Aegean Angiosperms. *Bot. Helv.* 96: 27-36.
- VAN LOON, J. C. (1987). *A cytotaxonomical atlas of the Balkan flora*. J. Cramer.
- VAN LOON, J. C. & H. DE JONG (1978). Chromosome numbers. In: LÖVE, A. (ed.), Chromosome number reports LIX. *Taxon* 27: 56-60.

Submitted on March 8, 2005
Accepted on October 31, 2005

Table 1. – Alphabetical list of investigated species (* first record, + new ploidy level).

Species	Voucher number	Number of individuals investigated	2n
<i>Ranunculus asiaticus</i>	13230	1	16
<i>R. asiaticus</i>	13545	9	16
<i>R. asiaticus</i>	13740	5	16
<i>R. asiaticus</i>	13744	9	16
<i>R. asiaticus</i>	13741	9	16
<i>R. creticus</i>	12793	9	16
<i>R. creticus</i>	13555	12	16
<i>R. creticus</i>	13559	8	16
<i>R. creticus</i>	13561	10	16
<i>R. cupreus</i>	13558	10	16*
<i>R. ficaria</i>	13557	13	32
<i>R. ficaria</i>	13560	6	32
<i>R. marginatus</i>	13717	6	16
<i>R. muricatus</i>	13718	7	48
<i>R. muricatus</i>	13720	15	48
<i>R. muricatus</i>	13723	4	48
<i>R. muricatus</i>	13725	9	48
<i>R. paludosus</i>	13231	3	32
<i>R. paludosus</i>	13814	14	40+
<i>R. paludosus</i>	13815	13	40




Fig. 1. – Somatic metaphases. **A.** *Ranunculus asiaticus* L. (M. Baltisberger & A. Widmer 13545; $2n = 2x = 16$); **B.** *Ranunculus creticus* L. (M. Baltisberger & A. Widmer 13561; $2n = 2x = 16$); **C.** *Ranunculus cupreus* Boiss. & Heldr. (M. Baltisberger & A. Widmer 13558; $2n = 2x = 16$); **D.** *Ranunculus paludosus* Poir. (M. Baltisberger 13815; $2n = 5x = 40$).




Fig. 2. – *Ranunculus cupreus* Boiss & Heldr. (M. Baltisberger & A. Widmer 13558). A. Habit; B. Basal leaf.




Fig. 3. – Achenes of *Ranunculus marginatus* d'Urv. (M. Baltisberger 13717) showing the surface variability from tuberculate (left) to smooth (right).