

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 61 (2006)

Heft: 1

Artikel: Historical and functional aspects of plant biodiversity : an example on the flora of the Vukova Gorica region (Central Croatia)

Autor: Alegro, Antun / Markovi, Ljerka / Antoni, Oleg

DOI: <https://doi.org/10.5169/seals-879246>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Historical and functional aspects of plant biodiversity – an example on the flora of the Vukova Gorica region (Central Croatia)

ANTUN ALEGRO,
LJERKA MARKOVIĆ,
OLEG ANTONIĆ
&
SANDRO BOGDANOVIĆ

ABSTRACT

ALEGRO, A., LJ. MARKOVIĆ, O. ANTONIĆ & S. BOGDANOVIĆ (2006). Historical and functional aspects of plant biodiversity: the example of the flora of the Vukova Gorica region (Central Croatia). *Candollea* 61: 135-166. In English, English and French abstracts.

Vukova Gorica region, located in the Western part of Central Croatia and displaying an area of 3.2 km², has a vascular flora of 531 taxa from 96 families. Seven main habitat types were recognised. For each type of habitat, biological and chorological spectra have been calculated, as well as a cross-spectrum representing the relationship between the chorological types and the life forms. The spectra were analysed using correspondence analysis (CA). The present state of diversity can be considered as the result of the natural and historical circumstances and the extensive anthropogenic impact. It is concluded that the human impact was essential for the richness of the flora and its chorological structure. The high α -diversity index recorded on the studied area shows that Vukova Gorica region is one of the floristically richest region of Croatia.

RÉSUMÉ

ALEGRO, A., LJ. MARKOVIĆ, O. ANTONIĆ & S. BOGDANOVIĆ (2006). Aspects historiques et fonctionnels de la biodiversité des plantes: l'exemple de la flore de la région de Vukova Gorica (Croatie centrale). *Candollea* 61: 135-166. En anglais, résumés anglais et français.

La région de Vukova Gorica se situe dans la partie occidentale du centre de la Croatie et comprend une aire de 3,2 km². Sa flore vasculaire est composée de 96 familles et de 531 taxa. Des spectres biologiques et chorologiques ont été calculés pour chaque type d'habitat de même qu'un spectre croisé représentant la relation entre les types chorologiques et les formes vivantes. Les spectres ont été analysés selon l'analyse de correspondance (AC). L'état actuel de la diversité peut être considéré comme étant le résultat des circonstances naturelles et historiques et d'un impact anthropogénique extensif. Il en est conclu que l'impact humain est essentiel pour la richesse et la structure chorologique de la flore. L'index de diversité- α élevé de la région étudiée montre que la région florale de Vukova Gorica est l'une des plus riches de Croatie.

KEY-WORDS: flora – diversity – chorological types – life forms – cross-spectrum – Vukova Gorica region – Croatia

Introduction

Biodiversity studies have often been considered only in terms of species richness for a given area. Species richness, however, is only one aspect of biodiversity. The complex regional patterns of species assemblages may be originated from the operation and interaction of historical, causal and functional factors (BISBY, 1995). A statistical approach to floristic data can provide some explanations for the patterns of distribution of different groups of plant species.

The historical and functional aspects of plant biodiversity are considered in this paper where (1) we describe the vascular flora of the studied area with the species richness and the main habitats where species are found, (2) we quantify the relationships between the phytogeographical affinities of species, reflecting their history, and their distribution within the habitat types, (3) we quantify the relationships between species life forms and their distribution within the habitat types, (4) we distribute the species in groups with same life form and chorotype in order to analyse the flora from a biogeographical and structural point of view, and (5) we compare the flora richness of the studied area with the other parts of Croatia using α -diversity index.

This is the first attempt to evaluate the flora of some part of Croatia by the simultaneous interconnection of chorological, life-form and habitat data. The flora of the continental part of Croatia has not been systematically studied until now. Furthermore, the analyses of a few areas provide independent presentations of chorological and life-form spectra, regardless to the habitat types (cf. ŠEGULJA, 1977; HULINA, 1989; STANČIĆ, 1994; ŠEGULJA & al., 1998; ŠOŠTARIĆ & MARKOVIĆ, 1998; REGULA-BEVILACQUA & ŠEGULJA, 2000). Similar approach of describing biodiversity by species richness and basic spectra was commonly used in a majority of traditional analyses of the floras around the world. Some of the first consistent and detailed statistical approaches to study associations between life-forms, phytogeographical elements and the distribution of species within the habitats have been carried out for the flora of Terra Nova National Park (Newfoundland, Canada) (CHAREST & al., 2000) and for the flora of Euganean Hills (North-Eastern Italy) (VILLANI & al., 2003).

Study area

The studied area of Vukova Gorica (Fig. 1) is located in Central Croatia and belongs to the West-Pannonian macroregion, spreading alongside the Kupa river (i.e. the border with the Republic of Slovenia) in a total length of 5 km. The area is bordered by the Kupa river on the North and North-West and roads on the South and South-East. The total surface of the area is 3.2 km². The elevation ranges from 149.5 to 255.6 m. The area includes the following small villages Johi, Glavica, Vukova Gorica and Gornje Prilišće. According to the climate classification by Köppen (BERTOVIĆ, 1975), it belongs to the temperate C climate and is located on the "continentally line", i.e. on the boundary between continental and maritime annual precipitation regime (FILIPČIĆ, 1996). Therefore, the precipitations are evenly distributed throughout the year with 680 mm during the vegetation period (April-October), and this represents 51% of the total annual amount. The average yearly temperature is 9.7°C (16-17°C during the vegetation period). The coldest month is January (average temperature of -1.6°C) and the hottest is July (21.1°C). The area belongs to the Croatian karst with geological basis composed by Jurassic limestone, Jurassic and Cretaceous dolomites and alluvial deposits on some parts of the Kupa bank. These rocks are covered with several types of brown soils, relict terra rossa and alluvial soils above river deposits. The landscape is wavy, spotted with numerous shallow dolines (sinkholes). The special peculiarity of the region is a 60 m deep canyon of the Kupa river, exposed to the North, with vertical rocks in some parts.

After the Turkish wars, the area has been repopulated beginning in year 1554. In year 1805, there were 826 inhabitants, in 1935 approximately 2000 inhab. (in broader area), and in 1991 only 183 inhab. until now. The result of centuries long, but moderate human activity, is a mosaic landscape composed of a variety of habitats. Besides natural forest vegetation (oak-hornbeam, beech and gallery woods alongside the river), also shrubs, grasslands, meadows, and other rural habitats are present.

Materials and methods

The area has been floristically studied in the period between 1996 and 2004. The specimens, collected in all seasons, have been prepared for identification. Regarding the allochthonous species, only naturalised and invasive plants *sensu* RICHARDSON & al. (2000) were collected, and not the cultivated ones.

The habitat types have been recognised during the field work and later grouped into seven major types:

- B (□) – beech forest (association *Lamio orvalae-Fagetum* (Horvat 1938) Borhidi 1963), developed in the Kupa canyon;
- OH (▲) – oak-hornbeam forest (association *Epimedio-Carpinetum betuli* (Horvat 1938) Borhidi 1963), climazonal vegetation of the region;
- G (◇) – grasslands and meadows (predominantly different communities from alliance *Bromion erecti* (Koch 1926) Br.-Bl. 1936) and few patches of association *Arrhenatheretum elatioris* Br.-Bl. ex Scherrer 1925);
- M (+) – moist shrubs and gallery woods alongside the river with *Salix* spp., *Alnus* spp., *Populus nigra* and *Fraxinus excelsior* as dominant tree species;
- S (■) – vegetation of margins and shrubs of woodland edges and roadsides;
- W (◆) – weeds and ruderal habitats;
- A (B) – aquatic habitats (plants in the river).

Flora Europaea (TUTIN & al., 1964-1980, 1993) was mainly used to identify the plant material. In the case of problematic taxa, other floras and monothematic keys were considered (e.g. KOPP & SCHNEEBELI-GRAF, 1998; DELFORGE, 2001; HÖRANDL & al., 2002).

The nomenclature used follows NIKOLIĆ (1994, 1997, 2000). Habitat type, life-form and chorological type were attributed to each taxon of the check-list. Some taxa growing in two or more habitat types were equally considered for each habitat type in the further analyses.

Life-form categories of Raunkier's system (RAUNKIER, 1934), as presented by ELLENBERG & al. (1991) and OBERDORFER (1994), were accepted:

- P – Phanaerophytes;
- Ch – Chamaephytes;
- H – Hemicryptophytes;
- T – Therophytes;
- G – Geophytes;
- Hy – Hydrophytes.

If more than one life form was quoted for a single taxon, the most appropriate one was chosen considering field observations in the studied area.

Chorological types follow system proposed by HORVATIĆ (1963, 1967-68):

1. medit – Mediterranean (incl. submediterranean);
2. illyr-balc – Illyrian-Balkan;
3. S-europ – South-European;
4. atlant – Atlantic;
5. E-europ-pont – East-European-Pontic;
6. SE-europ – Southern-East-European;

7. C-europ – Central-European;
8. europ – European;
9. europ-asiat – Eurasiatic;
10. circ-holoarct – Circumholarctic;
11. cosmop – Cosmopolites;
12. neoph – Neophytes.

For the taxa not listed there, OBERDORFER (1994), PIGNATI (1982), SIMON & al. (1992), HORVAT (1929), HORVAT & al. (1974) were used. When more than one chorological type was given by these sources for a single taxon, the most appropriate was chosen and accorded with Horvatić's system. The biological and chorological spectra were prepared for the total flora and for each type of habitat. For the purposes of statistical analysis, the Atlantic chorological group was unified with the European group. In addition, the E-European-Pontic and SE-European group were also considered as one group. The reason for these agglomerations was the low number of species in the Atlantic group (one species) and in the SE-European group (three species), which can decrease the resolution of statistical analysis. The cross-spectra were prepared by considering the number of species of every chorological group and life form for each habitat type, and by considering the number of species of every life-form for each chorological group. In order to visualize the data structure, a correspondence analysis (CA) was carried out on the basis of the two-way contingency tables with the values of cross-spectra.

The results of CA analysis are presented in ordination diagrams, where groupings of variables are emphasized by arbitrary superimposed ellipses. However, the studied data set has three dimensions and analysis of all possible two-way marginal tables cannot substitute the analysis of whole three-way table (LEGENDRE & LEGENDRE, 1998). Therefore, log-linear contingency table analysis was performed. Large proportion of zero frequencies (number of species in particular combinations of life form, chorological type and habitat) in the cited three-way contingency table allowed computation of log-linear analysis impossible. To avoid this, all zero frequencies of these combinations were replaced by value 1 and other frequencies were multiplied by 100. Positive differences between observed species frequencies and those expected by chance for particular combinations of habitat type, chorological type and life-form are presented graphically. Log-linear analysis was also used for the testing of differences between observed and expected number of species in two-way combinations of habitat and chorological types (regardless of the life forms), as well as habitat types and life forms (regardless of the chorological types). In order to compare the biodiversity of Vukova Gorica region with some other specific regions in Croatia, the logarithmic regression between area and number of species was computed (HOBOHM 2000a, 2000b) through all the regions. On the basis of the formula of the regression line, the α -diversity indices were computed and compared for all regions. Correspondence analysis was carried out using the PC-ORD 4.14 package (MCCUNE & MEFFORD, 1999), while log-linear analysis and regression were done using the Statistica 7 package (Stasoft, Inc.).

Results

In the course of the research, 531 taxa of vascular plants from 96 different families were found (Appendix 1). The largest group are dicots (*Magnoliatae*) with 410 taxa from 72 families or 77.2% of total flora found. Monocots (*Liliatae*) are represented by 99 species from 11 families, followed by ferns (*Pteridophyta*) in 18 taxa, and finally by gymnosperms (*Coniferophytina*) in 4 taxa. The most numerous families (Table 1) are *Asteraceae* s.l., *Poaceae*, *Fabaceae*, *Lamiaceae*, and *Rosaceae*.

Table 1. – Families represented in the flora of Vukova Gorica with 10 or more taxa.

Family	No of species	% of total flora
<i>Asteraceae s.l.</i>	60	11.30
<i>Poaceae</i>	44	8.29
<i>Asteraceae s.s.</i>	38	7.16
<i>Fabaceae</i>	35	6.59
<i>Lamiaceae</i>	32	6.03
<i>Rosaceae</i>	23	4.33
<i>Cichoriaceae</i>	22	4.14
<i>Brassicaceae</i>	21	3.95
<i>Ranunculaceae</i>	21	3.95
<i>Scrophulariaceae</i>	20	3.77
<i>Apiaceae</i>	19	3.58
<i>Caryophyllaceae</i>	17	3.20
<i>Liliaceae</i>	16	3.01
<i>Cyperaceae</i>	14	2.64
<i>Polygonaceae</i>	10	1.88
<i>Orchidaceae</i>	10	1.88

137 taxa were found in the beech forests, 94 in the oak-hornbeam forests, 160 in the grasslands, 59 in the moist shrubs and gallery woods, 115 in the shrubs and the vegetation of woodland edges, 57 in the weed and the ruderal vegetation, and 12 in the aquatic habitats.

The phytogeographical spectrum is presented in Table 2. The interrelationship between the habitat types and chorotypes given by CA (Fig. 2) shows grouping of the forest vegetation (B, OH) with the Illyrian-Balkan chorotype (I). The weed and ruderal vegetation (W) is closest to the neophytes (II). The aquatic habitats (A) have isolated position nearest to the Circumholarctic chorotype (III). The shrub and margin vegetation (S) is situated between SE-European and Mediterranean chorotype on one side, and neophytes on the other side (IV). The grasslands (G)

Table 2. – The percentages of chorological types in the habitat types of Vukova Gorica.

Chorotypes	Total No	Habitat types						
		B	OH	G	M	S	W	A
Medit	7	1.46	1.06	1.25	0.00	0.86	1.75	0.00
Illyr-balc	14	8.76	6.38	0.00	0.00	0.00	0.00	0.00
S-europ	56	13.14	12.77	15.63	3.39	5.17	3.51	7.69
SE-europ-pont	13	1.46	1.06	2.50	3.39	2.59	1.75	0.00
C-europ	48	15.33	15.96	8.75	5.08	5.17	0.00	0.00
Europ	50	9.49	11.70	10.00	15.25	7.76	7.02	0.00
Europ-asiat	164	30.66	34.04	32.50	38.98	37.07	15.79	7.69
Circ-holarct	45	8.76	7.45	6.88	15.25	3.45	3.51	38.46
Cosmop	116	10.95	8.51	22.50	15.25	32.76	50.88	46.15
Neoph	18	0.00	1.06	0.00	3.39	5.17	15.79	0.00

and the moist shrubs and gallery woods (M) have relatively central position, closest to the Central- and South-European, European and Eurasiatic chorotypes (V). Similar tendencies are also evident in results of log-linear analysis (Table 3, Fig. 3). Weed and ruderal vegetation (◆) is characterised by high frequency of cosmopolites and neophytes. High frequency of cosmopolites and Eurasiatic plants in the shrub and margin vegetation (■) is evident, and that was not so clear in previous analysis. Connection of grassland vegetation (◇) with S-European plants is more emphasized than in CA. Furthermore, it is clear shown that frequencies of Illyrian-Balkan and C-European species are higher in beech forest (□) than in oak-hornbeam forest (▲).

Table 3. – Comparisons of observed species frequencies and those expected by chance using Chi-square test (as a part of log-linear analysis of frequency tables) (see main text).

Comparison	Chi-square	Degrees of freedom	P
Habitats vs. chorological types*	205.14	54	<0.01
Habitats vs. life-forms*	770.61	30	<0.01
Chorological types vs. life-forms (for habitat 1)	1538.43	45	<0.01
Chorological types vs. life-forms (for habitat 2)	3578.83	45	<0.01
Chorological types vs. life-forms (for habitat 3)	4243.39	45	<0.01
Chorological types vs. life-forms (for habitat 4)	2473.56	45	<0.01
Chorological types vs. life-forms (for habitat 5)	3274.17	45	<0.01
Chorological types vs. life-forms (for habitat 6)	3521.27	45	<0.01
Chorological types vs. life-forms (for habitat 7)	3012.68	45	<0.01

Table 4 presents spectrum of life-forms. The hemicryptophytes are the most represented group in the grassland vegetation, the geophytes and phanerophytes in the forest vegetation, the therophytes in the weed and ruderal vegetation. The hydrophytes are naturally present only in the water habitats. These relations are clearly confirmed by the log-linear analysis (Fig. 4).

Table 4. – The percentages of life forms in the habitat types of Vukova Gorica.

Life forme	Total No	Habitat types						
		B	OH	G	M	S	W	A
P	60	11.68	31.91	0.63	18.64	11.30	0.00	0.00
Ch	21	2.92	5.32	6.25	3.39	0.87	1.75	0.00
H	285	55.47	37.23	71.88	54.24	62.61	21.05	8.33
T	84	2.19	3.19	8.75	11.86	20.87	73.68	0.00
G	72	27.74	22.34	12.50	11.86	4.35	3.51	16.67
Hy	9	0.00	0.00	0.00	0.00	0.00	0.00	75.00

The cross-spectrum representing the interrelationship between the chorological types and life forms (Table 5) shows that phanerophytes, hemicryptophytes and therophytes are composed of all chorological groups. The most uniform are hydrophytes represented with Circumholarctic chorotype and cosmopolites. Given by CA, Figure 5 shows relationship between the therophytes and neophytes (I), the isolated position of hydrophytes closest to the Circumholarctic chorotype (II), the close relation between geophytes and South- and Central-European plants (III), and the almost central position of hemicryptophytes closest to Illyrian-Balkan, European and Eurasiatic chorotypes (IV). Chamaephytes are also isolated to some degree and closest to the Central-European plants (V). The position of the phanerophytes reflects their diversity with regard to chorological types.

Table 5. – Cross-spectrum representing percentages of chorological types in each life form.

Chorotypes	Total No	Life forms [%]					
		P	Ch	H	T	G	Hy
Medit	7	1.67	0.00	1.42	1.18	1.35	0.00
Illyr-balc	14	1.67	0.00	2.48	1.18	6.76	0.00
S-europ	56	10.00	19.05	12.06	2.35	13.51	0.00
SE-europ-pont	13	5.00	0.00	2.48	2.35	1.35	0.00
C-europ	48	16.67	28.57	7.09	1.18	13.51	0.00
Europ	49	11.67	19.05	8.87	8.24	9.46	0.00
Europ-asiat	164	33.33	19.05	35.82	20.00	31.08	0.00
Circ-holarct	45	5.00	4.76	9.93	2.35	9.46	44.44
Cosmop	116	10.00	9.52	18.79	47.06	13.51	55.56
Neoph	18	5.00	0.00	1.06	14.12	0.00	0.00

The results of the log-linear analysis of the three-way contingency table are presented in Figure 6. The relations with the most outstanding frequencies important for the explanation of the structure of the flora are:

- Eurasiatic hemicryptophytes in grassland vegetation (◇);
- Complex structure of the vegetation of margins and shrubs (■) with cosmopolitan and neophytic therophytes;
- European and S-European hemicryptophytes.

High frequency of Eurasiatic and Illyrian-Balkan geophytes is characteristic for beech wood (□), and this latter also for oak-hornbeam forest (▲). However, for forest vegetation, important combinations of Central-European, European phanerophytes and S-European hemicryptophytes are occurring, the latter especially for beech forest. Weed and ruderal vegetation (◆) is characterised by cosmopolitan and neophytic therophytes. Aquatic vegetation (B) is characterised by hydrophytes of cosmopolitan and circumholarctic distribution. In the vegetation of moist shrubs and gallery woods (+), high frequencies have European phanerophytes and hemicryptophytes, as well as circumholarctic hemicryptophytes.

The logarithmic regression between the number of species and areas of nine floristically well studied specific regions, in the non Eumediterranean parts of Croatia, is shown (Fig. 7). On the basis of the regression formula, α -diversity indices have been calculated (Table 6). The Vukova Gorica area has comparatively a high α -diversity index, and therefore it can be considered as one of the floristically richest region of Croatia.

Table 6. – Surfaces, numbers of species and α -diversity indices of some areas in Croatia.

Areas	area/km ²	No of species	α -diversity index
Papuk	336.00	1500	0.0682
Vukova Gorica	3.20	530	0.0487
Učka	160.00	1200	0.0402
Krapinske toplice	3.00	492	0.0224
Risnjak	64.10	936	0.0173
Medvednica	228.26	1202	0.0079
Plitvice	294.82	1267	0.0070
Paklenica	102.00	809	-0.0892
Konjščina	32.00	584	-0.1230

Discussion

The great floristic diversity of the studied area cannot only be explained by the moderate climate and the specific geographical position between the South- and Central-European floristic regions, but also by habitat diversity formed as a result of extensive anthropogenic influence.

According to the map of the natural potential vegetation of Croatia (TRINAJSTIĆ, 1987), the studied area should be covered with oak-hornbeam (OH) and beech (B) forests, as it was in the past for the whole inland area of Croatia (ŠOŠTARIĆ, 2004). In both types of natural climazonal vegetation together, 177 species were found. If we add the species of the water habitats, tree, shrub and some herbaceous species of gallery vegetation of the river banks, as well as some species of woodland clearings, the total number of taxa is around 250. In comparison to the 531 taxa of real recent flora, it is clear that the majority of the species is living on the habitats created by a moderate human activity during the past centuries. A similar doubling of species number can be read out from the graphs presented by KORNECK & al. (1998) for the flora of Germany and by SUKOPP & TREPL (1987) for the flora of Central Europe. The history of flora can be read out from the spectra of chorological types. Hence, endemic Illyrian-Balkan plants are strictly limited to the beech and oak-hornbeam forests, which is also shown in log-linear analysis. This indicates that forests are in the area the oldest vegetation type, which has persisted long enough to create and maintain suitable conditions for specific plant species. Some of the plant species growing there (e.g. *Epimedium alpinum*, *Lamium orvala*, *Hacquetia epipactis*, *Omphalodes verna*, *Cardamine enneaphyllos*) can be considered as relicts of the old Tertiary flora extinct in Central Europe during the Ice Ages. This shows more or less continuous development of the forest vegetation in South-European refugia. Furthermore, ŠOŠTARIĆ (2004) states that during the postglacial periods, up to the Subatlantic, there was no natural or anthropogenic devastation resulting in the loss of forest cover. As a result, the beech woods of Croatian part of west Balkan are floristically the richest ones in the whole of its area of distribution in Europe (HORVAT, 1949, 1962; HORVAT & al., 1974). Furthermore, relatively high portions of S-European and Central-European plants (especially represented by geophytes) are also significant for the flora of forests, indicating the transitional position of the studied area between the two phytogeographical regions.

Grasslands (G) are seminatural habitats formed during prehistorical and historical times by timbering and burning the woodland in order to obtain room and food for live stocks (cf. BONN & POSCHOLD, 1998). According to ŠOŠTARIĆ (2004), anthropogenic impacts on the forest vegetation in the continental part of Croatia has beginning early in the Neolithic, and the development of grasslands for animal grazing started perhaps between the Bronze Age until the Iron Age. However, these processes have given traces in their recent flora; Illyrian-Balkan plants lack completely, and the total amount of cosmopolites is higher than in wood vegetation. Grasslands have central position between S-European, Central-European, European and Eurasiatic plants. The higher frequency of S-European plants, which corresponds with geographical position of the area, is pointed in log-linear analysis.

Both the wood and grassland vegetations are free from neophytes. This can be explained by the "immunity" of natural, undisturbed plant communities against hemerochoric species, or by the immigration conditions as discussed by SUKOPP & TREPL (1987).

The vegetation of the margins and shrubs of the woodland edges and the roadsides (S) has always been under strong and constant human impact. As a consequence, the flora is dominated by Eurasiatic plants and cosmopolites. In Figure 2, this vegetation type is located between SE-European-Pontic and Mediterranean plants on one side, and neophytes on the other. Such position indicates that these habitats are relatively warm and dry, enabling the existence of the plants originating from South and South-East, but some degree of instability has allowed the colonisation of neophytes. Results of log-linear analysis show higher frequencies of therophytic cosmopolites and therophytic neophytes in this type of vegetation. However, this tendency is even more prominent in the weed and ruderal vegetation (W). Constant and profound anthropogenic disturbance through tith of gardens, orchards, vineyards, crops, live stock breeding and similar activities, has given a flora characterised by a high amount of therophytic neophytes and cosmopolites.

Very specific habitats are those influenced by water. The aquatic habitats (A) are floristically the poorest and the simplest ones. The water is a main and dominant ecological factor and, therefore, the resulting flora has no special phytogeographical peculiarities. Circumholarctic species, besides the cosmopolites ones, are dominant. Moist shrubs and gallery woods alongside the river (M) comprise terrestrial and semiterrestrial (inundated) habitats and consequently are more complex. Mediterranean and Illyrian-Balkan plants lack, but the frequency of Eurasiatic and Circumholarctic hemicryptophytes and Eurasiatic phanerophytes is the highest among all terrestrial habitat types. This shows again the influence of water as an equalizing ecological factor.

The cross-spectrum shows that the geophytes, phanerophytes and hemicryptophytes are the closest group to many chorological groups. Therefore, plants belonging to these life forms are the main basis of diversity from chorological point of view. Hemicryptophytes, phanerophytes and therophytes are the most diverse life forms, composed of all chorological groups, but with the prevalence of Eurasiatic plants. The high share of European plants is significant for phanerophytes and especially for chamaephytes. Chamaephytes and geophytes are also characterised by high share of South- and Central-European plants. Therefore, these life forms, especially quantitatively superior geophytes, mainly cause the transitional phytogeographical position of the studied area as previously discussed. Therophytes are characterised, besides Eurasiatic group, by neophytes. As already presented, neophytes are connected mostly with the weed and ruderal vegetation. Therefore, this life form can also be an indication of human disturbance because this life form is typical for sinantropic and ruderal species. Hydrophytes, the plants which inhabit the Kupa river, are representatives of only two groups, Circumholarctic and cosmopolites. This also indicates that water flora does not reflect local phytogeographical peculiarities.

The α -diversity index of the studied area is high, and compared with areas of other specific regions in Croatia, it indicates that the Vukova Gorica region is one of the most floristically rich region of Croatia.

In conclusion, it can be stated that the flora of the studied area is the result of natural and historical circumstances and long yet extensive anthropogenic impact in one of the most diverse areas of Croatia.

Unfortunately, diminishing of the diversity of the flora is to be expected. The main threat to the present state of diversity is the loss of seminatural habitats. Grasslands, which are the richest habitat type in the studied area, are especially threatened. The process of grassland degradation started with the decline in population and substantial reduction in traditional methods of managing. Without cattle and regular mowing, grasslands gradually turned into shrubs and forests (RADOVIĆ, 2000), which are unsuitable habitats for most open habitat species.

REFERENCES

- BISBY, F. A. (1995). Characterization of biodiversity. In: HEYWOOD, V. H. & R. T. WATSON (ed.), *Global biodiversity assessment*: 21-107. Cambridge University Press.
- BERTOVIĆ, S. (1975). Prilog poznavanju odnosa klime i vegetacije u Hrvatskoj. Contribution to knowledge of relation between climate and vegetation in Croatia. *Prir. Istraž. Ser. Acta Biol.* 41: 89-216.
- BONN, S. & P. POSCHOLD (1998). *Ausbreitungsbiologie der Pflanzen Mitteleuropas*. Quelle & Meyer Verlag.
- CHAREST, R., L. BROUILLET, A. BOUCHARD & S. HAY (2000). The vascular flora of Terra Nova National Park, Newfoundland, Canada: a biodiversity analysis from a biogeographical and life form perspective. *Canad. J. Bot.* 78: 629-645.
- DELFORGE, P. (2001). *Guide des Orchidées d'Europe d'Afrique du Nord et du Proche-Orient*. 2nd édition. Delachaux et Niestlé.
- ELLENBERG, H., H. E. WEBER, R. DÜLL, V. WIRTH, W. WERNER, & D. PAULIBEN (1991). Zeigwerte von Pflanzen in Mitteleuropa. *Scripta Geobot.* 18: 1-248.

- FILIPČIĆ, A. (1996). Klima Hrvatske-Voda u atmosferi. *Hrvatsk. Zemljopis, Specijalno Izdanje* 1: 90-92.
- HOBOHM, C. (2000a). Biodiversität. Quelle & Meyer Verlag.
- HOBOHM, C. (2000b). Plant species diversity and endemism on islands and archipelagos with special reference to the Macaronesian Islands. *Flora* 195: 9-24.
- HÖRANDL, E., F. FLORINETH, & F. HADACEK (2002). *Weiden in Österreich und angrenzenden Gebieten. Eigenverlag des Arbeitsbereiches Ingenieurbiologie und Landschaftsbau*. Institut für Landschaftsplanung und Ingenieurbiologie. Universität für Bodenkultur Wien.
- HORVAT, I. (1929). Rasprostranjenje i prošlost mediteranskih, ilirskih i pontskih elemenata u flori sjeverne Hrvatske i Slovenije (Die Verbreitung und Geschichte der mediterranen, illyrischen und pontischen Florenelemente in Nordkroatien und Slovenien). *Acta Bot. Inst. Bot. Univ. Zagreb* 4: 1-34.
- HORVAT, I. (1949). *Nauka o biljnim zajednicama*. Nakladni zavod Hrvatske, Zagreb
- HORVAT, I. (1962). Die Vegetation Südosteuropas in klimatischem und bodenkundlichem Zusammenhang. *Mitt. Österr. Geogr. Ges.* 104: 136-160.
- HORVAT, I., V. GLAVAČ & H. ELLENBERG (1974). *Vegetation Südosteuropas*. G. Fischer Verlag.
- HORVATIĆ, S. (1963). Vegetacijska karta otoka Paga s općim pregledom vegetacijskih jedinica hrvatskog primorja. (Carte des groupements végétaux de l'île Nord-Adriatique de Pag avec un aperçu général des unités végétales du littoral Croate). *Prir. Istraž. Ser. Acta Biol.* 4: 1-187.
- HORVATIĆ, S., LJ. ILIJANIĆ & LJ. MARKOVIĆ-GOSPODARIĆ (1967-68). Biljni pokrov okoline Senja (Die Pflanzendecke der Umgebung von Senj). *Senjski zbornik* 3: 298-323
- HULINA, N. (1989). Prikaz i analiza flore na području Turopolja. *Acta Bot. Croat.* 48: 141-160.
- KOPP, E. & R. SCHNEEBELI-GRAF (1998). *Illustrierter Leitfaden zum Bestimmen der Farne und farnverwandten Pflanzen der Schweiz und angrenzender Gebiete*. SVF – Schweizerische Vereinigung der Farnfreunde.
- KORNECK, D., M. SCHNITTLER, F. KLINGENSTEIN, G. LUDWIG, M. TAKLA, U. BOHN & R. MAY (1998). Warum verarmt unsere Flora? Auswertung der Roten Liste der Farn- und Blütenpflanzen Deutschlands. *Schriftenreihe Vegetationsk.* 29: 299-444.
- LEGENDRE, P., L. LEGENDRE (1998). *Numerical Ecology*. 2nd ed. Elsevier.
- MCCUNE, B. & M. J. MEFFORD (1999). *PC-ORD. Multivariate Analysis of Ecological Data, Version 4*. MjM Software Design, Gleneden Beach, Oregon, USA.
- NIKOLIĆ, T. (ed.) (1994). Flora Croatica. Index Florae Croaticae. Pars 1. *Nat. Croatica* 3, *Suppl.* 2: 1-116.
- NIKOLIĆ, T. (ed.) (1997). Flora Croatica. Index Florae Croaticae. Pars 2. *Nat. Croatica* 6, *Suppl.* 1: 1-232.
- NIKOLIĆ, T. (ed.) (2000). Flora Croatica. Index Florae Croaticae. Pars 3. *Nat. Croatica* 9, *Suppl.* 1: 1-324.
- OBERDORFER, E. (1994). *Pflanzensoziologische Exkursionsflora*. E. Ulmer Verlag.
- PIGNATTI, S. (1982). *Fl. Ital.*: 1-3.
- RADOVIĆ, J. (ed.) (2000). *An overview of the state of biological and landscape diversity of Croatia*. Ministry of environmental protection and physical planning, Zagreb.
- RAUNKIER, C. (1934). *Life forms of plants and statistical plant geography*. Clarendon Press, Oxford.
- REGULA-BEVILACQUA, LJ. & N. ŠEGULJA (2000). Analyse der Flora des Gebirgszuges Strahinščica im Hrvatsko zagorje (Kroatien). *Acta Bot. Croat.* 59: 243-278.
- RICHARDSON, D. M., P. PYŠEK, M. REJMÁNEK, M. G. BARBOUR, F. D. PANETTA & C. J. WEST (2000). Naturalization and invasion of alien plants: concepts and definitions. *Diversity & Distrib.* 6: 93-107.
- ŠEGULJA, N. (1977). Analiza flore Vukomeričkih Gorica. *Biosistematika* 3: 45-59.
- ŠEGULJA, N., LJ. ILIJANIĆ & LJ. MARKOVIĆ (1998). Prikaz i analiza flore Zrinske gore (Flora of the Zrinska gora mountain (Croatia)). *Acta Bot. Croat.* 55/56: 65-99.
- SIMON, T., A. HORÁNSZKY, K. DOBLOYI, T. SZERDAHELYI & F. HORVÁTH (1992). A magyar edényes flóra értékelő táblázata. In: SIMON, T. & A. MAGYARORSZÁGI (ed.), *A magyarországi edényes flóra határozója*: 837-955. Nemzeti Tankönyvkiadó, Budapest.
- ŠOŠTARIĆ, R. (2004). The development of vegetation in the inland area of Croatia during the postglacial period. *Nat. Croatica* 13: 357-369.
- ŠOŠTARIĆ, R. & LJ. MARKOVIĆ (1998). The flora of Krapinske toplice (Croatia). *Acta Bot. Croat.* 55/56: 101-118.
- STANČIĆ, Z. (1994). Prikaz i analiza flore okolice Konjšćine (Hrvatska). *Acta Bot. Croat.* 53: 125-140.

- SUKOPP, H. & L. TREPL (1987). *Extinction and Naturalization of Plant Species as Related to Ecosystem Structure and Function*. Springer Verlag.
- TRINAJSTIĆ, I. (1987). Karta prirodne potencijalne vegetacije Jugoslavije 1: 1000000. *Glasn. Šumske Pokuse Posebno Izd.* 3: 269-276.
- TUTIN, T. G. & al. (1964-1980). *Fl. Eur.*: 1-5.
- TUTIN, T. G., N. A. BURGESS, A. O. CHATER, J. R. EDMONDSON, V. H. HEYWOOD, D. M. MOORE, D. H. VALENTINE, S. M. WALTERS, D. A. WEBB (ed.) (1993). *Fl. Eur.*: 1, ed. 2. Cambridge University Press, Cambridge.
- VILLANI, M., M. BRENTAN, A. TODARO, S. MARCHIORI & N. TORNADORE (2003). Statistical and ecological analysis for the evaluation of floristic diversity: the case of a volcanic complex in North-Eastern Italy (Colli Euganei, Padova, Italy). *Plant Biosyst.* 137: 293-304.

Submitted on April 29, 2005

Accepted on May 11, 2006

Appendix 1. – List of the taxa of vascular plants found in the Vukova Gorica region.

	Chorotype	Life form	Habitat
PTERIDOPHYTA			
Equisetaceae			
<i>Equisetum arvense</i> L.	10	H	W
<i>Equisetum hiemale</i> L.	10	H	M
<i>Equisetum sylvaticum</i> L.	10	H	B
Aspleniaceae			
<i>Asplenium ruta-muraria</i> L.	10	H	B,OH,S
<i>Asplenium scolopendrium</i> L.	10	H	B
<i>Asplenium trichomanes</i> L.	11	H	B,OH
Athyriaceae			
<i>Athyrium filix-femina</i> (L.) Roth	10	H	B,OH
<i>Cystopteris fragilis</i> (L.) Bernh.	11	H	B
<i>Matteuccia struthiopteris</i> (L.) Tod.	10	H	M
Dryopteridaceae			
<i>Dryopteris affinis</i> (Lowe) Fraser-Jenk.	8	H	B
<i>Dryopteris carthusiana</i> (Vill.) H. P. Fuchs	10	H	B
<i>Dryopteris dilatata</i> (Hoffm.) A. Gray	10	H	B
<i>Dryopteris filix-mas</i> (L.) Schott	11	H	B,OH
<i>Gymnocarpium robertianum</i> (Hoffm.) Newman	10	G	B
<i>Polystichum aculeatum</i> (L.) Roth	11	H	B
<i>Polystichum setiferum</i> (Forskål) Woyn.	10	H	B
Hypolepidaceae			
<i>Pteridium aquilinum</i> (L.) Kuhn	11	G	G
Polypodiaceae			
<i>Polypodium vulgare</i> L.	11	H	B,OH
SPERMATOPHYTA			
CONIFEROPHYTINA			
Cupressaceae			
<i>Juniperus communis</i> L.	10	P	G
Pinaceae			
<i>Abies alba</i> Mill.	3	P	B
<i>Pinus strobus</i> L.	12	P	OH
<i>Pinus sylvestris</i> L.	9	P	OH
MAGNOLIOPHYTINA			
Magnoliatae			
Aceraceae			
<i>Acer campestre</i> L.	8	P	B,OH
<i>Acer obtusatum</i> Willd.	2	P	OH
<i>Acer pseudoplatanus</i> L.	7	P	OH

	Chorotype	Life form	Habitat
Adoxaceae			
<i>Adoxa moschatellina</i> L.	10	G	OH
Amaranthaceae			
<i>Amaranthus hybridus</i> L.	11	T	W
<i>Amaranthus lividus</i> L.	11	T	W
<i>Amaranthus retroflexus</i> L.	11	P	W
Apiaceae			
<i>Aegopodium podagraria</i> L.	9	H	B,OH,M
<i>Aethusa cynapium</i> L.	9	T	S
<i>Angelica sylvestris</i> L.	9	H	G,M,S
<i>Anthriscus sylvestris</i> (L.) Hoffm.	9	H	B,OH
<i>Astrantia major</i> L.	8	H	B
<i>Carum carvi</i> L.	9	H	G
<i>Chaerophyllum hirsutum</i> L.	7	H	B
<i>Daucus carota</i> L.	9	H	G
<i>Eryngium amethystinum</i> L.	1	H	G
<i>Hacquetia epipactis</i> (Scop.) DC.	2	H	B
<i>Heracleum sphondylium</i> L.	9	H	G,M
<i>Pastinaca sativa</i> L.	9	H	G
<i>Peucedanum cervaria</i> (L.) Lapeyr.	5	H	G
<i>Peucedanum oreoselinum</i> (L.) Moench	8	H	G
<i>Pimpinella major</i> (L.) Huds.	8	H	S
<i>Pimpinella saxifraga</i> L.	9	H	G
<i>Sanicula europaea</i> L.	11	H	B,OH
<i>Selinum carvifolia</i> (L.) L.	9	H	S
<i>Torilis japonica</i> (Houtt.) DC.	9	H	S
Apocynaceae			
<i>Vinca minor</i> L.	7	Ch	OH
Araliaceae			
<i>Hedera helix</i> L.	7	P	B,OH
Aristolochiaceae			
<i>Asarum europaeum</i> L.	9	H	B,OH
Asteraceae			
<i>Achillea millefolium</i> L.	11	H	G
<i>Ambrosia artemisiifolia</i> L.	12	T	S,W
<i>Anthemis arvensis</i> L.	11	T	W
<i>Arctium lappa</i> L.	9	H	M
<i>Artemisia vulgaris</i> L.	11	Ch	S
<i>Bellis perennis</i> L.	7	H	G,S
<i>Buphthalmum salicifolium</i> L.	7	H	OH
<i>Carlina acaulis</i> L.	3	H	G
<i>Carlina vulgaris</i> L.	7	H	G

	Chorotype	Life form	Habitat
<i>Centaurea jacea</i> L.	9	H	G
<i>Centaurea macroptylon</i> Borbàs	8	H	G,S
<i>Centaurea nigrescens</i> Willd.	3	H	G,S
<i>Centaurea pannonica</i> (Heuff.) Simonk.	5	H	G
<i>Centaurea scabiosa</i> L.	9	H	G
<i>Cirsium arvense</i> (L.) Scop.	9	G	S,W
<i>Cirsium montanum</i> (Willd.) Spreng.	3	H	S
<i>Cirsium oleraceum</i> (L.) Scop.	9	H	M
<i>Cirsium vulgare</i> (Savi) Ten.	9	H	S
<i>Conyza canadensis</i> (L.) Cronquist	12	T	W
<i>Doronicum austriacum</i> Jacq.	3	H	B
<i>Erechtites hieracifolia</i> (L.) DC.	12	T	S
<i>Erigeron annuus</i> (L.) Pers.	12	H	W
<i>Eupatorium cannabinum</i> L.	9	H	S
<i>Galinsoga ciliata</i> (Raf.) S. F. Blake	12	T	W
<i>Galinsoga parviflora</i> Cav.	12	T	W
<i>Inula conyza</i> DC.	3	H	B
<i>Inula helenium</i> L.	12	H	S
<i>Inula salicina</i> L.	9	H	G
<i>Leucanthemum vulgare</i> Lam.	9	H	G
<i>Petasites hybridus</i> (L.) P. Gaertner, B. Mey. & Scherb.	9	G	M
<i>Pulicaria dysenterica</i> (L.) Bernh.	3	H	M
<i>Rudbeckia laciniata</i> L.	12	H	M
<i>Senecio erucifolius</i> L.	9	H	G
<i>Senecio nemoralis</i> subsp. <i>fuchsii</i> (C. C. Gmel.) Čelak.	7	H	B,M
<i>Senecio ovirensis</i> (Koch) DC.	7	H	B
<i>Serratula tinctoria</i> L.	9	G	OH
<i>Solidago virgaurea</i> L.	9	H	OH,G
<i>Telekia speciosa</i> (Schreb.) Baumg.	5	H	M
Balsaminaceae			
<i>Impatiens glandulifera</i> Royle	12	T	M
<i>Impatiens noli-tangere</i> L.	9	T	M
Berberidaceae			
<i>Berberis vulgaris</i> L.	9	P	OH
<i>Epimedium alpinum</i> L.	2	G	B,OH
Betulaceae			
<i>Alnus glutinosa</i> (L.) Gaertn.	9	P	M
<i>Alnus incana</i> (L.) Moench.	10	P	M
<i>Alnus x pubescens</i> Tausch	10	P	M
<i>Betula pendula</i> Roth	9	P	OH
Boraginaceae			
<i>Echium vulgare</i> L.	8	H	G,S
<i>Myosotis arvensis</i> (L.) Hill.	9	T	S
<i>Myosotis ramosissima</i> Rochel	9	T	W

	Chorotype	Life form	Habitat
<i>Myosotis scorpioides</i> L.	10	H	M
<i>Omphalodes verna</i> Moench	2	H	B
<i>Pulmonaria officinalis</i> L.	8	H	B,OH
<i>Symphytum officinale</i> L.	8	H	M,S
<i>Symphytum tuberosum</i> L.	3	G	B,OH
Brassicaceae			
<i>Alliaria petiolata</i> (M. Bieb.) Cavara & Grande	9	H	B
<i>Arabidopsis thaliana</i> (L.) Heynh.	11	T,H	W
<i>Arabis hirsuta</i> (L.) Scop.	11	H	S,W
<i>Armoracia rusticana</i> P. Gaertner, B. Mey. & Scherb	11	G	M
<i>Barbarea vulgaris</i> R. Br.	11	H	S
<i>Brassica rapa</i> L.	12	T	W
<i>Capsella bursa-pastoris</i> (L.) Medik.	11	T	S,W
<i>Cardamine amara</i> L.	9	H	M
<i>Cardamine bulbifera</i> (L.) Crantz	7	G	B
<i>Cardamine enneaphyllos</i> (L.) Crantz	3	G	B
<i>Cardamine flexuosa</i> With.	11	H	B
<i>Cardamine hirsuta</i> L.	11	T,H	S,W
<i>Cardamine impatiens</i> L.	9	H	B
<i>Cardamine pratensis</i> L.	10	H	G
<i>Cardamine trifolia</i> L.	2	G	B
<i>Cardamine waldsteinii</i> Dyer	2	H	B
<i>Cardaminopsis arenosa</i> (L.) Hayek	8	T	B
<i>Lunaria annua</i> L.	6	H	S
<i>Lunaria rediviva</i> L.	8	H	B
<i>Rorippa sylvestris</i> (L.) Besser	9	H	S
<i>Sinapis arvensis</i> L.	11	T	W
Campanulaceae			
<i>Campanula patula</i> L.	8	H	G
<i>Campanula persicifolia</i> L.	9	H	B
<i>Campanula trachelium</i> L.	9	H	B,OH,S
Cannabidaceae			
<i>Humulus lupulus</i> L.	9	H	M,S
Caprifoliaceae			
<i>Sambucus ebulus</i> L.	8	H	S
<i>Sambucus nigra</i> L.	8	P	S
<i>Viburnum opulus</i> L.	9	P	S
Carpinaceae			
<i>Carpinus betulus</i> L.	7	P	OH

	Chorotype	Life form	Habitat
Caryophyllaceae			
<i>Arenaria serpyllifolia</i> L.	11	T	W
<i>Cerastium glomeratum</i> Thuill.	11	T	W
<i>Cerastium sylvaticum</i> Waldst.	8	H	B,OH
<i>Cucubalus baccifer</i> L.	9	H	S
<i>Lychnis flos-cuculi</i> L.	9	H	G
<i>Minuartia viscosa</i> (Schreb.) Schinz & Thell	6	T,H	W
<i>Moehringia muscosa</i> L.	3	H	B
<i>Moehringia trinervia</i> (L.) Clairv.	9	B	
<i>Myosoton aquaticum</i> (L.) Moench	9	G	M
<i>Saponaria officinalis</i> L.	11	H	M
<i>Silene dioica</i> (L.) Clairv.	9	H	B
<i>Silene latifolia</i> subsp. <i>alba</i> (Mill.) Greuter & Burdet	9	T,H	S
<i>Silene vulgaris</i> subsp. <i>angustifolia</i> Hayek	3	Ch	G
<i>Stellaria graminea</i> L.	9	H	G
<i>Stellaria holostea</i> L.	9	Ch	B,OH
<i>Stellaria media</i> (L.) Vill.	11	T,H	W
<i>Stellaria nemorum</i> subsp. <i>glochidisperma</i> Murb.	7	H	B
Celastraceae			
<i>Euonymus europaeus</i> L.	9	P	OH,S
<i>Euonymus latifolius</i> (L.) Mill.	1	P	OH
<i>Euonymus verrucosus</i> Scop.	5	P	B
Chenopodiaceae			
<i>Chenopodium album</i> L.	11	T	W
<i>Chenopodium polyspermum</i> L.	9	T	W
Cichoriaceae			
<i>Aposeris foetida</i> (L.) Less.	3	H	B,OH
<i>Cichorium intybus</i> L.	11	H	G,S
<i>Crepis biennis</i> L.	7	H	G,S
<i>Crepis capillaris</i> (L.) Wallr.	11	T	G
<i>Hieracium laevigatum</i> agg.	10	H	OH
<i>Hieracium murorum</i> L.	9	H	B
<i>Hieracium piliferum</i> Hoppe	3	H	G
<i>Hieracium pilosella</i> L.	5	G	
<i>Hieracium sabaudum</i> L.	7	H	OH
<i>Hieracium virosum</i> Pall.	9	H	OH
<i>Hypochoeris radicata</i> L.	3	H	G
<i>Lactuca serriola</i> L.	11	H	W
<i>Lactuca virosa</i> L.	9	T,H	S
<i>Lapsana communis</i> L.	9	H	OH
<i>Leontodon autumnalis</i> L.	9	H	G
<i>Leontodon hispidus</i> L.	8	H	G
<i>Mycelis muralis</i> (L.) Dumort.	7	H	B
<i>Picris hieracioides</i> L.	9	H	S,W
<i>Sonchus asper</i> (L.) Hill	9	T	S

	Chorotype	Life form	Habitat
<i>Sonchus oleraceus</i> L.	11	T	S
<i>Taraxacum officinale</i> Weber	11	H	G,S
<i>Tussilago farfara</i> L.	9	H	S
Cistaceae			
<i>Helianthemum nummularium</i> (L.) Mill.	7	Ch	G
Convolvulaceae			
<i>Calystegia sepium</i> (L.) R. Br.	11	H	S,W
<i>Convolvulus arvensis</i> L.	11	H	S,W
Cornaceae			
<i>Cornus mas</i> L.	3	P	B,OH
<i>Cornus sanguinea</i> L.	8	P	B,OH
Corylaceae			
<i>Corylus avellana</i> L.	8	P	B,OH
Crassulaceae			
<i>Sedum sexangulare</i> L.	7	Ch	G
Cucurbitaceae			
<i>Echinocystis lobata</i> (Michx.) Torr. & A. Gray	12	T	M
Cuscutaceae			
<i>Cuscuta epithymum</i> (L.) L.	9	T	G
Dipsacaceae			
<i>Dipsacus fullonum</i> L.	11	H	S
<i>Knautia arvensis</i> (L.) Coult.	9	H	G
<i>Knautia drymeia</i> Heuff.	2	H	B,OH
<i>Scabiosa columbaria</i> L.	3	H	G
<i>Succisa pratensis</i> Moench	9	H	G
Ericaceae			
<i>Calluna vulgaris</i> (L.) Hull	4	Ch	G
Euphorbiaceae			
<i>Euphorbia brittingeri</i> Samp.	3	H	G
<i>Euphorbia carniolica</i> Jacq.	2	H	B
<i>Euphorbia cyparissias</i> L.	9	H	S
<i>Euphorbia dulcis</i> L.	7	G	B
<i>Euphorbia esula</i> L.	9	H	W
<i>Euphorbia exigua</i> L.	3	T	W
<i>Euphorbia helioscopia</i> L.	11	T	W
<i>Mercurialis perennis</i> L.	8	G	B,OH

	Chorotype	Life form	Habitat
Fabaceae			
<i>Anthyllis vulneraria</i> subsp. <i>polyphylla</i> (DC.) Nyman	7	H	G
<i>Astragalus glycyphyllos</i> L.	9	H	S
<i>Chamaecytisus supinus</i> (L.) Link	8	Ch	OH
<i>Chamaespartium sagittale</i> (L.) P. E. Gibbs	3	Ch	G
<i>Coronilla varia</i> L.	8	H	S
<i>Dorycnium pentaphyllum</i> subsp. <i>herbaceum</i> (Vill.) Rouy	3	H	G
<i>Genista germanica</i> L.	7	Ch	G
<i>Hippocrepis comosa</i> L.	3	H	G
<i>Lathyrus niger</i> (L.) Bernh.	8	G	OH
<i>Lathyrus pratensis</i> L.	9	H	S
<i>Lathyrus sylvestris</i> L.	8	H	S
<i>Lathyrus tuberosus</i> L.	9	H	G
<i>Lathyrus vernus</i> (L.) Bernh.	8	G	B
<i>Lotus corniculatus</i> L.	11	H	G
<i>Lotus uliginosus</i> Schkuhr	9	H	G
<i>Medicago lupulina</i> L.	11	T	G,S
<i>Medicago sativa</i> subsp. <i>falcata</i> (L.) Arcang.	9	H	M
<i>Medicago sativa</i> L. subsp. <i>sativa</i>	11	H	G,S
<i>Melilotus alba</i> Medik.	9	H	S
<i>Melilotus officinalis</i> (L.) Pall.	9	H	S
<i>Ononis spinosa</i> L.	8	Ch	G
<i>Robinia pseudacacia</i> L.	12	P	S
<i>Trifolium arvense</i> L.	9	T	G
<i>Trifolium campestre</i> Schreb.	11	T	G
<i>Trifolium fragiferum</i> subsp. <i>bonannii</i> (C. Presl) Soják	11	H	G
<i>Trifolium montanum</i> L.	9	H	G
<i>Trifolium patens</i> Schreb.	3	T	G
<i>Trifolium pratense</i> L.	9	H	G
<i>Trifolium repens</i> L.	11	H	G,S
<i>Vicia cassubica</i> L.	8	H	W
<i>Vicia cracca</i> L.	9	H	G,S
<i>Vicia dumetorum</i> L.	9	H	S
<i>Vicia sativa</i> subsp. <i>cordata</i> (Hoppe) Asch. & Graebn.	5	G	
<i>Vicia sepium</i> L.	9	H	M
<i>Vicia tetrasperma</i> (L.) Schreb.	9	T	G
Fagaceae			
<i>Castanea sativa</i> Mill.	3	P	OH
<i>Fagus sylvatica</i> L.	7	P	B
<i>Quercus cerris</i> L.	3	P	OH
<i>Quercus petraea</i> (Matt.) Liebl.	7	P	OH
Fumariaceae			
<i>Corydalis cava</i> (L.) Schweigg. & Körte	9	G	B
<i>Corydalis solidia</i> (L.) Clairv.	9	G	B
<i>Pseudofumaria alba</i> (Mill.) Lidén	1	H	B

	Chorotype	Life form	Habitat
Gentianaceae			
<i>Centaurium erythraea</i> Rafn.	11	T,H	G
<i>Gentiana asclepiadea</i> L.	3	H	B,OH
<i>Gentiana cruciata</i> L.	9	H	G
Geraniaceae			
<i>Geranium columbinum</i> L.	9	T	S
<i>Geranium phaeum</i> L.	3	H	B
<i>Geranium robertianum</i> L.	11	T,H	B,OH
<i>Geranium sanguineum</i> L.	3	H	G
Globulariaceae			
<i>Globularia punctata</i> Lapeyr.	3	H	G
Haloragaceae			
<i>Myriophyllum spicatum</i> L.	11	Hy	A
Hydrophyllaceae			
<i>Phacelia tanacetifolia</i> Benth.	12	T	W
Hypericaceae			
<i>Hypericum hirsutum</i> L.	11	H	S
<i>Hypericum humifusum</i> L.	11	T,H	S
<i>Hypericum perforatum</i> L.	11	H	G
Juglandaceae			
<i>Juglans regia</i> L.	12	P	OH
Lamiaceae			
<i>Ajuga genevensis</i> L.	9	H	G
<i>Ajuga reptans</i> L.	9	H	G,S
<i>Calamintha sylvatica</i> Bromf.	3	H	B
<i>Clinopodium vulgare</i> L.	11	H	G,S
<i>Galeopsis pubescens</i> Besser	7	T	M
<i>Galeopsis speciosa</i> Mill.	8	T	M
<i>Glechoma hederacea</i> L.	10	H	G,S
<i>Glechoma hirsuta</i> Waldst. & Kit.	3	H	B,OH
<i>Lamiaeum galeobdolon</i> (L.) Ehrend. & Polatschek	9	Ch	B,OH
<i>Lamium maculatum</i> L.	9	H	M
<i>Lamium orvala</i> L.	2	H	B
<i>Lamium purpureum</i> L.	9	T	W
<i>Lycopus europaeus</i> L. subsp. <i>europaeus</i>	9	H	M
<i>Lycopus europaeus</i> subsp. <i>mollis</i> (A. Kern.) Skalický	9	H	M
<i>Melissa officinalis</i> L.	1	H	S
<i>Melittis melissophyllum</i> L.	7	H	OH
<i>Mentha arvensis</i> L.	10	H	W
<i>Mentha longifolia</i> (L.) Huds.	11	H	S
<i>Origanum vulgare</i> L.	9	H	G

	Chorotype	Life form	Habitat
<i>Prunella laciniata</i> (L.) L.	3	H	G
<i>Prunella vulgaris</i> L.	11	H	G
<i>Salvia glutinosa</i> L.	9	H	B
<i>Salvia pratensis</i> L.	8	H	G
<i>Salvia verticillata</i> L.	3	H	S
<i>Saturea montana</i> L.	3	Ch	OH
<i>Scutellaria galericulata</i> L.	10	M	
<i>Stachys annua</i> (L.) L.	8	T	W
<i>Stachys officinalis</i> (L.) Trevis.	8	H	G
<i>Stachys palustris</i> L.	10	G	G
<i>Stachys sylvatica</i> L.	9	H	B,OH
<i>Teucrium chamaedrys</i> L.	3	Ch	G
<i>Thymus pulegioides</i> L.	9	Ch	G
Linaceae			
<i>Linum catharticum</i> L.	11	T	G
<i>Linum usitatissimum</i> L.	12	T	S
Lythraceae			
<i>Lythrum salicaria</i> L.	11	H	M
Malvaceae			
<i>Malva alcea</i> L.	3	H	S
<i>Malva sylvestris</i> L.	11	H	S
Oleaceae			
<i>Fraxinus ornus</i> L.	3	P	OH
<i>Ligustrum vulgare</i> L.	7	P	OH
<i>Fraxinus excelsior</i> L.	8	P	M
Onagraceae			
<i>Circaea lutetiana</i> L.	11	H	B
<i>Epilobium hirsutum</i> L.	9	H	S
<i>Epilobium parviflorum</i> Schreb.	9	H	S,W
<i>Epilobium tetragonum</i> subsp. <i>lamyi</i> (F. W. Schultz) Nyman	7	H	B
Oxalidaceae			
<i>Oxalis acetosella</i> L.	11	H,G	B,OH
<i>Oxalis stricta</i> L.	12	T	W
Papaveraceae			
<i>Chelidonium majus</i> L.	11	H	S
<i>Papaver rhoeas</i> L.	11	T	W
Plantaginaceae			
<i>Plantago lanceolata</i> L.	11	H	G
<i>Plantago major</i> L.	9	H	S
<i>Plantago media</i> L.	9	H	G

	Chorotype	Life form	Habitat
Polygalaceae			
<i>Polygala comosa</i> Schkuhr	9	H	G
<i>Polygala vulgaris</i> L.	9	H	G
Polygonaceae			
<i>Fallopia convolvulus</i> (L.) Á. Löve	11	T	M,S
<i>Polygonum aviculare</i> L.	11	T	S,W
<i>Polygonum lapathifolium</i> L.	11	T	M
<i>Polygonum mite</i> Schrank	8	T	M
<i>Polygonum persicaria</i> L.	11	T	W
<i>Rumex acetosa</i> L.	11	H	G
<i>Rumex acetosella</i> L.	11	H,G	G
<i>Rumex crispus</i> L.	11	H	G
<i>Rumex obtusifolius</i> L.	9	H	G
<i>Rumex patientia</i> L.	5	H	S
Portulacaceae			
<i>Portulaca oleracea</i> L.	11	T	W
Primulaceae			
<i>Anagalis arvensis</i> L.	11	T	W
<i>Cyclamen purpurascens</i> Mill.	2	G	B,OH
<i>Lysimachia nummularia</i> L.	8	Ch	W
<i>Lysimachia punctata</i> L.	5	H	S
<i>Lysimachia vulgaris</i> L.	9	H	S
<i>Primula vulgaris</i> Huds.	3	H	OH,G
Pyrolaceae			
<i>Monotropa hypopitys</i> L.	10	G	B
Ranunculaceae			
<i>Aconitum lycoctonum</i> subsp. <i>vulparia</i> (Rchb.) Nyman	9	H	B
<i>Aconitum variegatum</i> L.	7	H	B
<i>Actaea spicata</i> L.	9	H	B
<i>Anemone nemorosa</i> L.	11	G	B,OH
<i>Anemone ranunculoides</i> L.	9	G	B,OH
<i>Clematis vitalba</i> L.	7	P	B,OH,S
<i>Helleborus dumetorum</i> Waldst. & Kit.	7	G	B,OH
<i>Helleborus niger</i> subsp. <i>macranthus</i> (Freyn) Schiffn.	2	G	B,OH
<i>Hepatica nobilis</i> Schreb.	11	H	B
<i>Isopyrum thalictroides</i> L.	9	G	B
<i>Ranunculus acris</i> L.	11	H	G
<i>Ranunculus auricomus</i> L.	9	H	B
<i>Ranunculus bulbosus</i> L.	9	G	G
<i>Ranunculus ficaria</i> L.	8	H	M
<i>Ranunculus lanuginosus</i> L.	7	H	B
<i>Ranunculus repens</i> L.	11	H	G,S
<i>Ranunculus sardous</i> Crantz	11	H	G

	Chorotype	Life form	Habitat
<i>Ranunculus trichophyllus</i> Chaix	10	Hy	A
<i>Thalictrum aquilegifolium</i> L.	9	H	B
<i>Thalictrum lucidum</i> L.	8	H	M
<i>Thalictrum minus</i> L.	11	H	G
Rhamnaceae			
<i>Frangula alnus</i> Mill.	11	P	S
<i>Rhamnus catharticus</i> L.	9	P	S
Rosaceae			
<i>Agrimonia eupatoria</i> L.	10	H	G
<i>Aruncus dioicus</i> (Walter) Fernald	11	H	B
<i>Crataegus laevigata</i> (Poir.) DC.	8	P	B,OH
<i>Crataegus monogyna</i> Jacq.	9	P	B,OH
<i>Filipendula vulgaris</i> Moench	9	H	G
<i>Fragaria vesca</i> L.	11	H	B,OH
<i>Geum urbanum</i> L.	11	H	S
<i>Potentilla erecta</i> (L.) Raeusch.	9	H	G
<i>Potentilla heptaphylla</i> L.	7	H	G
<i>Potentilla micrantha</i> DC.	3	H	B
<i>Potentilla reptans</i> L.	11	H	G
<i>Prunus avium</i> L.	7	P	OH
<i>Prunus cerasifera</i> Ehrh.	5	P	OH
<i>Prunus fruticosa</i> Pall.	9	P	OH
<i>Prunus spinosa</i> L.	9	P	S
<i>Pyrus piraster</i> Burgsd.	9	P	B,OH
<i>Rosa arvensis</i> Huds.	7	Ch	B
<i>Rosa canina</i> L.	11	P	S
<i>Rubus bifrons</i> Tratt.	11	P	S
<i>Rubus caesius</i> L.	9	P	S
<i>Rubus hirtus</i> Waldst. & Kit.	7	P	S
<i>Sanguisorba minor</i> Scop.	9	H	G
<i>Sorbus torminalis</i> (L.) Crantz	9	P	B
Rubiaceae			
<i>Cruciata glabra</i> (L.) Ehrend.	3	H	G
<i>Cruciata laevipes</i> Opiz	3	H	G,S
<i>Galium album</i> Mill.	7	H	G
<i>Galium aparine</i> L.	11	T	S
<i>Galium mollugo</i> L.	9	H	G
<i>Galium odoratum</i> (L.) Scop.	9	G	B,OH
<i>Galium sylvaticum</i> L.	7	G	B
<i>Galium verum</i> L.	11	H	G

	Chorotype	Life form	Habitat
Salicaceae			
<i>Populus nigra</i> L.	11	P	M
<i>Populus tremula</i> L.	9	P	OH,S
<i>Salix alba</i> L. subsp. <i>alba</i>	9	P	M
<i>Salix alba</i> subsp. <i>vitellina</i> (L.) Arcang.	9	P	M
<i>Salix caprea</i> L.	9	P	OH,S
<i>Salix purpurea</i> L.	9	P	M
<i>Salix triandra</i> L.	9	P	M
Saxifragaceae			
<i>Chrysosplenium alternifolium</i> L.	10	H	B
<i>Saxifraga cuneifolia</i> L.	3	H	B
Scrophulariaceae			
<i>Chaenorhinum minus</i> (L.) Lange	8	T	W
<i>Cymbalaria muralis</i> P. Gaertner, B. Mey. & Scherb.	3	H	B,OH
<i>Euphrasia stricta</i> J. F. Lehm.	8	T	G
<i>Kickxia elatine</i> (L.) Dumort.	1	T	W
<i>Kickxia spurina</i> (L.) Dumort.	9	T	W
<i>Lathraea squamaria</i> L.	9	G	B,OH
<i>Linaria vulgaris</i> Mill.	9	G	G,S
<i>Melampyrum pratense</i> L.	9	T	OH
<i>Melampyrum velebeticum</i> Borbás	2	T	OH
<i>Rhinanthus minor</i> L.	10	T	G
<i>Scrophularia nodosa</i> L.	10	H	S
<i>Verbascum blattaria</i> L.	11	H,T	S
<i>Veronica agrestis</i> L.	9	T	W
<i>Veronica beccabunga</i> L.	10	Hy	A
<i>Veronica chamaedrys</i> L.	9	Ch	G
<i>Veronica montana</i> L.	3	H	B
<i>Veronica officinalis</i> L.	9	H	G
<i>Veronica persica</i> Poir.	11	T	W
<i>Veronica serpyllifolia</i> L.	11	H	G
<i>Veronica urticifolia</i> Jacq.	7	Ch	B
Solanaceae			
<i>Physalis alkekengi</i> L.	3	H	S
<i>Scopolia carniolica</i> Jacq.	2	H	B
<i>Solanum dulcamara</i> L.	11	Ch	M
<i>Solanum nigrum</i> L.	11	T	S
Staphyleaceae			
<i>Staphylea pinnata</i> L.	3	P	B
Thymeleaceae			
<i>Daphne mezereum</i> L.	9	P	B,OH

	Chorotype	Life form	Habitat
Tiliaceae			
<i>Tilia cordata</i> Mill.	8	P	OH
<i>Tilia platyphyllos</i> Scop.	7	P	OH
Ulmaceae			
<i>Ulmus minor</i> Mill.	5	P	M
<i>Ulmus glabra</i> Huds.	11	P	B
Urticaceae			
<i>Urtica dioica</i> L.	11	H	S
<i>Urtica urens</i> L.	11	T	S
Valerianaceae			
<i>Valeriana officinalis</i> L.	9	H	G,S
Verbenaceae			
<i>Verbena officinalis</i> L.	10	T,H	S
Violaceae			
<i>Viola alba</i> subsp. <i>scotophylla</i> (Jord.) Nyman	3	H	G
<i>Viola alba</i> Besser subsp. <i>alba</i>	3	H	G
<i>Viola arvensis</i> Murray	11	T	W
<i>Viola hirta</i> L.	9	H	G
<i>Viola odorata</i> L.	8	H	G,S
<i>Viola reichenbachiana</i> Boreau	7	H	B,OH
Vitaceae			
<i>Vitis vinifera</i> L.	11	P	S
MAGNOLIOPHYTINA			
Liliatae			
Alismataceae			
<i>Alisma plantago-aquatica</i> L.	11	Hy	A
Amaryllidaceae			
<i>Galanthus nivalis</i> L.	9	G	B,OH
<i>Leucojum vernum</i> L.	8	G	M
Araceae			
<i>Arum maculatum</i> L.	7	G	B
Cyperaceae			
<i>Carex buekii</i> Wimm.	8	G,Hy	M
<i>Carex caryophyllea</i> Latourr.	9	G	G
<i>Carex digitata</i> L.	9	H	B,OH
<i>Carex divulsa</i> subsp. <i>leersii</i> (Kneuck.) W. Koch	7	H	S
<i>Carex elata</i> All.	8	H	M
<i>Carex flacca</i> Schreb.	11	G	G,S

	Chorotype	Life form	Habitat
<i>Carex hirta</i> L.	9	H	M,S
<i>Carex pallescens</i> L.	10	H	G
<i>Carex pilosa</i> Scop.	9	H	B,OH
<i>Carex remota</i> L.	10	H	M
<i>Carex sylvatica</i> Huds.	8	H	B,OH
<i>Eleocharis palustris</i> (L.) Roem. & Schult.	11	A	
<i>Scirpus lacustris</i> L.	10	G,Hy	A
<i>Scirpus sylvaticus</i> L.	10	G	A
Dioscoreaceae			
<i>Tamus communis</i> L.	3	G	M
Iridaceae			
<i>Crocus vernus</i> (L.) Hill	3	G	G
Juncaceae			
<i>Juncus effusus</i> L.	9	H	S
<i>Juncus inflexus</i> L.	9	H	S
<i>Luzula pilosa</i> (L.) Wild.	10	H	B,OH
<i>Juncus compressus</i> Jacq.	11	G	A
<i>Juncus tenuis</i> Willd.	11	H	S
<i>Luzula campestris</i> (L.) Lam. & DC.	11	H	G
Liliaceae			
<i>Allium carinatum</i> L.	8	G	G
<i>Allium ursinum</i> L.	7	G	B,M
<i>Allium vineale</i> L.	7	G	G
<i>Anthericum ramosum</i> L.	7	H	G
<i>Asparagus tenuifolius</i> Lam.	3	G	OH
<i>Colchicum autumnale</i> L.	7	G	G,S
<i>Convallaria majalis</i> L.	10	G	OH
<i>Erythronium dens-canis</i> L.	2	G	B
<i>Gagea lutea</i> (L.) Ker-Gawl.	9	G	B
<i>Lilium martagon</i> L.	9	G	B,OH
<i>Ornithogalum pyramidale</i> L.	3	G	G
<i>Ornithogalum umbellatum</i> L.	3	G	G
<i>Paris quadrifolia</i> L.	9	G	B
<i>Polygonatum multiflorum</i> (L.) All.	10	G	B
<i>Scilla bifolia</i> L.	3	G	B
<i>Veratrum album</i> L.	9	H	M
Orchidaceae			
<i>Cephalanthera damasonium</i> (Mill.) Druce	3	G	G
<i>Dactylorhiza fuchsii</i> (Druce) Soó	9	G	B
<i>Dactylorhiza maculata</i> (L.) Soó	9	G	B
<i>Listera ovata</i> (L.) R. Br.	9	G	B,OH
<i>Neottia nidus-avis</i> (L.) Rich.	9	G	B,OH
<i>Orchis morio</i> L.	7	G	G

	Chorotype	Life form	Habitat
<i>Orchis simia</i> Lam.	1	G	G
<i>Orchis tridentata</i> Scop.	3	G	G
<i>Platanthera bifolia</i> (L.) Rich.	9	G	G
<i>Epipactis helleborine</i> (L.) Crantz	9	G	B
Poaceae			
<i>Agrostis capillaris</i> L.	10	H	G
<i>Agrostis stolonifera</i> L.	9	H	A
<i>Anthoxanthum odoratum</i> L.	9	H	G
<i>Arrhenatherum elatius</i> (L.) J. & C. Presl	8	H	G
<i>Brachypodium rupestre</i> (Host) Roem. & Schult.	11	H	G
<i>Brachypodium sylvaticum</i> (Huds.) P. Beauv.	9	H	B,OH
<i>Briza media</i> L.	9	H	G
<i>Bromus commutatus</i> Schrad.	8	T	G
<i>Bromus erectus</i> Huds.	11	H	G
<i>Bromus hordeaceus</i> L.	11	T	G
<i>Bromus ramosus</i> Huds.	9	H	B
<i>Bromus sterilis</i> L.	11	T	S
<i>Calamagrostis epigejos</i> (L.) Roth	8	G	G
<i>Calamagrostis varia</i> (Schrad.) Host	9	H	OH
<i>Cynodon dactylon</i> (L.) Pers.	11	G	S,W
<i>Cynosurus cristatus</i> L.	11	H	G
<i>Dactylis glomerata</i> L.	9	H	G,S
<i>Deschampsia caespitosa</i> (L.) P. Beauv.	11	H	M
<i>Dichanthium ischaemum</i> (L.) Roberty	3	H	W
<i>Digitaria sanguinalis</i> (L.) Scop.	11	T	W
<i>Echinochloa crus-galli</i> (L.) P. Beauv.	11	T	W
<i>Elymus repens</i> (L.) Gould	11	G	G,S
<i>Festuca gigantea</i> (L.) Vill.	9	H	B
<i>Festuca pratensis</i> Huds.	9	H	G
<i>Glyceria maxima</i> (Hartm.) Holmb.	9	H,Hy	M
<i>Glyceria plicata</i> (Fr.) Fr.	11	H,Hy	M
<i>Holcus lanatus</i> L.	9	H	G
<i>Koeleria pyramidata</i> (Lam.) P. Beauv.	7	H	G
<i>Lolium perenne</i> L.	8	H	G
<i>Melica nutans</i> L.	5	B	
<i>Melica uniflora</i> Retz.	7	G	B,OH
<i>Molinia caerulea</i> subsp. <i>arundinacea</i> (Schrank) H. Paul	10	H	M
<i>Panicum capillare</i> L.	12	T	W
<i>Phalaris arundinacea</i> L.	10	H,Hy	M
<i>Phleum pratense</i> L.	10	H	G
<i>Poa angustifolia</i> L.	11	H	G
<i>Poa annua</i> L.	11	T,H	S,W
<i>Poa palustris</i> L.	10	H	OH
<i>Poa pratensis</i> L.	10	H	G
<i>Poa trivialis</i> L.	9	H	G
<i>Sesleria tenuifolia</i> Schrad.	1	H	B

	Chorotype	Life form	Habitat
<i>Setaria pumilla</i> (Poir.) Schult.	11	T	W
<i>Sorghum halepense</i> (L.) Pers.	11	G	G
<i>Trisetum flavescens</i> (L.) P. Beauv.	10	H	G
Potamogetonaceae			
<i>Potamogeton lucens</i> L.	10	Hy	A
<i>Potamogeton natans</i> L.	11	Hy	A
<i>Potamogeton perfoliatus</i> L.	11	Hy	A




Fig. 1. – Geographical position of the study area.




Fig. 2. – Correspondence analysis (CA) explaining the relationships between chorological types and habitats.




Fig. 3. – Log-linear analysis explaining the differences between observed species frequencies and those expected by chance (according to log-linear analysis of frequency tables) for particular combinations of habitat type and chorological type (negative differences not shown). Symbols used (in brackets): B (□) – beech forest; OH (▲) – oak-hornbeam forest; G (◇) – grasslands and meadows; M (+) – moist shrubs and gallery woods; S (■) – margins and shrubs; W (◆) – weeds and ruderal habitats; and A (B) – aquatic habitats.




Fig. 4. – Log-linear analysis explaining the differences between observed species frequencies and those expected by chance (according to log-linear analysis of frequency tables) for particular combinations of habitat type and life-form (negative differences not shown). Symbols used (in brackets): B (□) – beech forest; OH (▲) – oak-hornbeam forest; G (◇) – grasslands and meadows; M (+) – moist shrubs and gallery woods; S (■) – margins and shrubs; W (◆) – weeds and ruderal habitats; and A (B) – aquatic habitats.




Fig. 5. – Correspondence analysis (CA) explaining the relationships between chorological types and life forms.




Fig. 6. – Log-linear analysis explaining the differences between observed species frequencies and those expected by chance (according to log-linear analysis of frequency tables) for particular combinations of habitat type, chorological type and life-form (negative differences not shown). Symbols used (in brackets): B (□) – beech forest; OH (▲) – oak-hornbeam forest; G (◇) – grasslands and meadows; M (+) – moist shrubs and gallery woods; S (■) – margins and shrubs; W (◆) – weeds and ruderal habitats; and A (B) – aquatic habitats.




Fig. 7. – Logarithmic regression between the number of species (S) and area (A in km²) of some specific regions in Croatia. The line follows equation: $\log S = 0,2139\log A + 2,5675$.