

<b>Zeitschrift:</b>	Candollea : journal international de botanique systématique = international journal of systematic botany
<b>Herausgeber:</b>	Conservatoire et Jardin botaniques de la Ville de Genève
<b>Band:</b>	61 (2006)
<b>Heft:</b>	2
<b>Artikel:</b>	Euploca rodaliae J. I. M. Melo & Semir : a new species of Euploca (Heliotropiaceae) from Brazil
<b>Autor:</b>	Melo, José Iranildo Miranda de / Semir, João
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-879259">https://doi.org/10.5169/seals-879259</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Euploca rodaliae J. I. M. Melo & Semir – a new species of Euploca (Heliotropiaceae) from Brazil

JOSÉ IRANILDO MIRANDA DE MELO  
&  
JOÃO SEMIR

## ABSTRACT

MELO, J. I. M. & J. SEMIR (2006). *Euploca rodaliae* J. I. M. Melo & Semir – a new species of *Euploca* (Heliotropiaceae) from Brazil. *Candollea* 61: 453-456. In English, English and French abstracts.

A new species of the genus *Euploca* (Heliotropiaceae) from the states of Pernambuco and Bahia in Brazil is described and illustrated. *Euploca rodaliae* J. I. M. Melo & Semir is characterized mainly by characters of the leaf blade associated to the floral structure.

## RÉSUMÉ

MELO, J. I. M. & J. SEMIR (2006). *Euploca rodaliae* J. I. M. Melo & Semir – une nouvelle espèce d'*Euploca* (Heliotropiaceae) pour le Brésil. *Candollea* 61: 453-456. En anglais, résumés anglais et français.

Une nouvelle espèce du genre *Euploca* (Heliotropiaceae) pour les états de Pernambuco et Bahia au Brésil est décrite et illustrée. *Euploca rodaliae* J. I. M. Melo & Semir est caractérisé principalement par ses caractères foliaires associés à des structures florales.

**KEY WORDS:** HELIOTROPIACEAE – *Euploca* – Northeast Brazil

## Introduction

*Euploca* was established by NUTTALL (1837) based on *E. convolvulacea* Nutt. It was included in *Heliotropium* by GRAY (1874) and lately reestablished by HILGER & DIANE (2003). This genus belongs to the family Heliotropiaceae Schrad. and comprises approximately 120 species distributed in tropical, subtropical and temperate regions, most of them concentrated in the dry tropical zones. It includes plants predominantly herbaceous, subshrubs or rarely shrubs, associated to scorioideoes inflorescences or single flowers, usually with white and yellow mouth or yellow corollas. The fruit is squizocarpic, dry, constituted by four mericarps denominated nutlets. This important character is broadly adopted since the first treatment focusing on *Euploca* (MELO & SALES, 2004).

*Euploca* is widely distributed in Brazil, including the North, Northeast, Western-Center, South-east and South regions (MELO & SALES, 2004) and it is found in coastal habitats, forests, and more frequently in caatinga, cerrado and campo rupestre vegetation, a few species being as weeds in cultures.

During the study of the Brazilian *Euploca* species, a taxonomic novelty, *Euploca rodaliae* J. I. M. Melo & Semir, was recorded from the states of Pernambuco and Bahia (Northeast Brazil). Its description is based on the terminology proposed by HARRIS & HARRIS (1994). For the foliar typification, the venation and the vestiture patterns, RIZZINI (1977), HICKEY (1973) and PAYNE (1978), respectively, are followed.

***Euploca rodaliae* J. I. M. Melo & Semir, spec. nova** (Fig. 1).

**Typus:** BRAZIL. Bahia: Rio-Bahia road, 10.III.1958, fl., fr., Andrade-Lima 58-2919 (holo-: PEUFR 462; iso-: IPA 11482).

*Species suffruticosa, foliis discoloribus, ovalibus, ovalo-lanceolatis vel raro spatulatis et obovatis, petiolo decurrenti; spicis terminalibus et axillaribus, dichotomis; calycis laciniis subcarnosis et alatis; ovario stipitato; stylo circa 1 mm longo.*

**Subshrub** 0.3-1.1 m high. **Stem** striate, branches cylindrical, scabrous, sericeous. **Leaves** alternate or rarely pseudo-opposite just in the internodes region, petiole decurrent, blade 1.3-3 × 0.5-1 cm, discolour, membranaceous to subcarnose, ovate, ovate-lanceolate, or rarely espatulate to obovate, apices mucronate to acute, base truncate, margin slightly revolute to sinuate, ciliate, puberulous or estrigose in the upper surface, densely estrigose to setaceous in the lower surface, whitish, trichomes bulbous on both surfaces, hiphodromous, showing impressed veins in the upper surface and prominent veins in the lower surface. **Inflorescence** 1.9-8.4 cm, terminal and axial, dichotomous, lax, slightly escorpioideous, peduncle 0.7-1.4 cm, bracts 4.5 × 1.5 mm, lanceolate, estrigose. **Flowers** 5. 5-7 mm, short pedicellate. **Calyx** 4.5-5.5 mm, obpyramidal, laciniae 4-6 × 0.4-1 mm, subcarnose, linear to lanceolate, setaceous outside, glabrous inside, margin ciliate, whitish, presenting sparse blackish punctations, central region greenish. **Corolla** 5-6.6 mm, white, funnel-shaped, sericeous outside, villose inside just in the stamen attachment, tube 3.5-4 mm, lobes ca. 2 × 2 mm, patent, largely deltoideous, waved-plicate. **Stamens** subsessile, attached to ca. 1.5 mm of the base of the corolla's tube, anthers 1 mm, connate, ovate to ovate-lanceolate, apices with glands and trichomes. **Ovary** 0.5-0.7 mm, depressed-globose, sulcate, stipitate. **Style** ca. 1 mm, cylindrical. **Stigma** 0.5-0.7 mm, fertile region conical, slightly 2-furcate in the apice, puberulous. **Fruit squizocarp** 2-2.5 mm diameter, depressed-globose, puberulous, calyx and stigma attached, nutlets 4 with 1 mm, presenting three faces. **Seed** 1 for nutlet, 1 to 1.2 mm, elliptical, curved embryo.

This species belongs to the genus *Euploca* for its bracteate inflorescences, connate anthers with appendages, glands and trichomes, in their apices, as well as for its fruits composed by four nutlets and seeds with a curved embryo. It is separated from the other Brazilian *Euploca* species by its discolour leaves, a decurrent petiole, exclusively dichotomous inflorescences and mainly for its short pedicellate flowers, a calyx presenting subcarnose laciniae and an estipitate ovary.

**Etymology.** – The specific epithet was given in honor to Dr. Maria Jesus Nogueira Rodal of the Universidade Federal Rural de Pernambuco at Recife, PE, Brazil, researcher of the Brazilian Northeast's flora.

**Distribution.** – Known from Brazil: states of Pernambuco and Bahia. In Pernambuco, it is found in caatinga vegetation. In Bahia, it is distributed in Northwest and Southeast in caatinga and cerrado vegetation.

**Ecology.** – The species occurs isolated or in small assemblages, generally on the road-sides, at elevations between 560-940 m. However, in the cerrado vegetation, *E. rodaliae* grows on sandy soils associated to rocks.

*Phenology.* – Flowering from February to November and fruiting from February to September.

*Paratypes.* – **BRAZIL. Bahia:** Barreiras, IX.1978, fl., fr., *L. Coradin & al.* 1177 (CEN); Caetité, 13°59'35"S, 42°12'27"W, 8.III.1997, fl., fr., *E. Saar & al.* 5242 (ALCB 35414, HUEFS 29030); Caetité, 13°51'17"S, 42°16'12"W, II.1997, fl., fr., *B. Stannard & al.* 5199 (ALCB, CEPEC, HUEFS); Santaluz, 11°11'16"S, 39°25'48"W, XI.1986, fl., *L. P. Queiroz & al.* 1127 (HUEFS). **Pernambuco:** Caruaru, 1998, fl., fr., *V. C. Lima s.n.* (IPA 65497).

#### ACKNOWLEDGEMENTS

We are grateful to CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for a PhD fellowship to the first author, to Mrs. Maria de Fátima de Araújo Lucena for comments and suggestions on the first manuscript's version, to Mr. Frank Valdomiro Silva for the drawings, to Mr. Enio Wocilly Dantas and Mr. Raphael Arias for the French Abstract and to Mr. Antonio Travassos for the revision of this text.

#### REFERENCES

- GRAY, A. (1874). Contributions to the botany of North America II: Notes on Boraginaceae. *Proc. Amer. Acad. Arts* 10: 48-61.
- HARRIS, J. G. & M. W. HARRIS (1994). *Plant identification terminology: an illustrated glossary*. Spring Lake Publishing, Utah.
- HICKEY, L. J. (1973). Classification of the architecture of dicotyledonous leaves. *Amer. J. Bot.* 60: 17-33.
- HILGER, H. H. & N. DIANE (2003). A systematic analysis of Heliotropiaceae (Boraginales) based on trnL and ITS1 sequence data. *Bot. Jahrb. Syst.* 125: 19-51.
- MELO, J. I. M. & M. F. SALES (2004). *Heliotropium* L. (Boraginaceae – Heliotropioideae) de Pernambuco, Nordeste do Brasil. *Rodriguésia* 84: 65-87.
- NUTTALL, T. (1837). Collection towards a flora of the territory of Arkansas. *Trans. Amer. Philos. Soc.* 5: 137-203.
- PAYNE, W. W. (1978). A glossary of plant hair terminology. *Brittonia* 30: 239-255.
- RIZZINI, C. T. (1977). Sistematização terminológica da folha. *Rodriguésia* 42: 103-125.

Submitted on June 7, 2005

Accepted on July 4, 2006

---

Addresses of the authors: JIMM: Programa de Pós-Graduação em Botânica, Universidade Federal Rural de Pernambuco, Rua Dom Manoel de Medeiros, s/n, Dois Irmãos, 52171-900, Recife, PE, Brasil. Email: [jimmelo@zipmail.com.br](mailto:jimmelo@zipmail.com.br)

JS: Departamento de Botânica, Instituto de Biologia, Universidade Estadual de Campinas, C.P. 6109, 13083-970, Campinas, SP, Brasil.




Fig. 1. – *Euploca rodaliae* J. I. M. Melo & Semir. **A.** Habit; **B.** Leaf; **C.** Flower; **D.** Sepal; **E.** Bract; **F.** Corolla dissected; **G.** Detail of the androecium, showing anthers connate; **H.** Gynoecium; **I.** Fruit in superior view.

[**A-D** and **F-I**: B. Stannard & al. 5199; **E**: E. Saar & al. 5242] (Drawing Frank Valdomiro Silva)