

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	61 (2006)
Heft:	2
Artikel:	Holothrinx klimkoana Szlach. & Marg. (Orchidaceae, Orchidoideae) : a new species from Angola
Autor:	Szlachetko, Dariusz L. / Margoska, Hanna B.
DOI:	https://doi.org/10.5169/seals-879261

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Holothrix klimkoana Szlach. & Marg. (Orchidaceae, Orchidoideae), a new species from Angola

DARIUSZ L. SZLACHETKO
&
HANNA B. MARGOŃSKA

ABSTRACT

SZLACHETKO, D. L. & H. B. MARGOŃSKA (2006). *Holothrix klimkoana* Szlach. & Marg. (Orchidaceae, Orchidoideae), a new species from Angola. *Candollea* 61: 467-470. In English, English and French abstracts.

Holothrix klimkoana Szlach. & Marg. (Orchidaceae, Orchidoideae), new species from Angola, is described, illustrated and compared to its closest relative, *H. longiflora* Rolfe.

RÉSUMÉ

SZLACHETKO, D. L. & H. B. MARGOŃSKA (2006). *Holothrix klimkoana* Szlach. & Marg. (Orchidaceae, Orchidoideae), une nouvelle espèce décrite d'Angola. *Candollea* 61: 467-470. En anglais, résumés anglais et français.

Holothrix klimkoana Szlach. & Marg. (Orchidaceae, Orchidoideae), nouvelle espèce d'Angola, est décrite, illustrée et comparée au taxon le plus proche, *H. longiflora* Rolfe.

KEY WORDS: ORCHIDACEAE – ORCHIDOIDEAE – *Holothrix* – Africa

The genus *Holothrix* was described by LINDLEY (1835). It embraces about 50-60 species distributed widely in Africa with a few in tropical Arabia. They are characterized by 1 or 2 small, ellipsoid or ovoid tubers. The one or two leaves are sessile, reniform, ovate to orbicular, radical, often papillose or hairy. The stem is erect with or without caudine bracts, glabrous, papillose or hairy. The inflorescence is terminal, many-flowered. The flowers are resupinate, often secund, sessile to pedicellate, tubular or widened apically. The sepals are usually smaller than the petals, often hairy. The petals are usually divided into 3 or more fleshy, finger-like or filiform segments. The lip is similar to the petals with a cylindrical spur (SZLACHETKO & OLSZEWSKI, 1998; PRIDGEON & al., 2001). The gynostemium is adnate to the lip basally. The stigma is 2-lobed, with lobes divergent, small. The rostellum is shelf-like, truncate, indistinctly 3-lobed, middle lobe small, folded, side lobes flat, separated from one another. The viscidia are minute, two, with very short caudicles. The anther is more or less ovoid to ellipsoid, erect, connective relatively wide, with locules parallel. Anther basal projections obscure the rostellum side lobes and viscidia. The pollinia are sectile. Auricles absent (SZLACHETKO & RUTKOWSKI, 2000).

While working on “Contribution to the Orchid Flora of Central West Africa (Tchad, Cameroon, Republic of Central Africa, Equatorial Guinea, Principe, Sao Tome, Congo, Gabon, Angola)”, we had access to herbarium and liquid preserved specimens kept in various collections. Among materials from Angola, deposited at the Muséum National d’Histoire Naturelle in Paris, we found specimens belonging to the genus *Holothrix*, with lip and petals deeply dissected into filiform segments. At first it appeared to be similar to *H. longiflora* Rolfe, from which it is easily distinguished by its secondarily divided petals and sepals. In our opinion the specimens from Angola represent an unknown taxon, which deserves to be recognized as a new species.

***Holothrix klimkoana* Szlach. & Marg., spec. nova** (Fig. 1).

Type: ANGOLA: Humbe et bords du Cunene, Tyipelongo-Mucope-Donguena, 1910-1920, Bonneforex & Villain 62 (holo-: P!; iso-: P!).

Species haec H. longiflorae affinis sed petalis secundariter ramosis recedit.

Stem 30-38 cm tall, erect, stout, densely hairy, with 2 caudate bracts above. Leaves 2, one larger than the other, 2.5-3.5 cm long, 2.5-4 cm wide, adpressed to ground, opposite, cordate to reniform, obtuse, sparsely hairy. Inflorescence 6.5-12 cm long, densely 10-18-flowered, secund. Floral bracts up to 12 mm long, densely hairy. Dorsal sepal 8.5 mm long, 3 mm wide, oblong-ovate, apiculate, concave, ciliate. Petals up to 35 mm long, 1-2 mm wide in the basal part, ribbon-like, above slightly expanded and divided into 3 slender lobes, which are secondarily divided into 3 filiform segments each. Lateral sepals 8 mm long, 3 mm wide, obliquely lanceolate-ovate, apiculate, ciliate. Lip up to 32 mm long, ribbon-like in the lower half, then much expanded and divided into 3 main lobes, which are secondarily divided into numerous filiform elements. Spur 4 mm long, conical, forming a complete loop, acute. Gynostemium 2 mm high. Ovary 8-10 mm long, hairy.

Etymology. – Dedicated to Prof. dr hab. Małgorzata Klimko, Polish taxonomist and botanist.

Ecology. – No data available.

General distribution. – Endemic to Angola?

ACKNOWLEDGMENTS

We are grateful to the Curators of P for the loan of herbarium specimens and their hospitality during our personal visits. We are indebted to Prof. Dr. hab. Ryszard Ochyra for the Latinization of the diagnosis and also to M. Sc. Agnieszka Kowalkowska for the drawings.

REFERENCES

- LINDLEY, J. (1835). *Holothrix. Gen. Sp. Orchid. Pl.*: 257.
- PRIDGEON, A. M., P. J. CRIBB, M. W. CHASE & F. N. RASMUSSEN (2001). *Genera Orchidacearum* 2: 1-416. Oxford University Press, New York.
- SZLACHETKO, D. L. & T. S. OLSZEWSKI (1998). Orchidacées. Vol. 1. In: MORAT, P. (ed.), *Flore du Cameroun* 34: 1-327.
- SZLACHETKO, D. L. & P. RUTKOWSKI (2000). Gynostemia Orchidálium 1. Apostasiaceae, Cypripediaceae, Orchidaceae (Thelymitroideae to Vanilloideae). *Acta Bot. Fenn.* 169: 1-380.

Submitted on May 30, 2005

Accepted on December 4, 2006




Fig. 1. – *Holothrix klimkoana* Szlach. & Marg. **A**: flower; **B**: lip; **C**: petal; **D**: dorsal sepal; **E**: lateral sepal.

[Bonneforex & Villain 62, P] (Drawings by Agnieszka Kowalkowska)