

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	62 (2007)
Heft:	2
Artikel:	Begonia callosa L. Kollmann (Begoniaceae) : a new species from the Atlantic forest in the State of Espírito Santo, Brazil
Autor:	Kollmann, Ludovic Jean Charles
DOI:	https://doi.org/10.5169/seals-879172

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Begonia callosa L. Kollmann (Begoniaceae), a new species from the Atlantic forest in the State of Espírito Santo, Brazil

Ludovic Jean Charles Kollmann

Abstract

KOLLMANN, L. J. C. (2007). *Begonia callosa* L. Kollmann (Begoniaceae), a new species from the Atlantic forest in the State of Espírito Santo, Brazil. *Candollea* 62: 141-144. In English, English and French abstracts.

Begonia callosa L. Kollmann is a new species described and illustrated. It is related to *Begonia reniformis* Dryand., from which it differs by the apical calluses on the petioles and hypopodia, the carinate stipules, and the strongly canaliculate and zigzag stems.

Key-words

BEGONIACEAE – *Begonia* – Brazil – State of Espírito Santo – Atlantic Forest

Résumé

KOLLMANN, L. J. C. (2007). *Begonia callosa* L. Kollmann (Begoniaceae), une nouvelle espèce de la forêt atlantique de l'Etat d'Espírito Santo, Brésil. *Candollea* 62: 141-144. En anglais, résumés anglais et français.

Begonia callosa L. Kollmann est une nouvelle espèce décrite et illustrée. Cette espèce est proche de *Begonia reniformis* Dryand., mais s'en différencie par la présence de callosités à l'extrémité des pétioles et des hypopodes, ses stipules fortement carinés et des entremiseaux caulinaires fortement caniculés et en zigzag.

Address of the author: Museu de Biologia Prof. Mello Leitão, 4 Av. José Ruschi CEP 29650-000 Santa Teresa, ES, Brazil.
Email: ludovic@limainfo.com.br

Submitted on January 9, 2007. Accepted on August 10, 2007.

Introduction

The family *Begoniaceae* comprises two genera: *Hillebrandia* Oliver, with one species from the Hawaiian Islands, and *Begonia* L. (CLEMENT & al., 2004), pantropical and with about 1400 species (DOORENBOS & al., 1998), of which about 240 are native to Brazil (SMITH & al., 1986), mostly along the Atlantic Forest (DUARTE, 1961). The Atlantic Forest was recently classified as one of the 25 most important world hotspots for conservation (SIMÕES & al., 2002; MITTERMEIER & al., 1999) because of the threats to its conservation, of its large biological wealth, and of its high level of endemism. In the Atlantic Forest, *Begonia* species occur in different habitats such as marshes, sunlight place and wet soil, rain and dry forest, altitude vegetation, and xerophytic vegetation (BRADE, 1961).

During fieldwork in the mountainous region of central Espírito Santo State, we collected a species of *Begonia* that we are proposing as new and describing herein.

Result and discussion

Begonia callosa L. Kollmann, spec. nova (Fig. 1)

Typus: BRAZIL. Espírito Santo: Fundão, Goiapaba-açu, 21.X.2005, L. Kollmann 8388 & A. P. Fontana (holo-: MBML; iso-: RB).

Species haec B. reniformi Dryand. affinis, sed petiolis et hypopodiis apicis callosis, stipulis carinatis, caulinibus angulosis et caniculatis differt.

Suffrutescent herbs 2-3 m tall, pilose, trichomes simple. Stems 1-2.2(-4.5) cm in diameter, green, brown with age, striate, lenticels present, internodes 3-10(-35) cm long. Petioles 13-21 cm, green, base and apex reddish, adaxial face sulcate, apically sulcate and callose, lenticels present. Stipules deciduous, 2-4.5 cm long, green, ovate to triangular, carinate. Leaf blades (20-)26-45(-55) × (14-)18-27(-42) cm, green, abaxial face reddish green, pilose, base cordate, apex acuminate; margins serrate, lightly ciliate; venation actinodromous, veins 6-7, pilose, stomates single. Inflorescence 26-69 cm long, with 6-9 nodes, green, dichasial, the first hypopodium sulcate with an apical callus, the others lightly sulcate, all with apical calluses, lenticels present. Bracts persistent, opposite, 2-10 mm long, triangular, glabrous, apex acute and apiculate. Staminate flowers: peduncles 5.5-6 mm long, reddish pink; tepals 4, white, the outer 0.5-1 × 0.4-0.6 cm, concave, obelliptic, margins reddish, apex rounded to obtuse, the inner 0.5-0.7 × 0.15-0.3 cm, oblong to obovate, apex acute; stamens ca. 27, yellow, rimose, obovate, filaments unequal, the connective projecting beyond the anthers. Pistillate flowers: pedicels 5.5-14 mm long, white; tepals 5, 0.55-1 × 0.15-0.6 cm, unequal, ovate to obovate, apex acute; styles 3, yellow, bifurcate, spirally twisted, united at base, stigma with rows of stigmatic papillae on the

branches, placenta one per locule, ovules present on both surfaces. Capsules 0.8-1.2 × 0.4-0.5 cm, glabrous, basally dehiscent, wings 3, the larger 0.8-1.2 × 1.1-1.6 cm, the smaller two 0.8-1.3 × 1-2 mm, glabrous. Seeds cylindrical, oblong.

Paratypi. – **BRAZIL. Espírito Santo:** Fundão municipality, Alto Piaba, 27.VII.1988, W. Boone 1219 & al. (MBML, RB); Goiapaba-açu, 18.III.2005, L. Kollmann 7470 & al. (MBML); Santa Teresa municipality, Valsugana Velha, 24.V.2000, L. Kollmann 2959 & al. (MBML); Rio Saltinho, 7.VI.2001, L. Kollmann 3852 & al. (MBML); 21.X.2005, L. Kollmann 8373 & al. (MBML); Valsugana Velha, Estação Biológica de Santa Lúcia, 19.VI.2001, L. Kollmann 4095 & al. (MBML); 11.IX.2005, L. Kollmann 8430 & al. (MBML); Nova Lombardia, Reserva Biológica Augusto Ruschi, 16.X.2001, L. Kollmann 4871 & al. (MBML); Santa Leopoldina municipality, Suíça, 29.I.2006, A. P. Fontana 1933 & A. M. Assis (MBML); Atílio Vivácqua municipality, Moitão, 27.IV.2007, L. Kollmann & al. 9763 (MBML).

Etymology. – The epithet refers to the apical calluses of petioles and hypopodia.

Begonia callosa grows in sunlight in the leaf litter and humus of rocky openings of the Atlantic Forest, at 200 to 800 m. It is also found growing in low canopy forests from open environments on sandy soil.

Begonia callosa resembles *B. reniformis* Dryand., another Brazilian species of sect. *Pritzelia* (Klotzsch) A. DC., due to its lobate leaves, inflorescences, flowers and fruits. Nevertheless, it can be distinguished from *B. reniformis* by the presence of a callus at the apex of the petioles and hypopodia, petioles adaxially sulcate, the strongly carinate stipules and the strongly canaliculate and zigzag stems.

Acknowledgments

We acknowledge the staff of Mello Leitão Biological Museum, especially Helio de Queiroz Boudet Fernandes, Director of the Museum and Curator of MBML, for their collaboration. Marcos Sobral for suggestions and help with the English version and Latin diagnosis. The Petrobras to financing the project.

References

- BRADE, A. C. (1961). O porte das begônias brasileiras e os ambientes onde ocorrem. *Arch. Jard. Bot. Rio de Janeiro* 17: 51-55.
- CLEMENT, W. L., M. C. TEBBITT, L. L. FORREST, J. E. BLAIR, L. BROUILLET, T. ERIKSSON & S. M. SWENSEN (2004). Phylogenetic position and biogeography of *Hillebrandia sandwicensis* (Begoniaceae): a rare Hawaiian relict. *Amer. J. Bot.* 91: 905-917.

DOORENBOS, J., S. M. SOSEF & J. J. F. E. DE WILDE (1998). The sections of Begonia. Studies in Begoniaceae VI. *Wageningen Agric. Univ. Pap.* 98(2): 1-266.

DUARTE, A. P. (1961). Considerações acerca do comportamento e dispersão de algumas espécies de Begônias do Estado da Guanabara. *Arch. Jard. Bot. Rio de Janeiro* 17: 57-105.

MITTERMEIER, R. A., N. MYERS & C. G. MITTERMEIER (1999). *Hotspots Earth's biologically richest and most endangered terrestrial ecoregions*. Conservation International.

SIMÕES, L. L. & C. F. LINO (2002). *Sustentável Mata Atlântica: a exploração de seus recursos florestais*. Editora Senac.

SMITH, L. B., D. C. WASSHAUSEN, J. GOLING & C. E. KAREGEANNES (1986). Begoniaceae. Part I: Illustrated key. Part II: Annotated Species List. *Smithsonian Contr. Bot.* 60: 1-584.




Fig. 1. – *Begonia callosa* L. Kollmann. **A.** Habit; **B.** Petiole callus apice; **C.** Cross section of petiole callus; **D.** Cross section of petiole base; **E.** Stipule, lateral view; **F.** Stipule, abaxial face extended; **G.** Petal male flower; **H.** Sepal male flower; **I.** Stamen; **J.** Tepal femal flower; **K.** Stigma, adaxial face; **L.** Stigma, abaxial face; **M.** Fruit; **N.** Seed.

[Drawn by the author]

