

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	62 (2007)
Heft:	2
Artikel:	Two new epitypes in Danaea (Marattiaceae, Pteridophyta) selected from original historical collections in Paris
Autor:	Christenhusz, Maarten J. M.
DOI:	https://doi.org/10.5169/seals-879183

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Two new epitypes in *Danaea* (Marattiaceae, Pteridophyta) selected from original historical collections in Paris

Maarten J. M. Christenhusz

Abstract

CHRISTENHUSZ, M. J. M. (2007). Two new epitypes in *Danaea* (Marattiaceae, Pteridophyta), selected from original historical collections in Paris. *Candollea* 62: 221-230 In English, English and French abstracts.

Charles Plumier (1646-1704) completed nearly 200 drawings of ferns and clubmosses during his travels in the French Caribbean between 1689 and 1695. Many of these plates have been cited by Linnaeus in his *Species Plantarum* (1753), and were subsequently designated as lectotypes for the taxa they represent. Surveys of historical specimens from the herbarium in the National Museum of Natural History in Paris have revealed many original fern specimens annotated by Plumier that have most probably been collected by Surian. Several of these specimens should be designated as epitypes, in addition to the lectotypified Plumier illustrations, because not all distinguishing characters necessary to unambiguously identify the species are represented in the Plumier plates. The advantage of these epitypes is that they are specimens, rather than illustrations, and show characters that are not visible in Plumier's illustrations. We have chosen the genus *Danaea* Sm. (Marattiaceae) as an example and provide new synonymy and a complete taxonomic history of the genus. Two new epitypes of *Danaea* are proposed here, *Danaea alata* Sm. and *Danaea nodosa* (L.) Sm. The name *Danaea angustifolia* C. Presl is lectotypified.

Key-words

MARATTIACEAE – *Danaea* – Antilles – Plumier – Epitypes – Ferns – Historical specimens – Pre-Linnaean illustrations

Address of the author: MJMC: Department of Biology, University of Turku, 20014 Turku, Finland.
Email: maarten.christenhusz@utu.fi

Submitted on December 19, 2006. Accepted on October 30, 2007.

ISSN: 0373-2967

Candollea 62(2): 221-230 (2007)

© CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 2007

Résumé

CHRISTENHUSZ, M. J. M. (2007). Deux nouveaux épitypes de *Danaea* (Marattiaceae, Pteridophyta), sélectionnés à partir des collections originales historiques à Paris. *Candollea* 62: 221-230. En anglais, résumés anglais et français.

Charles Plumier (1646-1704) a réalisé près de 200 illustrations de fougères et de lycopodes durant son voyage dans les Caraïbes françaises entre 1689 et 1695. Ces planches ont été citées, entre autres, par Linné dans son *Species Plantarum* (1753), et ont été désignées comme les lectotypes des taxons qu'elles représentent. Des recherches récentes dans les collections historiques de l'herbier du Muséum national d'Histoire naturelle à Paris ont permis de découvrir de nombreux spécimens annotés par Plumier et probablement collectés par Surian. Plusieurs de ces spécimens devraient être désignés comme épitypes, en complément des illustrations de Plumier classées comme lectotypes, du fait de l'absence dans les planches de Plumier de certains caractères distinctifs nécessaires à une identification sans ambiguïté des espèces. Ces épitypes présentent l'avantage d'avoir conservé des caractères qui ne sont pas visibles sur les illustrations. Nous avons choisi le genre *Danaea* Sm. (Marattiaceae) comme exemple et en donnons une nouvelle synonymie ainsi qu'un historique taxinomique. Deux nouveaux épitypes de *Danaea* sont proposés ici, *Danaea alata* Sm. et *Danaea nodosa* (L.) Sm. Le nom *Danaea angustifolia* C. Presl est lectotypifié.

Introduction

The French missionary Charles Plumier (1646–1704), a member of the catholic order of the Minims, sketched numerous plants during his stay on Martinique and Hispaniola. He travelled in the Caribbean between 1689 and 1695, and had a close working relationship with Joseph Donat Surian, who later returned to France with a large collection of dried specimens (many of which are still preserved in P). Plumier did not collect specimens himself, but instead drew and described the plants he studied in the field, which were subsequently collected by Surian. These drawings and descriptions formed the basis of a series of works: PLUMIER (1693, 1703, 1705), describing the flora of the French Caribbean. In PLUMIER (1705), 50 plates of his 1693 publication were reproduced. In total nearly two hundred ferns and clubmosses were described and illustrated by Plumier (PENNELL, 1945).

LINNAEUS (1753) based several species descriptions on Plumier's illustrations, since he had been able to study copies of these in Groningen, the Netherlands. Several authors such as FÉE (1866), JENMAN (1888), URBAN (1925: 273–397), PROCTOR (1977, 1985, 1989), LELLINGER & PROCTOR (1983) and PROCTOR & LOURTEIG (1990) designated Plumier plates as lectotypes for many Linnaean fern species. Also in several West Indian fern floras, PROCTOR (1977, 1985, 1989) selected lectotypes from among Plumier's drawings.

Despite the good quality and detail of the Plumier plates, a plate as a lectotype can be problematic if the image does not depict all characters necessary to distinguish between species. Some characters such as hairs, scales, glands or venation may not be well illustrated. According to MCNEILL & al. (2006), “when a holotype, a lectotype, or a neotype is an illustration, the specimen or specimens upon which that illustration is based should be used to help determine the application of the name” (*recommendation 8A.1*). For several species the Plumier plates are demonstrably ambiguous, which would imply the designation of the original specimens in the Surian and Tournefort collections that were annotated by Plumier, as epitypes (MCNEILL & al., 2006, *article 9.7*). The lectotype and epitype should both be considered when deciding on the identity of a species. It is in this spirit that I have conducted the study of the specimens in the Tournefort collection and other historical collections at P.

Many of the original specimens on which Plumier based his illustrations are preserved in the herbaria of Tournefort (P-TRF), Vaillant (P), and Surian (P). In cases where the illustration is not sufficient for the unambiguous identification of the plant, many of these specimens may be designated as epitypes.

Because I am currently revising the genus *Danaea* Sm. (*Marattiaceae, Pteridophyta*), I chose this genus as an example for epitypifying specimens linked to Plumier plates. The

identity of the two species depicted by Plumier is ambiguous, because the venation, pinna margins and petiole nodes are not clearly illustrated, and these names have therefore often been misapplied (i.e. PRESL, 1845; JENMAN, 1898; ROLLERI, 2004). The illustrations are thus not sufficient to distinguish these species they depict from related taxa. The epitypes selected in this study will clarify this issue and may prevent confusion of *D. alata* Sm. and *D. nodosa* (L.) Sm. with related species in the future.

Taxonomic History of *Danaea*

The first mention of taxa now belonging to the genus *Danaea* was by PLUMIER (1705), who published two plates that he named: “*Lingua cervina nodosa minor*” (Fig. 1) and “*Lingua cervina nodosa maior*” (Fig. 2). LINNAEUS (1753) applied “*Lingua cervina nodosa major*” to *Acrostichum nodosum* L. Even though PLUMIER (1705) stated that the species was observed in either Martinique or Hispaniola, LINNAEUS (1753) wrote “*Habitat in Dominica*”, and further indicated “*stipitis articulis nodosis*”, an unfortunate error because this species does not bear nodes on the petioles. LINNAEUS (1763) reconsidered the placement of *Acrostichum nodosum* and transferred it to *Asplenium*.

The second plate of PLUMIER (1705: 91, tab. 109) “*Lingua cervina nodosa minor*”, was not mentioned by LINNAEUS (1753), but SMITH (1793) described it as *Danaea alata*, indicating “*Habitat in Martinica, Plumier in herb. Linn.*” as the original material (not present at LINN). In this article SMITH (1793) described *Danaea* in honor of “*Bot. Prof. J. Petri Mariae Dana*”, and he based his new genus on *Asplenium nodosum* (L.) L. (= *Acrostichum nodosum* L.), citing “*Plumier, Fil. 90, l. 108*” and referring to “*a specimen in the Linnaean Herbarium*”. The latter was probably an assumption made at the time, for there is no original *Danaea* material known in LINN.

RUDGE (1805: 24) added a third species, *D. simplicifolia* Rudge, which is remarkable for its undivided blades. SMITH (1808) revised his genus for Rees' *Cyclopaedia*, and added a fourth species *D. elliptica* Sm. The description of this species was ambiguous, and therefore this name became very widely but variously applied. However, the material (*Herb. Sloane 1: 85, BM-SL*) indicated in the original description by SMITH (1808) and lectotypified by PROCTOR (1977), is a juvenile of *D. nodosa*, and thus the name *D. elliptica* becomes a synonym of that species (CHRISTENHUSZ & TUOMISTO, 2006).

In subsequent years, six new species were added to the genus (DESVAUX, 1811; RADDI, 1825: 75–76; KUNZE, 1837, 1840: 55–57, 1843: 137–138), until PRESL (1845) revised the genus. He described four additional species, erroneously synonymised *D. geniculata* Raddi under *D. elliptica*, and placed the 11 species

known to him into three sections: *Eudanaea* C. Presl, *Arthrodanaea* C. Presl and *Holodanaea* C. Presl, based on inconsistent morphological characters. Additionally two species were placed into segregate genera: *Heterodanaea* C. Presl and *Danaeopsis* C. Presl. The lectotype of *Heterodanaea* [*Danaea stenophylla* Kunze; Guadeloupe, l'Herminier 213 ex herb. Bory, *Danaea* nr. 37 (K!)], designated by PROCTOR (1977)] is a specimen identical to *Danaea alata* (ROLLERI, 2004). It therefore becomes a synonym of that species, and *Heterodanaea* is thus a synonym of *Danaea*. The other genus *Danaeopsis* was based on *Danaea paleacea* Raddi of which the holotype is: *Raddi* s.n. (P!) from Serra d'Estrella, Est. Sao Paulo, Brazil. This specimen has reticulate venation, a character that does not occur in *Danaea*, and is *Bolbitis serratifolia* (Kaulf.) Schott (PICHI-SERMOLLI & BIZZARRI, 2005), which coincidentally is the type species of the genus *Bolbitis* Schott (Dryopteridaceae). To confuse matters, the fossil genus *Danaeopsis* Heer (HEER, 1865) is in use for certain marattialean fossils. The correct name for these fossils is *Marantoidea* Jaeger (WEBB, 2001). Presl's genera, sections and classification were not widely accepted and were rarely followed by subsequent authors on *Danaea*.

Over the following 57 years, little taxonomic attention was given to *Danaea* apart from listings in various indices (i.e. MOORE, 1861: 285-288), and the occasional addition of new species (LIEBMANN, 1849; FÉE, 1869; REICHENBACH, 1872; BAKER, 1881, 1891; JENMAN, 1898).

UNDERWOOD (1902) published the first review of the genus. He described, typified, and redefined several species and provided an analytical key for 12 species, of which five were new to science. This work was adapted for the North American Flora, where UNDERWOOD (1909) determined the type locality for *D. nodosa* as "Near Port de Paix, Haiti", and where he provided descriptions of a dozen species, which, remarkably, were a different set of species than he presented in his 1902 review. Meanwhile several new species were described (CHRIST, 1905, 1907, 1909, 1910; ROSENSTOCK, 1907, 1909), and CHRISTENSEN (1906, 1913) provided an overview of the genus listing a total of 32 species names.

SMITH (1793) did not explicitly establish that *D. nodosa* was the type of *Danaea*. So PICHI-SERMOLLI (1957) designated it as such. Additionally he proposed to conserve the genus *Danaea* Sm. over *Danaa* All. (*Apiaceae*) and *Danaa* Colla (*Asteraceae*), which was approved by MCNEILL & al. (2006).

It took nearly a century, during which only a few new *Danaea* species were described (MAXON, 1924; ROSENSTOCK, 1925; MORTON, 1951; STOLZE, 1987), before TUOMISTO & MORAN (2001) revised the, until then, poorly known and challenging genus *Danaea*. They recognised a total of 18 species for Ecuador alone, of which eight were newly described. Their

work was the first comprehensive regional monograph of the genus, based on field work and herbarium studies. Subsequently, ROLLERI (2004) accepted a total of 17 species of *Danaea* in her revision of the genus. Recent studies (CHRISTENHUSZ & TUOMISTO, 2005; CHRISTENHUSZ, 2006; CHRISTENHUSZ & TUOMISTO, 2006; CHRISTENHUSZ & al., in press) have shown however that several species, not recognised by ROLLERI (2004), are morphologically and genetically distinct and warrant recognition. I estimate this exclusively Neotropical genus to consist of approximately 50 species, but several species complexes need to be unravelled before a more exact species number can be established.

New epitypes in *Danaea*

Several specimens in P-TRF have been annotated by Plumier. Many of these specimens are available as lecto- or epitypes of species that are based on Plumier's descriptions and illustrations. As an example of how typifications may need to be revised in the light of this collection, I epitypify two *Danaea* species below, and we provide their full synonymy.

1. *Danaea alata* Sm. in Mém. Acad. Roy. Sci. Turin 5: 420. 1793.

Lectotypus (designated by PROCTOR, 1977): Martinique, Morne de la Calebasse (PLUMIER, 1705: tab. 109) (Fig. 1).

Epitypus (designated here): *Ibitoboua Surian* 648, Herbarium Vaillant 56 (P!) (Fig. 2).

- = *Danaea fendleri* Underw. in Bull. Torrey Bot. Club 29: 673. 1902. **Typus: TRINIDAD & TOBAGO. Trinidad:** *A. Fendler* 147 (holo-: NY!; iso-: BM!, GH!, K!, US!, YU!).
- = *Danaea stenophylla* Kunze, Farnkraüter 1: 55. 1840. = *Heterodanaea stenophylla* (Kunze) C. Presl, Suppl. Tent. Pterid.: 38. 1845. **Typus: GUADELOUPE: l'Herminier 213 ex herb. Bory, Danaea nr. 37** (holo-: LZ [destroyed]; lecto-: K!; isolecto-: P!, NY! (fragment of K), P! (fragment), designated by PROCTOR, 1977).

The material cited by SMITH (1793) was "Plum. Fil. 91., t. 109" and "Habitat in Martinica, Plumier in herb. Linn." The venation of the epitype is clearly that of *D. alata*, and matches modern specimens collected in Martinique (i.e. Christenhusz 2711, 2713, TUR!). The venation depicted in the plate is variable and unclear. Since this character is very important in distinguishing this species from the otherwise similar *D. mazaeana* Underw. and *D. moritziana* C. Presl, an epitype showing the venation is necessary.

Since its publication the name *D. alata* has been very variously applied, and some authors have erroneously synonymised species (such as *D. moritziana* and *D. mazaeana*)

with this name (i.e. PRESL, 1845; JENMAN, 1898; ROLLERI, 2004). Revision of this complex requires the application of the name *D. alata* to an original specimen, hence this epitype is selected.

Additional historical specimen. – Herbarium Tournefort 5383 (P-TRF!).

2. *Danaea nodosa* (L.) Sm. in Mém. Acad. Roy. Sci. Turin 5: 420. 1793.

= *Acrostichum nodosum* L., Sp. Pl.: 1070. 1753.

= *Asplenium nodosum* (L.) L., Sp. Pl. ed. 2: 1539. 1763.

Lectotypus (designated by UNDERWOOD, 1909): Haiti, Port de Paix (PLUMIER, 1705: tab. 108) (Fig. 3).

Epitypus (designated here): *Herbarium Vaillant* 43 (P!) (Fig. 4).

= *Danaea angustifolia* C. Presl, Suppl. Tent. Pterid.: 35. 1845. = *Danaea nodosa* var. *angustifolia* (C. Presl) T. Moore, Index Fil. 2: 287. 1861. **Lectotypus** (designated here): Antilles, s.l., Bertero s.n. (P!; isolecto-: PRC!, fragment).

= *Danaea longifolia* Desv. in Mag. Ges. Naturf. Freunde Berlin 5: 307. 1811. = *Danaea nodosa* var. *longifolia* (Desv.) Hassl. in Trab. Inst. Bot. Farmac. 45: 89. 1928. **Lectotypus** (designated by PROCTOR, 1985): “*Habitat in Antillis*”, collector unknown [perhaps Lavallée s.n. anno 1896], ex herb. Desvaux (P!).

= *Danaea elliptica* Sm. in Rees, Cycl. 11: Danaea n° 2. 1808. **Lectotypus** (designated by PROCTOR, 1977): Jamaica, Mount Diablo, *Herb. Sloane* 1: 85 (BM-SL!). [It is not the lectotype suggested by LELLINGER (2000): West Indies, *Herb. Smith* No. 1645.7 (LINN-SM!). The lectotype of PROCTOR (1977) has priority (CHRISTENHUSZ & TUOMISTO, 2006)].

Danaea nodosa is part of a complex of closely related species, which has often been misunderstood. This has incorrectly resulted in a very broad application of the name *D. nodosa*. Revision of this group requires epitypification of *D. nodosa* to match this name to an actual specimen, rather than an illustration, permitting an unambiguous definition. The epitype will be an additional aid in defining the morphological differences between species of the *D. nodosa* group (CHRISTENHUSZ & TUOMISTO, 2005; CHRISTENHUSZ & al., in press) and it will facilitate the distinction of *D. nodosa* from closely related species (i.e. *D. cartilaginea* Christenb. & Tuomisto, *D. grandifolia* Underw., *D. kalevala* Christenb., *D. media* Liebm., *D. nigrescens* Jenm., *D. sellowiana* Underw., and *D. ushana* Christenb.).

Additional historical specimen. – Herbarium Tournefort 5377 (P-TRF!).

Acknowledgements

Many thanks to Gerard Aymonin, from the National Museum of Natural History in Paris, for sharing his knowledge of the historical collections. I am also grateful to the herbarium staff at Paris (P), and especially France Rakotondrainibe who permitted me to study the historical specimens, and who provided the photographs. In addition, I thank George Proctor (IJ) and Hanna Tuomisto (TUR) for their support and valuable comments, as well as Molly McMullen (DUKE), Michel Boudrie (CAY) and Frédéric Tronchet (P) for commenting on the manuscript.

References

- BAKER, J. G. (1881). On a collection of ferns made by Mr. W. Kalbreyer in New Grenada. *J. Bot.* 10: 208.
- BAKER, J. G. (1891). A summary of the new ferns which have been discovered or described since 1874. *Ann. Bot.* 5: 498-499.
- CHRIST, H. (1905). Filices Uleanae Amazoniae. *Hedwigia* 44: 368-370.
- CHRIST, H. (1907). Fougères nouvelles ou peu connues. *Bull. Herb. Boissier* 7: 927-928.
- CHRIST, H. (1909). Primitiae Florae Costaricensis. *Bull. Soc. Bot. Genève* 1: 234-235.
- CHRIST, H. (1910). Filices Costaricensis. *Repert. Spec. Nov. Regni Veg.* 8: 19-20.
- CHRISTENHUSZ, M. J. M. (2006). Three new species of *Danaea* (Marattiaceae) from French Guiana and the Lesser Antilles. *Ann. Bot. Fenn.* 43: 212-219.
- CHRISTENHUSZ, M. J. M. & H. TUOMISTO (2005). Some notes on the taxonomy, biogeography and ecology of *Danaea* (Marattiaceae). *Fern Gaz.* 17: 217-222.
- CHRISTENHUSZ, M. J. M. & H. TUOMISTO (2006). Five new species of *Danaea* (Marattiaceae) from Peru and a new status for *D. elliptica*. *Kew Bull.* 61: 17-30.
- CHRISTENHUSZ, M. J. M., H. TUOMISTO, J. S. METZGAR, & K. M. PRYER (in press). Evolutionary relationships within the Neotropical, euphorangioid fern genus *Danaea* (Marattiaceae). *Molec. Phylogen. Evol.*
- CHRISTENSEN, C. F. A. (1906). *Index filicum*. Copenhagen.
- CHRISTENSEN, C. F. A. (1913). *Index filicum, supplementum 1906-1912*. Copenhagen.
- DESVAUX, A. N. (1811). Observations sur quelques nouveaux genres de fougères et sur plusieurs espèces nouvelles de la même famille. *Mag. Ges. Naturf. Freunde Berlin* 5: 307.
- FÉE, A. L. A. (1866). Histoire des Fougères et des Lycopodiacées des Antilles. In: FÉE, A. L. A. (ed.), *Mémoires sur les familles des fougères* 11. Strasbourg.
- FÉE, A. L. A. (1869). Danaeacées. In: FÉE, A. L. A. (ed.), *Cryptogames Vasculaires du Brésil*: 214-217. Strasbourg.

- HEER, O. (1865). *Die Urwelt der Schweiz*. Zürich.
- JENMAN, G. S. (1888). Plumier's American ferns. *Gard. Field Forest* 4: 78.
- JENMAN, G. S. (1898). Ferns: Synoptical List - LIV. *Bull. Bot. Dept., Jamaica* 5: 208.
- KUNZE, G. (1837). *Analecta Pteridographica seu descriptio et illustratio filicum aut novarum, aut minus cognitarum*. Leipzig.
- KUNZE, G. (1840). *Schkahr's Farrnkräuter, Supplement* 28. Leipzig.
- KUNZE, G. (1843). *Schkahr's Farrnkräuter, Supplement* 60. Leipzig.
- LELLINGER, D. B. (2000). On the lectotypification of *Danaea elliptica*. *Amer. Fern J.* 90: 100-103.
- LELLINGER, D. B. & G. R. PROCTOR (1983). The ascriptions of Plumier's fern plates. *Taxon* 32: 565-571.
- LIEBMAN, F. (1849). Mexicos Bregner, en systematisk, critisk, plantegeographisk Undersögelse. *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.* 1: 151-333.
- LINNAEUS, C. (1753). *Species Plantarum*. Stockholm.
- LINNAEUS, C. (1763). *Species Plantarum*. Second edition. Stockholm.
- MAXON, W. R. (1924). Further notes on Hispaniola ferns. *J. Washington Acad. Sci.* 14: 195-196.
- MCMILLAN, J., F. R. BARRIE, H. M. BURDET, V. DEMOULIN, D. L. HAWKSWORTH, K. MARHOLD, D. H. NICOLSON, J. PRADO, P. C. SILVA, J. E. SKOG, J. H. WIERSEMA & N. J. TURLAND (2006). International code of botanical nomenclature (Vienna Code) adopted by the seventeenth International Botanical Congress, Vienna, Austria, July-August 2005. *Regnum Veg.* 146.
- MOORE, T. (1861). *Index Filicum*. Ed. 2. London.
- MORTON, C. V. (1951). A new fern of the genus *Danaea* from Colombia. *J. Washington Acad. Sci.* 41: 276.
- PENNELL, F. W. (1945). Historical Sketch. In: VERDOORN, F. (ed.), *Plants and Plant Science in Latin America*: 35-48. Waltham.
- PICHI-SERMOLLI, R. E. G. (1957). Names and types of fern genera 2, Danaeaceae. *Webbia* 12: 352-353.
- PICHI-SERMOLLI, R. E. G. & M. P. BIZZARRI (2005). A revision of Raddi's pteridological collection from Brazil (1817-1818). *Webbia* 60: i-vii, 1-393.
- PLUMIER, C. (1693). *Description des plantes de l'Amérique*. Paris.
- PLUMIER, C. (1703). *Nova plantarum americanarum genera*. Paris.
- PLUMIER, C. (1705). *Traité des Fougères de l'Amérique*. Paris.
- PRESL, C. B. (1845). Marattiaceae. In: Presl, C. B. (ed.), *Genera Filicacearum, Supplementum Tentaminis Pteridographiae*: 7-40. Prague.
- PROCTOR, G. R. (1977). Marattiaceae. In: Howard, R. A. (ed.), *Fl. Lesser Antilles* 2: 45-49. Arnold Arboretum, Jamaica Plain.
- PROCTOR, G. R. (1985). *Ferns of Jamaica*. London.
- PROCTOR, G. R. (1989). Ferns of Puerto Rico and the Virgin Islands. *Mem. New York Bot. Gard.* 53.
- PROCTOR, G. R. & A. LOURTEIG (1990). Nomenclatura Plantarum Americanum XIII, Pteridophyta. *Bradea* 5: 1-4.
- RADDI, I. (1825). *Plantarum Brasiliensium Nova Genera et species novae, vel minus cognitae collegit et descripsit I. Raddius* 1. Florence.
- REICHENBACH, H. G. (1872). Botanische Notizen. *Bot. Zeitung (Berlin)* 30: 487-491.
- ROLLERI, C. H. (2004). Revisión del género *Danaea* (Marattiaceae - Pteridophyta). *Darwiniana* 42: 271-301.
- ROSENSTOCK, E. (1907). Beiträge zur Pteridophytenflora Sübrasiliens. *Hedwigia* 46: 161-164.
- ROSENSTOCK, E. (1909). Filices Spruceanae adhuc nondum descriptae, in Herbario Rolandi Bonapartii Principis asservatae. *Repert. Spec. Nov. Regni Veg.* 7: 289-310.
- ROSENSTOCK, E. (1925). Filices novae a coll. Alfred et Curt Brade in Costa Rica collectae. *Repert. Spec. Nov. Regni Veg.* 22: 23.
- RUDGE, E. (1805). *Plantarum Guianae rariorum icones et descriptiones hactenus ineditae* 1. London.
- SMITH, J. E. (1793). *Danaea*. *Mém. Acad. Roy. Sci. Turin* 5: 420.
- SMITH, J. E. (1808). *Danaea*. In: REES, A. (ed.), *Cyclopaedia* 2 (D-Deluge). Philadelphia.
- SPRENGEL, C. (1800). Genera et species filicum. *J. Bot. (Schrader)* 1799(2): 272.
- STOLZE, R. G. (1987). A new species of *Danaea* from Peru. *Amer. Fern J.* 77: 33-35.
- TUOMISTO, H. & R. C. MORAN (2001). 7. Marattiaceae. *Fl. Ecuador* 66: 23-68.
- UNDERWOOD, L. M. (1902). American Ferns V. A review of the genus *Danaea*. *Bull. Torrey Bot. Club* 29: 669-679.
- UNDERWOOD, L. M. (1909). Marattiaceae. *N. Amer. Fl.* 16: 17-21.
- URBAN, I. (1925). *Symbolae Antillanae* 9. Berlin.
- WEBB, J. A. (2001). A New Marattialean Fern from the Middle Triassic of Eastern Australia. *Proc. Linn. Soc. New South Wales* 123: 215-224.




Fig. 1. – Lectotype of *Danaea alata* Sm.

[© Bibliothèque centrale du Muséum National d'Histoire Naturelle, Paris. Reproduced with permission]



Fig. 2. – Epitype of *Danaea alata* Sm.

[Surian 648, Herbarium Vaillant 56, P] [© Herbier du Muséum National d'Histoire Naturelle, Paris. Reproduced with permission]




Fig. 3. – Lectotype of *Danaea nodosa* (L.) Sm.

[© Bibliothèque centrale du Muséum National d'Histoire Naturelle, Paris. Reproduced with permission]




Fig. 4. – Epitype of *Danaea nodosa* (L.) Sm.

[Herbarium Vaillant 43, P] [© Herbier du Muséum National d'Histoire Naturelle, Paris. Reproduced with permission]

