

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	62 (2007)
Heft:	2
Artikel:	Typification of some critical taxa of Gagea Salisb. (Liliaceae) from W Asia
Autor:	Peruzzi, Lorenzo / Zarrei, Mehdi
DOI:	https://doi.org/10.5169/seals-879185

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Typification of some critical taxa of *Gagea* Salisb. (Liliaceae) from W Asia

Lorenzo Peruzzi & Mehdi Zarrei

Abstract

PERUZZI, L. & M. ZARREI (2007). Typification of some critical taxa of *Gagea* Salisb. (Liliaceae) from W Asia. *Candollea* 62: 237-244. In English, English and French abstracts.

Gagea amblyopetala var. *elata* A. Terracc., *Gagea aitchisoniana* A. Terracc., *Gagea linearifolia* A. Terracc. (≡ *Gagea sintenisii* Pascher) and *Gagea foliosa* var. *micrantha* Boiss. (basionym of *Gagea micrantha* (Boiss.) Pascher) are typified. Each type is illustrated and discussed. The taxonomic value of these taxa is discussed.

Key-words

LILIACEAE – *Gagea* – Typification

Résumé

PERUZZI, L. & M. ZARREI (2007). Typification de quelques taxons critiques de *Gagea* Salisb. (Liliaceae) de l'Asie de l'Ouest. *Candollea* 62: 237-244. En anglais, résumés anglais et français.

Gagea amblyopetala var. *elata* A. Terracc., *Gagea aitchisoniana* A. Terracc., *Gagea linearifolia* A. Terracc. (≡ *Gagea sintenisii* Pascher) et *Gagea foliosa* var. *micrantha* Boiss. (basionyme de *Gagea micrantha* (Boiss.) Pascher) sont typifiés. Chaque type est illustré et discuté. La valeur taxonomique de ces taxa est discuté.

Addresses of the author: LP: Dipartimento di Biologia, Unità di Botanica generale e sistematica, Università di Pisa, 56126, Pisa, Italy.

Email: lperuzzi@biologia.unipi.it

MZ: Department of Biology, Faculty of Science, University of Tehran, PO Box 14155-6455, Tehran, Iran and Department of Biology, Faculty of Science, Shahed University of Tehran, PO Box 18151-159, Tehran, Iran.

Submitted on February 26, 2007. Accepted on September 24, 2007

Gagea amblyopetala var. *elata* A. Terracc. in Bull. Herb. Boissier ser. 2, 6: 118. 1906.

Lectotypus (here designated): TURKEY: Gandali, mt. près Andiamyi (Asie M^e), près de la neige, 11.VI.1833, *Gustav Cocquebert de Montbret s.n.* (FI-Webb! n° 183378, individual on the right) (Fig. 1).

In describing *G. amblyopetala* var. *pumila* A. Terracc. and *G. amblyopetala* var. *elata* A. Terracc. as two tallest variants of *G. amblyopetala* s.str., TERRACCIANO (1906) affirms that many intermediate forms occur among these two taxa. However, he affirms also that “typical” var. *elata* can be recognised in two specimens: “Ras-Beyrut (Peyron s.n.!)” and “Gandali (herb. Webb)”. We traced the latter herbarium sheet, which is doubtless original material, in the Herbarium Webb, conserved at FI. Among *Gagea* individuals, there is also (on the extreme left) a plant referable to another taxon (possibly *Ornithogalum* sp.).

Taxonomic discussion. – On the basis of the typical collection, it seems that this variety is synonymous with *G. amblyopetala* Boiss. & Heldr.

Gagea aitchisoniana A. Terracc. in Boll. Soc. Ort. Palermo 2(4): 5-6. 1904.

Lectotypus (here designated): AFGHANISTAN: [illeg.], 12.VI.1880, Aitchison 828 (K!, individual on the right) (Fig. 2).

In the protologue (TERRACCIANO, 1904b), only one collection is cited: “Ho esaminato gli esemplari esistenti col n°828 nell’erbario di Kew, raccolti da I. E. Aitchison il 12 giugno 1880 nel suo viaggio attraverso l’Affghanistan”. We traced this herbarium specimen at K; and one well preserved individual of this collection is here designated as the lectotype.

Taxonomic discussion. – TERRACCIANO (1904b) considers this species as related to *G. reticulata* group. This species complex grows usually in drier part of the distribution range. The distinguishing characters for this group are so few that make it too difficult to handle them. *Gagea aitchisoniana* appears closely related to *G. setifolia* Baker and *G. anonyma* Rech. f. Indeed, WENDELBO & RECHINGER (1990) cited this species as a synonym of *G. setifolia*. *Gagea aitchisoniana* has shorter pedicels and broader tepals in relation with *G. setifolia*. The types for both species have been collected from Afghanistan by Aitchison during his trip. New investigation reveals the morphological variability in *G. setifolia* complex (incl. *G. anonyma*, *G. tehranica* Gand. *G. perpusilla* Pascher) is very high and there are no correlations between different morphological forms. It seems that all belong to one species (M. Zarrei, unpubl. data).

Gagea linearifolia A. Terracc. in Boll. Soc. Ort. Palermo 2(3): 7. 1904.

= *G. sintenisii* Pascher in Sitzungber. Deutsch. Naturwiss.-Med. Vereins Böhmen “Lotos” Prag 24: 125. 1904.

Lectotypus (here designated): IRAQ: Mesopotamia, Biredjik, in declivibus ad Euphratem, 12.III.1888, *Sintenis* 123 (LD! n°1038046, complete individual at the centre bottom) (Fig. 3).

In the protologue (TERRACCIANO, 1904a), only one collection is cited: “Mesopotamia, Biredjik, in declivibus ad Euphratem (18 martio 1889, leg. Sintenis! Iter orientale 1889; exs. N.°123, sub *G. Liotardi*, in herb. *Vindobonensi*)”. Unfortunately, this collection is no more present in Vienna herbaria, neither at W nor at WU. Consequently, we asked for information on duplicates to all those herbaria which notoriously hold Sintenis collections. We found only three specimens, at BHUPM!, G! and LD! As already noted by PERUZZI & TISON (2006), *Sintenis* n° 123 collection is heterogenous, merging several taxa. Indeed, TERRACCIANO (1904a) explicitly affirms that in the same collection (but in Berlin Herbarium) he found part of the material seen by the author when describing *G. syriaca* A. Terracc. For this reason, the choice of the lectotype of the name *G. linearifolia* was not so obvious. We selected in the herbarium sheet at LD, a single complete specimen well fitting with the protologue description.

Taxonomic discussion. – The selected lectotype fully confirms the close relationship (likely synonymy) between *G. linearifolia* and *G. luteoides* Stapf, as already suggested by RIX (1984) and LEVICHÉV (1990).

Gagea foliosa var. *micrantha* Boiss., Fl. Orient. 5: 205. 1881.

= *Gagea micrantha* (Boiss.) Pascher in Sitzungber. Deutsch. Naturwiss.-Med. Vereins Böhmen “Lotos” Prag 24: 122. 1904.

Lectotypus (here designated): ISRAEL: In summis alpinis humosis prope nives deliquescentes, alt. 8-9000 ped., 6.VI.1855, Kotschy Iter Syriacum 1855: 245 (G-BOIS! second individual on the left bottom; iso-: K!, individual on the left) (Fig. 4).

In the protologue (BOISSIER, 1881), cited “Habitat ad cedros et alibi (Bl.!), in Hermone Antilibani ad nives deliquescentes 8-9000’ (Ky. 245 sub *G. Billardieri*!), in subalpinis montis Argaei Cappadociae 8500’ (Ky. Suppl. 274!)”. We have found at G-BOIS all the herbarium material related to the name *G. foliosa* var. *micrantha*. A complete individual of the Kotschy’s collection n° 245 is here selected as the lectotype. A duplicate of the same collection was found at K. Incidentally, this herbarium collection is cited also by PASCHER (1904), who

elevated this taxon to specific rank. HEYN & DAFNI (1977) cited this same herbarium sheet as an isosyntype of *G. micrantha*. The syntypical Kotschy's collection n° 274 clearly represents instead a different taxon, belonging to sect. *Fistulosae* (Pasch.) Davlian., which conflicts with the protologue of *Gagea foliosa* var. *micrantha*.

Taxonomic discussion. – *Gagea micrantha* is a plant with two not-fistulose basal leaves, likely belonging to sect. *Didymobulbos* (K. Koch) Boiss. According to HEYN & DAFNI (1977) it is characterized one-sided inflorescence and distinctly pilose perianth. It is noteworthy to say that the collection here designated as the lectotype is also an isosyntype of the name *G. syriaca* A. Terracc. However, according to the lectotypification effected by PERUZZI & TISON (2006), the latter name is heterotypic and applicable to plants with single hollow basal leaf, close to *G. fragifera* (Vill.) Ehr. Bayer & G. López (sect. *Fistulosae*).

Acknowledgements

Curators of BHUPM, BREM, FI, G, GH, HAC, K, PAL, VT, W, WAG, WU herbaria are gratefully acknowledged. Kind thanks are due to Dr. Dimitar Uzunov (University of Calabria, Italy) for information about Sintenis specimens at W and WU herbaria.

References

- BOISSIER, E. (1881). *Gagea* Salisb. *Fl. Orient.* 5: 203-211.
- HEYN C. H. & A. DAFNI (1977). Studies in the genus *Gagea* (Liliaceae) II, The non-platyspermous species from the Galilee, the Golan Heights and Mt. Hermon, *Israel J. Bot.* 26: 11-22.
- LEVICHEV, I. G. (1990). The synopsis of the genus *Gagea* (Liliaceae) from the Western Tien-Shan. *Bot. Zhurn. (Moscow & Leningrad)* 75: 225-234.
- PASCHER, A. (1904). Übersicht über die Arten der Gattung *Gagea*. *Sitzungber. Deutsch. Naturwiss.-Med. Vereins Böhmen "Lotos"* Prag 24: 109-131.
- PERUZZI, L. & J.-M. TISON (2006). Typification of the names and taxonomic status of six taxa of *Gagea* Salisb. (Liliaceae) conserved at Firenze (FI). *Candollea* 61: 293-303.
- RIX, E. M. (1984). *Gagea* Salisb. In: DAVIS, P. H. (ed.), *Fl. Turkey* 8: 312-327.
- TERRACCIANO, A. (1904a). *Gagearium novarum* diagnoses. *Boll. Soc. Ortic. Palermo* 2(3): 3-10.
- TERRACCIANO, A. (1904b). Per la priorità delle mie *Gagearium novarum* diagnoses. *Boll. Soc. Ortic. Palermo* 2(4): 1-7.
- TERRACCIANO, A. (1906). *Gagearium* species florae orientalis. Pars secunda. *Bull. Herb. Boissier* ser. 2, 6: 105-120.
- WENDELBO, P. & K. H. RECHINGER (1990). *Gagea*. In: RECHINGER, K. H. (ed.), *Fl. Iran.* 165: 13-57.



Fig. 1. – Lectotypus of the name *Gagea amblyopetala* var. *elata* A. Terracc.
[Gustav Cocquebert de Montbret s.n., Fl] [© Museo di Storia Naturale, Florence. Reproduced with permission]




Fig. 2. – Lectotypus of the name *Gagea aitchisoniana* A. Terracc.

[Aitchison 828, K] [© The Board of Trustees of the Royal Botanic Garden, Kew. Reproduced with permission]




Fig. 3. – Lectotypus of the name *Gagea linearifolia* A. Terracc.
[Sintenis 123, LD] [© Botanical Museum, Lund. Reproduced with permission]




Fig. 4. – Lectotypus of the name *Gagea foliosa* var. *micrantha* Boiss.
[Kotschy 245, G-BOIS] [© Conservatoire et Jardin botaniques de la Ville de Genève]

