Zeitschrift: Candollea: journal international de botanique systématique =

international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 63 (2008)

Heft: 1

Artikel: Chloroplast and microsatellite markers in Astronium urundeuva

(Allemão) Engl. and close species of Anacardiaceae: toward the

definition of a species complex?

Autor: Caetano, Sofia / Nusbaumer, Louis / Naciri, Yamama

DOI: https://doi.org/10.5169/seals-879224

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 18.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Chloroplast and microsatellite markers in Astronium urundeuva (Allemão) Engl. and close species of Anacardiaceae: toward the definition of a species complex?

Sofia Caetano, Louis Nusbaumer & Yamama Naciri

Abstract

CAETANO, S., L. NUSBAUMER & Y. NACIRI (2008). Chloroplast and microsatellite markers in Astronium urundeuva (Allemão) Engl. and close species of Anacardiaceae: toward the definition of a species complex? *Candollea* 63: 115-130. In English, English and French abstracts.

Tree species belonging to genus Astronium Jacq. (Anacardiaceae R. Br.) can be sometimes difficult to identify on the sole basis of the morphological traits of the leaves. Individuals collected from Paraguay and Argentina were a priori identified in the field as Astronium urundeuva (Allemão) Engl., Astronium balansae Engl. and Astronium fraxinifolium Spreng., although no flowers nor fruits were found at the time of sampling. These were analysed with six microsatellites and two chloroplast markers. Using a bayesian approach on the microsatellite data, each individual was assigned to one of three distinct clusters that subsequently appeared to correspond to the previously described species. Astronium fraxinifolium was identified as the most differentiated species with both microsatellites and chloroplast data, although its leaf morphology is close to the one of Astronium urundeuva. The lower differentiation levels reported among Astronium urundeuva and Astronium balansae, despite their different leaf morphologies, was attributed to a more recent divergence of

Résumé

CAETANO, S., L. NUSBAUMER & Y. NACIRI (2008). Chloroplaste et marqueurs microsatellites chez Astronium urundeuva (Allemão) Engl. et autres espèces apparentées d'Anacardiaceae: vers une définition du complexe d'espèces? *Candollea* 63: 115-130. En anglais, résumés anglais et français.

Les espèces d'arbres appartenant au genre Astronium Jacq. (Anacardiaceae R. Br.) sont parfois difficiles à identifier sur la seule base des caractères morphologiques des feuilles. Des individus, a priori identifiés lors de la récolte comme étant Astronium urundeuva (Allemão) Engl., Astronium balansae Engl. et Astronium fraxinifolium Spreng., bien que ne présentant ni fleurs ni fruits au moment de l'échantillonnage, ont été collectés au Paraguay et en Argentine. Ils ont été analysés à l'aide de six marqueurs nucléaires microsatellites et de deux intergènes chloroplastiques. Sur la base des résultats obtenus sur les microsatellites, et en utilisant une approche bayésienne qui ne fait aucune utilisation de l'appartenance spécifique supposée, chaque individu a été assigné à l'un des trois groupes désignés par l'analyse. Ces trois groupes se sont avérés être exactement superposables aux trois espèces supposées. Astronium fraxinifolium est l'espèce la plus différenciée des deux autres, à la fois pour les microsatellites et les marqueurs chloroplastiques, bien que la morphologie de ses feuilles soit très

Addresses of the authors: Laboratoire de Systématique et de Biodiversité, Unité de Phylogénie et de Génétique Moléculaires, Conservatoire et Jardin botaniques de la Ville de Genève, chemin de l'Impératrice 1, CH-1292 Chambésy.

 $\label{prop:mail} Email~(YN): \textbf{yamama.naciri@ville-ge.ch}$

Submitted on 27 August 2007. Accepted on 16 January 2008.

the species or to a case of introgressive hybridization. This study underlines the advantages of using a Bayesian assignment procedure whenever molecular studies based on one particular species is to be undertaken. It furthermore confirms the species status of the three analysed taxa, and represents a first step toward a more comprehensive analysis of the three species.

Key-words

ANACARDIACEAE – Astronium – Species complex – Chloroplast markers – trnH-psbA – trnS-trnG – Nuclear microsatellites – Bayesian statistics

proche de celle d'Astronium urundeuva. La plus faible différentiation génétique trouvée entre Astronium urundeuva et Astronium balansae, bien que leurs feuilles soient assez différentes morphologiquement, est expliquée par une divergence plus récente des deux espèces ou à des phénomènes d'introgression et d'hybridation entre espèces. Cette étude souligne l'intérêt des procédures d'assignement Bayésiennes lorsque des études moléculaires sur une espèce doivent être entreprises. Elle confirme par ailleurs le statut d'espèce des trois taxons analysés, et constitue un premier pas en direction d'une analyse plus poussée de ces espèces.

Introduction

The conceptual recognition of species constitutes the starting point for many detailed investigations (MALLET, 1995), such as biodiversity assessments, red list publications, establishment of conservation strategies or genetic studies focusing on the evolutionary processes within well-defined species. Because different approaches can be used in species definition, alternative concepts co-exist, among which the biological concept of Mayr, the Hennigian concept, the phylogenetic concept or the evolutionary concept (WHEELER & MEIER, 2000). The basic procedure of accurate and efficient delimitation of species relies on the identification and comparison of diagnostic morphological or chemical characters, which are inferred to be invariant or to vary within a specified range among samples of different geographical origins (WIENS & Servedio, 2000). In practice both uncertainty and controversy about how to recognize species nevertheless exist, and are mostly related to four different issues: the concept of species presently used, the intrinsical properties of the species, the processes liable for their existence, and the methods used for inferring species boundaries (QUEIROZ, 2005).

Population researchers are frequently interested in describing the genetic structure of a group of populations more or less connected by gene flow, and in such a case, it is of major interest to ensure that no other reproductive entity is present in the data set to be analysed. DNA barcoding has been invoked to provide a rapid and reproducible identification of the species (Chase & al., 2005; Chase & al., 2007; Kress & Erickson, 2007), but much debate about its performance still exists and this method is not yet widely accepted. Accordingly, DNA barcoding, as currently applied, is unsuitable in particular situations where closely related species coexist and/or hybridization and introgression is occurring. Natural hybridization and introgression are indeed common phenomena of evolutionary importance that retain much of the researchers' attention across many fields within biology (ARNOLD, 1992; ANDER-SON & THOMPSON, 2002).

Due to the continuous improvement of the techniques for developing species-specific markers (Selkoe & Toonen, 2006) and to the recent development of numerous new statistical analyses, such as bayesian assignment techniques (Pritchard & al., 2000; Anderson & Thompson, 2002), other methods allowing the identification of related species have been introduced (Duminil & al., 2006). Hence, whenever nuclear microsatellite markers are available in a group of congeneric species, the correspondence between genotypic clusters and the previously described taxonomic species can be easily checked. Additionally, correspondence between nuclear microsatellite genotypes and other sources of genetic information, such as chloroplast haplotypes, can be very useful in cases of introgression. In this study, we aim at validating the

strategy proposed by Duminil & al. (2006) within the genus *Astronium* Jacq. (*Anacardiaceae*). Based on molecular tools, we seek to differentiate three congeneric species, *A. urundeuva* (Allemão) Engl., *A. balansae* Engl. and *A. fraxinifolium* Spreng., and to detect possible introgression among them. No clear agreement exists concerning the species name of *A. urundeuva* and *A. balansae*, with two recent revisions that either classified the species into *Astronium* genus (e.g., Muñoz, 1990) or moved it into the genus *Myracrodruon* Allemão (Santin & Leitão-Filho, 1991). Several aspects concerning the phylogenetic relationships among the two genus are still unresolved, so it is difficult to know which classification is the most suitable, biologically speaking. We will accordingly use the *Astronium* genus name in the following study.

A total of 13 Astronium species have been described in Central and South America, distributed from Mexico and the Antilles to Argentina (Muñoz, 1990). Astronium urundeuva is widespread in Eastern and central Brazil, Bolivia, Paraguay and Northwest Argentina. Astronium balansae has been described in Paraguay and Northwest Argentina, whereas A. fraxinifolium has been found from Central Brazil to the Eastern Bolivia and Western Paraguay. Close ecological requirements have been found for A. urundeuva and A. fraxinifolium, and because they also share similar distribution patterns (Fig. 1), co-occurrence has been reported in different patches of dry forests (e.g. Monte Occidental and Parque Chaqueño in Paraguay (Muñoz, 1990) or southwest Brazil (Oliveira-Filho, pers. comm.). On the other hand, A. balansae is rather reported in more dense and less dry forests, although some overlapping distribution exists in Paraguay with the two former species in the most humid parts of Parque Chaqueño or with A. urundeuva in Parque del Rio Paraguay. All three species are dioecious, wind or insect pollinated and produce small seeds that are wind dispersed. In Paraguay and North Argentina, the flowering period overlaps for the three species during September and October.

For a taxonomist, species differentiation is ideally based on floral characteristics (Muñoz, 1990), but in the absence of flowers and/or fruits, which often happens when sampling for population genetic purposes, identification of the species is still possible using a combination of several other characters. In the particular case of these three Astronium species, vegetative criteria often overlap (e.g. A. urundeuva is 5-30 m high, A. balansae 6-25 m and A. fraxinifolium 3-16 m), and the morphological traits of the leaves are the most useful character for recognizing the different species when the trees are in a vegetative state. After Muñoz (1990) and after a comparative survey of herbarium samples in the Geneva collection, the leaves of A. urundeuva measure 10-30 cm long and the leaflets are ovate to oblong-ovate, entire to slightly crenulated-dentate, shortly petiolulate (0.2-0.4 cm long excepted terminal one reaching 2 cm long), subacute to




Fig. 1. – Geographical range of the three Astronium species (○: A. urundeuva (Allemão) Engl.; △: A. fraxinifolium Spreng.; ■: A. balansae Engl.). In the detailed map, the same symbols described before for the three species have been kept: their distribution is represented by white symbols and the seven populations used in this study appear in black.

acuminate and mucronate at the apex, concolour to slightly discolour and generally pubescent. The leaves of *A. fraxinifolium* measure 12-30 cm long and the leaflets are ovate to triangular or oblong to ovate, entire to slightly undulatedentate, shortly petiolulate (0.2-0.4 cm), slightly acuminate, concolour to discolour and glabrous to hairy especially at the base of the main nerve. The leaves of *A. balansae* are 10-17 cm long and the leaflets are oblique lanceolate, distinctly dentate, long petiolulate (0.3-1.2 cm), acuminate at the apex, distinctly discolour and glabrous.

Despite the clear identification key for adult *Astronium* individuals, confusion can still persist due to the plasticity inherent to each species, depending on the geographic location, age of the tree and age of the selected branches, which can lead to possible sampling errors, especially when one single species is the object of the study. In this perspective, the risk of confusion

among *A. urundeuva* and *A. fraxinifolium* is indeed considerable, because their leaves are very alike. This risk is furthermore increased by their similar geographical range and ecological requirements. On the other hand, recognition of *A. balansae* is generally easier (morphologically distinct leaves and occurrence in a different habitat), and confusion is more uncommon.

Because of their chemical composition, *Astronium* species have been investigated for pharmacological ends and several properties have been identified: *A. urundeuva* has been screened for immunomodulatory activity (Deharo & al., 2004), tested as an antioxidant agent acting in anti-inflammatory processes (Desmarchelier & al., 1999), and proved to be successful in treating gastrointestinal transit disturbances (Menezes & Rao, 1988). *Astronium balansae* revealed the presence of bactericidal components (Salvat & al., 2004). Moreover, this genus is traditionally known for its very good timber quality, which

has resulted in the exploitation of several species for commercial purposes (BARANY & al., 2003). In this perspective, it seems obvious that a clear identification of the species is fundamental, either for pharmacological researches, or for the implementation of conservation strategies, even in the absence of flowers.

In this study we used Bayesian assignment analysis of multilocus nuclear genotypes in different populations of *Astronium* sp. in Paraguay and Argentina, to identify genotypic clusters. Recognition of *A. urundeuva*, *A. balansae* and *A. fraxinifolium* individuals was systematically done *a priori*, based on the morphological traits of the leaves, but this information was only used *a posteriori* to check for the consistency of the assignment results. Moreover, we examined the correspondence between nuclear and chloroplast molecular markers to search for possible cases of introgression.

Material and Methods

Astronium individuals were a priori identified on the basis of morphological characteristics of the trees, in particular their leaves. Identification of the three species was done in the field by local botanists: Dr. D. Prado, from the Rosario University

(Argentina) for *A. balansae*, and K. Elizeche, forest engineer at the Asuncion University (Paraguay) for both *A. fraxinifolium* and *A. urundeuva*.

A total of 165 non flowering adult individuals were sampled within seven populations:

- 1. 33 A. fraxinifolium from two populations in Paraguay;
- 2. 19 A. balansae from one Argentinean population;
- 3. 113 *A. urundeuva* from four populations in Paraguay (Table 1 and Fig. 1).

Genomic DNA was extracted with the DNeasy Plant Kit (Qiagen) and used as template for both chloroplast and microsatellite markers amplification.

Microsatellite markers

Each individual was genotyped with six nuclear microsatellites (Table 2) developed from *Astronium urundeuva: Auru.* A392, *Auru.* B209, *Auru.* D094, *Auru.* D167, *Auru.* D282 (CAETANO & al., 2005) and *Auru.* H207 (CAETANO & al., 2008). Analyses were performed on an ABI377 automated sequencer using Genescan software (Applied Biosystems), and Genescan-400 Rox (Applied Biosystems) as size standard for each individual PCR product.

Table 1. - Description of the seven Astronium populations used in this study (Nb Ind.: number of individuals).

Population	Code	Longitude	Latitude	Nb Ind.	ID Species
San Luis	Pa_StL	-57.439	-22.624	46	A. urundeuva
Cerro Léon	Pa_CLe	-60.317	-20.432	33	A. urundeuva
Altos	Pa_Alt	-57.238	-25.257	14	A. urundeuva
Cordillera	Pa_Cor	-56.420	-25.117	20	A. urundeuva
Paso de Patria	Ag_PsP	-59.815	-22.604	19	A. balansae
Loma Plata	Pa_LmP	-59.717	-22.490	13	A. fraxinifolium
Laguna Capitan	Pa_LgC	-58.610	-27.330	20	A. fraxinifolium

Table 2. – Characterization of the six microsatellite loci over the seven populations and the three Astronium species ($N_{A:}$ number of alleles; Allele Size: size of alleles [in base pairs]; $H_{T:}$ total expected heterozygosity; $F_{IS:}$ average allelic correlation within individuals relative to each population averaged over populations; $F_{ST:}$ measure of genetic divergence among the seven populations; **: p < 0.01, ***: p < 0.001).

Locus	GeneBank	N _A	Allele Size	H _T	F _{IS}	F _{ST}
Auru.A392	AY640260	14	178-214	0.823	0.003	0.308***
Auru.B209	AY509817	19	186-236	0.873	-0.004	0.280***
Auru.D094	AY640267	8	105-127	0.697	-0.105	0.448***
Auru.D167	AY640268	3	134-138	0.551	-0.166	0.277***
Auru.D282	AY640270	33	177-245	0.945	0.066	0.139***
Auru.H207	AY509818	19	127-173	0.848	0.173**	0.271***

Deviations from Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium were tested with 10000 iterations using ARLEQUIN 3.11 software (Excoffier & al., 2005). Genetically homogeneous groups of individuals were detected using a Bayesian-based approach, implemented in the Structure program (PRITCHARD & al., 2000). With Structure, the number of clusters K was tested in the range of one to seven (i.e. the total number of populations studied), assuming the admixture model with correlated allele frequencies. Ten independent runs for each value of K were performed, using a burn-in and MCMC lengths of 30000 and 100000 iterations, respectively. Convergence was achieved in all runs. The number of clusters was determined by calculating the $\triangle K$ statistic (EVANNO & al., 2005) and samples were then placed into the cluster for which they showed the highest assignment probabilities, as far as this probability was higher than 50%. Differentiation among the identified clusters, as well as between the seven populations was estimated using pairwise $F_{\rm ST}$ and computed in ARLE-QUIN 3.11 software.

A hierarchical analysis of molecular variance was also used to estimate the fraction of variance due to differences among clusters, among populations within clusters and among populations (Excoffier & al., 1992). In order to estimate null allele frequencies within each identified cluster, we then estimated the maximum-likelihood allele frequencies from the observed data, using 10000 iterations and an Expectation-Maximization (EM) algorithm implemented in ARLEQUIN. Measures of diversity were calculated within population and within cluster: number of polymorphic loci ($N_{\rm PL}$), observed number of alleles ($N_{\rm A}$) and private alleles ($N_{\rm PA}$), observed heterozygosity ($H_{\rm O}$) and expected

heterozygosity ($H_{\rm E}$) corrected for small samples. Allelic richness ($R_{\rm S}$) was computed using the rarefaction method implemented in FSTAT software (GOUDET, 2007).

Chloroplast Markers

The chloroplast DNA (cpDNA) polymorphisms were analysed by sequencing two different spacers in all *Astronium* spp. individuals: the HA locus using *trn*H and *psb*A primers, the SG locus using *trn*S and *trn*G (HAMILTON, 1999). PCR products were sequenced from both ends using BigDye Terminator v3.1 Cycle Sequencing Kit and an ABI 377 automated sequencer (Applied Biosystems). Each haplotype sequence was deposited in GenBank (Table 3). Nucleotide sequences were aligned using Clustal W (THOMPSON & al., 1994) implemented in the BIOEDIT program (HALL, 1999) and revised manually.

Genotypic linkage among loci was tested with the ARLE-QUIN package (Excoffier & al., 2005), and the results allowed the combination of the two haplotypes for each individual. In order to visualize the relationships between the combined haplotypes, a median-joining network was drawn using Network software (Bandelt & al., 1999). Insertion-deletion events (indels), as well as microsatellites, were included in the analysis, as their usefullness in intrageneric studies has already been shown (Hamilton & al., 2003; Ingvarsson & al., 2003).

Table 3. – Characterization of chloroplast haplotypes found for *trnH-psbA* and *trnS-trnG* spacers (S: substitution; ID: indel; mutation, including indels of one base pair, and indel positions, are numbered from the end of the *trnH* and *trnS* primers, respectively).

							trr	nH-psbA	(607bp)								
Haplotype	Gene Bank	ID 36	\$ 60	S 147	ID 192	S 218	S 273	\$ 295	\$ 342	\$ 352	\$ 399	ID 415	S 467	\$ 481	\$ 500	\$ 510	ID 535	ID 540
Α	EF513743	-	T	T	_	Α	С	Α	T	Α	T	_	С	G	Α	Α	G	
В	EF513744	С	T	T	- HERESTONIA - 1	Α	С	Α	T	Α	T	-	С	G	Α	Α	G	-
C	EF513745	С	T	T	_	Α	С	Α	G	Α	T	_	С	G	Α	Α	G	_
D	EU053211	-	С	G	T	T	Τ	G	T	С	G	Α	G	С	С	G	_	Α

	trnS-trnG (650bp)							
Haplotype	Gene Bank	\$ 108	S 230	ID 253	\$ 308	\$ 468	\$ 601	
Α	EF513746	С	С	Α	T	G	G	
D	EU053212	Α	Α	_	С	T	T	

Results

Bayesian assignment tests indicated that the most likely number of clusters was K = 3, as suggested by the distribution of ΔK values (Appendix 1, supplementary data). Assignment probabilities were very high, with 98% of the values being higher than 0.9. Four individuals showed lower performances (assignment probabilities to cluster A ranging between 0.616 to 0.878) but they still displayed a sharp difference with the second best cluster (between 0.081 and 0.381) and could, therefore, still be easily assigned (Appendix 2).

Our results indicated complete correspondence between taxonomic species and genotypic clusters: cluster A included all 113 individuals a priori identified as A. urundeuva, and clusters B and C corresponded respectively to a priori identified as A. balansae and A. fraxinifolium. Differentiation between the three Bayesian drawn clusters was very high ($F_{\rm CT}=0.344$; p<0.01) accounting for 90% of the total differentiation overall samples ($F_{\rm ST}=0.383$; p<0.0001). This is a further indication that the three clusters correspond to distinct biological entities, for which the average differentiation between conspecific populations was comparatively small ($F_{\rm SC}=0.059$, p<0.0001). Astronium fraxinifolium was the most differentiated cluster/species among the three, measured in terms of pairwise $F_{\rm ST}$, which is also reflected by the differential sharing of alleles among clusters (Table 4). The highest frequency of null

alleles (i.e. alleles that do not amplify because of mutations in the primer regions) was observed in *A. fraxinifolium* (0.169 \pm 0.074 averaged over loci, *A. urundeuva*: (0.027 \pm 0.043), *A. balansae* (0.026 \pm 0.042). Accordingly, clusters corresponding to *A. urundeuva* and *A. balansae* were found at Hardy-Weinberg equilibrium, as evidenced by their non significant $F_{\rm IS}$, but the opposite was true for *A. fraxinifolium* (Table 5).

Analysing the clusters/species separately, differentiation among populations within *A. fraxinifolium* cluster ($F_{\rm ST}=0.088$; p<0.0001) was about twofold the level reported among *A. urundeuva* populations ($F_{\rm ST}=0.048$; p<0.0001, the pairwise values varying between 0.021 and 0.075). $F_{\rm ST}$ could not be computed for *A. balansae*, as it was represented by a single population. In terms of diversity, *A. urundeuva* was the most diverse species (Table 5), as reflected by the highest diversity indices, namely $R_{\rm S}$ and $H_{\rm E}$.

Four chloroplast haplotypes were detected with *trn*H-*psb*A spacer (HA) and only two with *trn*S-*trn*G (SG, Table 3), which resulted in four combined haplotypes (Fig. 2). Based on consensus sequences, the HA fragment was 607bp long, and the haplotypes were characterized by four indels of 4, 5, 7 and 13bp, one mononucleotide repeat displaying two variants and twelve nucleotide substitutions. For SG, which corresponded to a 650bp fragment, the two haplotypes were characterized by five nucleotide substitutions, and two variants of a mononu-

Table 4. – Pairwise F_{ST} (below the diagonal) and number of alleles shared among each cluster of *Astronium* Jacq. (above the diagonal), estimated with six microsatellite loci. All F_{ST} values were highly significant (p < 0.001).

	A. urundeuva	A. balansae	A. fraxinifolium
A. urundeuva		15	9
A. balansae	0.295	·····································	2
A. fraxinifolium	0.346	0.631	

Table 5. – Variability indices in the three *Astronium* species based on microsatellite and chloroplast data (N_l , N_p : number of individuals and number of populations; N_{PL} , N_A , N_{PA} , N_{NA} : total number of polymorphic loci, of alleles, of private alleles, and of null alleles respectively \pm standard deviation; R_S : mean allelic richness averaged over populations \pm standard deviation; H_C , H_C : observed and expected heterozygosities averaged over populations \pm standard deviation; H_C : the average allelic correlation within individuals relative to each species, all populations confounded and corresponds to a measure of inbreeding; H_C : measure of genetic divergence among populations; for the chloroplast data the number of individuals displaying the combined haplotypes AA, BA, CA and DD is indicated (the first letter refers to trnH-psbA and the second one to trnS-trnG); trnB-trnB: trnB-

	Microsatellite data										Ch	loropl	ast da	ta	
	N _I	N _P	N_{PL}	N _A	N _{PA}	N _{NA}	R _S	Ho	H _E	F _{IS}	F _{ST}	AA	BA	CA	DD
A. urundeuva	133	4	6	71	49	0.027 ± 0.043	8.2 ± 4.2	0.701 ± 0.119	0.723 ± 0.140	-0.003	0.048***	52	59	2	-
A. balansae	19	1	3	17	4	0.026 ± 0.042	3.3 ± 3.4	0.270 ± 0.301	0.278 ± 0.299	0.029	_	_	19	_	_
A. fraxinifolium	33	2	4	28	20	0.169 ± 0.074	4.3 ± 4.2	0.375 ± 0.283	0.439 ± 0.338	0.108*	0.088**	-	-	-	33




Fig. 2. – Median Joining Network of Astronium haplotypes considering all mutations. The radius of each circle is proportional to the number of individuals displaying the haplotype and colours assigned correspond to each species (dark grey: A. urundeuva (Allemão) Engl.; diagonal cross: A. balansae Engl.; light grey: A. fraxinifolium Spreng.). Mutation names starting with HA and SG referred to mutations observed in trnH-psbA and trnS-trnG, respectively. Indels are in italics and substitutions are underlined.

cleotide repeat. We found absolute correspondence between combined haplotype DD and the cluster C, previously identified as being entirely constituted by *A. fraxinifolium* individuals. Combined haplotypes AA and CA corresponded to some *A. urundeuva* individuals, assigned to cluster A, and for the haplotype BA, correspondence was found with the remaining *A. urundeuva* individuals constituting cluster A and all *A. balansae* individuals assigned to cluster B (see Appendix 2 for details).

Discussion

A good delimitation of the three *Astronium* species was obtained using the nuclear markers and the Bayesian approach. However a major inconsistency concerning the identification of *A. urundeuva* and *A. balansae* was observed when considering the chloroplast results. Three haplotypes A, B and C, were found in the first species with the HA spacer, whereas all *A. balansae* individuals displayed haplotype B. Moreover, the locus SG could not discriminate these two species (haplotype A for both). On the other hand, whether with HA or SG, *A. fraxinifolium* displayed a new haplotype, D. Consequently, on the basis of both the higher microsatellite pairwise $F_{\rm ST}$ and the chloroplast results, *A. fraxinifolium* was the most

differentiated cluster/species, whilst a much weaker level of differentiation was reported between A. urundeuva and A. balansae, with possible confusion between the species when using the chloroplast markers alone. These results raise two additional topics in the discussion; the first concerns the sharp contradiction with the morphology, according to which A. urundeuva and A. fraxinifolium are closer to each other, and the second the status of species complex for A. urundeuva and A. balansae.

Divergence of A. fraxinifolium

Both types of markers agree well in identifying A. fraxinifolium as the most differentiated species. Given that discernment between A. fraxinifolium and A. urundeuva based on morphological traits of the leaves is tricky, and that these species share the same ecological conditions and often occur in sympatry, these results are rather reassuring. On the one hand, such results indicate that the close morphological traits of the leaves are most probable related to some selection pressure due to the same ecological features being shared among species. This is a clear example where the morphological similarity does not reflect the genetic relationships between species, as evaluated by a priori neutral markers. On the other, these results reveal that genetic structure surveys based on microsatellites (CAETANO & al., 2008) can be performed in all confidence, as delimitation of the species has been safely accomplished in places where a confusion might have had occurred (e.g. Central Brazil to Eastern Bolivia and Western Paraguay).

Relationship between A. urundeuva and A. balansae

The second issue concerns the relationship between *A. urundeuva* and *A. balansae*, and four major hypotheses can be put forward to explain the discrepancy between chloroplast and nuclear genes:

- it could result from homoplasy of this chloroplast region
 (i.e. identity by state but not identity by descent);
- it could reflect the existence of a still shared ancestral haplotypes;
- it could reflect the differential evolutionary rates of the chloroplast and nuclear polymorphisms analysed here;
- it could reflect chloroplast capture, *i.e.* the introgression of the chloroplast from one species to another through interspecific hybridization. Hybridization is indeed known to be one of the major factors leading to genetic incongruence between nuclear and cytoplasmic markers in plants (Soltis & Kuzoff, 1995), as illustrated by numerous examples reporting cases of chloroplast capture in several taxa (Soltis & Kuzoff, 1995; Comes & al., 1997; Manen & al., 2002; Okuyama & al., 2005).

While it is highly unlikely that homoplasy is in the origin of such results, especially given the nature of the mutation differentiating haplotypes HA_A and HA_B (indel of 7bp long), we are unable at this point to undoubtedly decide which of the second, third or fourth explanation is the more plausible. Since A. urundeuva and A. balansae appear to be closer to each other than to A. fraxinifolium, it can be hypothesized that they have diverged more recently than did A. fraxinifolium. In this context, the BA haplotype shared between A. urundeuva and A. balansae could indeed be viewed as an ancestral one still occurring in the two species despite their split into differentiated biological entities. Closely related to this second explanation is the differential mutation rates reported in the literature for chloroplast spacers and nuclear microsatellites that are known to evolve much faster. The rate of indel mutations in the intergenic regions of chloroplast genomes has for instance been recently estimated to be $\approx 0.8 \pm 0.04 \times 10^{-9}$ per site per generation (YAMANE & al., 2006), whilst mutation rates for nuclear microsatellites are in general substantially higher, of the order of 10⁻⁴-10⁻² per generation (JARNE & LAGODA, 1996; HANCOCK, 1999).

The fourth explanation, i.e. introgressive hybridization between A. urundeuva and A. balansae could be hypothesized, but the direction in which this event would have occurred cannot be assessed, because of the poor representation of A. balansae in the dataset. It is interesting to notice that the four individuals more poorly assigned to A. unrundeuva cluster were all secondarily assigned to A. balansae cluster, with p ranging from 0.081 to 0.381. Generally speaking, the introgression signal displayed by individual loci is still poorly understood (OKUYAMA & al., 2005) and depends on each locus evolutionary history, although it is known that chloroplast capture is generally promoted by cyto-nuclear incompatibilities and facilitated by partial selfing events under certain conditions (TSITRONE & al., 2003). In our case, such hybridization between A. urundeuva and A. balansae could be (or have been) promoted by their overlapping flowering period, the ability of pollen to be wind-dispersed and the co-occurrence of the two species in Parque del Rio Paraguay and in Parque Chaqueño, although they have different ecological requirements in terms of soil humidity.

It should be noticed finally, that the last three explanations are not exclusive since the more recent the two species are, the more chance they have to hybridize, because the barrier to gene flow is not strong enough, the more chance they have to share ancestral alleles or haplotypes, and the more chance they have that no mutation occurred since they diverged, specially with the chloroplast. In this perspective, it is reasonable to question a species complex status for *A. urundeuva* and *A. balansae*, and we therefore strongly encourage a more detailed study in light to what has been done in Europe for *Quercus petraea* (Matt.) Liebl. and *Q. robur* L. (MUIR AND SCHLÖTTERER, 2006).

This result is in agreement with the finding of Santin & Leitão-Filho (1991) that *A. urundeuva* and *A. balansae* share some ovary features, a reason why they suggested moving the two species together to *Myracrodruon* genus.

Diversity within species

A higher level of diversity was found within A. urundeuva when compared to the two other species, as reflected by both gene diversity and allelic richness. The unbalanced sampling can explain the differences observed between species. Allelic richness is specifically drawn to compare samples of different sizes, but it uses the observed allele frequencies that are better estimated for A. urundeuva than for the two other species because of a larger sampling size. A second explanation may arise from the use of specific A. urundeuva markers. Because microsatellite primers were designed from a unique A. urundeuva individual, the former result could primarily indicate some kind of ascertainment effect, according to which microsatellite amplifications are more prone to failure when used on congeneric species (TREUREN, 1998). Accordingly, higher frequencies of null alleles have been found for the most differentiated species A. fraxinifolium, whilst for A. urundeuva and A. balansae, weaker and very similar frequencies were observed. Therefore, significant biases can effectively arise if these A. urundeuva specific microsatellites are to be used in intraspecific analyses within other divergent Astronium species. This is clearly observed within A. fraxinifolium, and might also explain the higher F_{ST} that was found between the two A. fraxinifolium populations (0.088), when compared with global F_{ST} obtained for A. urundeuva (0.048), whose populations were distributed in a much wider way.

Conclusion

In this study, and contrary to most surveys of genetic variation, the a priori information about Astronium species based on morphological characters were not used to delineate the clustering of individuals. Individuals were instead grouped together blindly, i.e. with a Bayesian assignment method applied on multiple variable codominant nuclear markers. The morphological determination of each individual done in the field by local botanists was only used as an afterwards confirmation tool. The simple procedure used here proved to be very efficient in the discrimination between A. urundeuva, A. balansae and A. fraxinifolium, and showed that the leaf morphology represents well the floral characteristics, demonstrating this way the value of this character for the distinction of these species. Only few microsatellite loci were thus sufficient to achieve good assignment accuracy, probably because differentiation between the clusters was quite strong (Vähä & PRIMMER, 2006). Moreover, the high level of interspecific differentiation also justifies the use of Evannos's ΔK method, as it performs better with strong differentiation levels (WAPLES & GAGGIOTTI, 2006).

The recognition of the species based on molecular approaches has been largely criticized by taxonomists, who argue that these new identification methods, such as DNA barcoding (Chase & al., 2005; Kress & al., 2005; Chase & al., 2007; Kress & Erickson, 2007), could result in incorrect species recognition, diminishing the traditional morphologybased approaches. Because of their high specificity, the DNA barcode can not rely on microsatellites and it is instead based on the use of chloroplast markers (Chase & al., 2007). We demonstrate here that the two chloroplast locus used in this study were unable to differentiate between A. urundeuva and A. balansae, among which the HA locus that was suggested as a good barcode tool in association with a portion of the coding rbcL gene (Kress & Erickson, 2007). HA spacer is indeed known to be one of the most variable plastid region in angiosperms between and within genera (KRESS & al., 2005), which has led to its choice for DNA barcoding (CHASE & al., 2005), although it has been proved to be also variable within species (Hamilton & al., 2003; Ingvarsson & al., 2003; Naciri & Gaudeul, 2007).

The procedure employed in this study differs greatly from the classical phylogenetic reconstruction based on a few gene trees and on a restricted sampling scheme, as it uses population genetic principles based on multilocus data, obtained on a batch of individuals for each species. The increased availability of high polymorphic nuclear markers associated with the development of new powerful statistical methods greatly assists the circumscription of species, and does not compromise the traditional taxonomic approach, and we suggest that cross validation of species identification using different approaches is desirable.

Acknowledgements

We wish to thank Prof. F. Méréles from Asuncion University (Paraguay), Dr D. Prado from Rosario University (Argentina), Dr A. Oliveira-Filho from Lavras University (Brazil), Dr S. Beck from the Botanical Garden of La Paz (Bolivia), P. Silveira, K. Elizeche, M. Soloaga and L. Oakley, for their help in sampling, H. Geser, L. Turin for help at the CJBG lab, Prof. L. Excoffier and Dr E. Poloni for helpful comments during this project. We wish to thank the four referees for their helpful comments and suggestions. This work was supported by the Swiss National Foundation (grants n° 3100A0/100806-1 & 2), the Conservatoire et Jardin botaniques of Geneva and the three following Societies, which contributed to travel expenses and lab consumables (Société Académique de Genève, Société de Physique et d'Histoire Naturelle and Swiss Zoological Society).

References


- Anderson, E. C. & E. A. Thompson (2002). A model-based method for identifying species hybrids using multilocus Genetic Data. *Genetics* 160: 1217-1229.
- Arnold, M. L. (1992). Natural hybridization as an evolutionary process. *Annual Rev. Ecol. Syst.* 23: 237-261.
- Bandelt, H.-J., P. Forster & A. Röhl (1999). Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16: 37-48
- Barany, M., A. Hammett & P. Araman (2003). Lesser used wood species of Bolivia and their relevance to sustainable forest management. *Forest Prod. J.* 53.
- CAETANO, S., D. PRADO, T. PENNINGTON, S. BECK, A. OLIVEIRA-FILHO, R. SPICHIGER & Y. NACIRI (2008). The history of seasonally dry tropical forests in eastern South America: inferences from the genetic structure of the tree Astronium urundeuva (Anacardiaceae). *Mol. Ecol.* 17: 3147-3159.
- CAETANO, S., P. SILVEIRA, R. SPICHIGER & Y. NACIRI-GRAVEN (2005). Identification of microsatellite markers in a neotropical seasonally dry forest tree, Astronium urundeuva (Anacardiaceae). *Mol. Ecol. Notes* 5: 21-23.
- CHASE, M., N. SALAMIN, M. WILKINSON, J. DUNWELL, R. KESANA-KURTHI, N. HAIDAR & V. SAVOLAINEN (2005). Land plants and DNA barcodes: short-term and long-term goals. *Phil. Trans., Ser. B* 360: 1889-1895.
- CHASE, M., R. COWAN, P. HOLLINSWORTH & al. (2007). A proposal for a standardised protocol to barcode all land plants. *Taxon* 56: 295-299.
- COMES, H., J. KADEREIT, A. POHL & R. ABBOTT (1997). Chloroplast DNA and isozyme evidence on the evolution of Senecio vulgaris (Asteraceae). *Pl. Syst. Evol.* 206: 375-392.
- Deharo, E., R. Baelmans, A. Gimenez, C. Quenevo & G. Bourdy (2004). In vitro immunomodulatory activity of plants used by the Tacana ethnic group in Bolivia. *Phytomedicine* 11: 516-522.
- DESMARCHELIER, C., R. LISBOA-ROMÃO, J. COUSSIO & G. CICCIA (1999). Antioxidant and free radical scavenging activities in extracts from medicinal trees used in the "Caatinga" region in northeastern Brazil. *J. Ethnopharmacol.* 67: 69-77.
- DUMINIL, J., H. CARON, I. SCOTTI, S.-O. CAZAL & R. J. PETIT (2006). Blind population genetics survey of tropical rainforest trees. *Mol. Ecol.* 15: 3505-3513.
- EVANNO, G., S. REGNAUT & J. GOUDET (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.* 14: 2611-2620.
- EXCOFFIER, L., G. LAVAL & S. SCHNEIDER (2005). Arlequin, ver. 3.0: An integrated software package for population genetics data analysis. *Evol. Bioinf. Online* 1: 47-50.
- EXCOFFIER, L., P. E. SMOUSE & J. M. QUATTRO (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131: 479-491.

- GOUDET, J. (2007). FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3) available from the following link: [http://www.unil.ch/izea/softwares/fstat.htlm].
- HALL, T. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.* 41: 95-98.
- Hamilton, M. B. (1999). Four primer pairs for the amplification of chloroplast intergenic regions with intraspecific variation. *Mol. Ecol.* 8: 513-525.
- Hamilton, M. B., J. M. Braverman & D. F. Soria-Hernanz (2003). Patterns and relative rates of nucleotide and insertion/deletion evolution at six chloroplast intergenic regions in New World species of the Lecythidaceae. *Mol. Biol. Evol.* 20: 1710-1721.
- Hancock, J. (1999). Microsatellites and other simple sequences: genomic context and mutational mechanisms. *In*: Goldstein, D. B. & C. Schlotterer (ed.), *Microsatellites Evolution and Applications*: 1-9. Oxford University Press.
- INGVARSSON, P. K., S. RIBSTEIN & D. R. TAYLOR (2003). Molecular evolution of insertions and deletion in the chloroplast genome of Silene. *Mol. Biol. Evol.* 20: 1737-1740.
- JARNE, P. & P. J. L. LAGODA (1996). Microsatellites, from molecules to populations and back. *Trends Ecol. Evol.* 11: 424-429.
- Kress, W. J. & D. Erickson (2007). A two-locus global DNA barcode for land plants: the coding *rbcL* gene complements the non-coding trnH-psbA spacer region. *Plos One* 6(e508).
- Kress, W. J., K. J. Wurdack, E. A. Zimmer, L. A. Weigt & D. H. Janzen (2005). Use of DNA barcodes to identify flowering plants. *Proc. Natl. Acad. Sci. U.S.A.* 102: 8369-8374.
- Mallet, J. (1995). A species definition for the Modern Synthesis. *Trends Ecol. Evol.* 10: 294-299.
- MANEN, J.-F., M. BOULTER & Y. NACIRI-GRAVEN (2002). Complexity of the phylogeography of the genus Ilex. *Pl. Syst. Evol.* 235: 79-98.
- MENEZES, A. & V. RAO (1988). Effect of Astronium urundeuva (aroeira) on gastrointestinal transit in mice. *Brazil. J. Med. Biol. Res.* 21: 531-533.
- Muir, G. & C. Schlötterer (2006). Moving beyond single-locus studies to characterize hybridization between oaks (Quercus spp.). *Mol. Ecol.* 15: 2301-2304.
- MUÑOZ, J. (1990). Anacardiaceae. In: SPICHIGER, R. & L. RAMELLA (ed.), Fl. Paraguay 14. Conservatoire et Jardin botaniques de Genève.
- NACIRI, Y. & M. GAUDEUL (2007). Phylogeography of the endangered Eryngium alpinum L. (Apiaceae) in the European Alps. *Mol. Ecol.* 16: 2721-2733.
- OKUYAMA, Y., N. FUJII, M. WAKABAYASHI, A. KAWAKITA, M. ITO, M. WATANABE, N. MURAKAMI & M. KATO (2005). Nonuniform concerted evolution and chloroplast capture: heterogeneity of observed introgression patterns in three molecular data partition phylogenies of Asian Mitella (Saxifragaceae). *Mol. Biol. Evol.* 22: 285-296.
- PRITCHARD, J. K., M. STEPHENS & P. DONNELLY (2000). Inference of population structure using multilocus genotype data. *Genetics* 155: 945-959.

- QUEIROZ, K. DE (2005). Different species problems and their resolution. *Bioessays* 27: 1263-1269.
- SALVAT, A., L. ANTONACCI, R. FORTUNATO, E. SUAREZ & H. GODOY (2004). Antimicrobial activity in methanolic extracts of several plant species from northern Argentina. *Phytomedicine* 11: 230-234.
- Santin, D. & H. Leitão-Filho (1991) Restabelecimento e revisão taxonomica do género Myracrodruon Freire Alemão (Anacardiaceae). *Revista Brasil. Bot.* 14: 133-145.
- Selkoe, K. A. & R. J. Toonen (2006). Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecol. Letters* 9: 615-629.
- Soltis, D. E. & R. K. Kuzoff (1995). Discordance between Nuclear and Chloroplast Phylogenies in the Heuchera Group (Saxifragaceae). *Evolution* 49: 727-742.
- THOMPSON, J., D. HIGGINS & T. J. GIBSON (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucl. Acids Res.* 22: 4673-4680.
- TSITRONE, A., M. KIRKPATRICK & D. LEVIN (2003). A model for chloroplast capture. *Evolution* 57: 1776-1782.

- Treuren, R. van (1998). Estimating null allele frequencies at a microsatellite locus in the oystercatcher (Haematopus ostralegus). *Mol. Ecol.* 7: 1413-1417.
- VÄHÄ, J.-P. & C. R. PRIMMER (2006). Efficiency of model-based Bayesian methods for detecting hybrid individuals under different hybridization scenarios and with different numbers of loci. *Mol. Ecol.* 15: 63-72.
- Waples, R. S. & O. Gaggiotti (2006). What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Mol. Ecol.* 15: 1419-1439.
- Wheeler, Q. D. & R. Meier (2000). *Species concepts and phylogenetic theory*. Columbia University Press.
- WIENS, J. & M. SERVEDIO (2000). Species delimitation in systematics: inferring diagnostic differences between species. *Proc. Roy. Soc. Biol. Sci. Ser. B* 267: 631-636.
- Yamane, K., K. Yano & T. Kawahara (2006). Pattern and rate of indel evolution inferred from whole chloroplast intergenic regions in sugarcane, maize and rice. *DNA Res.* 13: 197-204.

Appendix 1. – Detection of the number of clusters within the seven sampled populations of *Astronium* Jacq. Left y-axis is the mean likelihood over the ten runs as a function of $K(\bullet)$, the right y-axis $\triangle K$ following EVANNO & al. (2005) as a function of $K(\circ)$.



Appendix 2. – Assignment probabilities obtained for each individual of *Astronium* Jacq. based on six microsatellite loci and chloroplast haplotypes found for *trn-psbA* (HA) and *trnS-trnG* (SG) spacers.

Ind.	M	icrosatellite da	ita	Chloroplast data			
	Cluster A	Cluster B	Cluster C	HA	SG		
Pa_StL_0001	0.994	0.003	0.003	Α	Α		
Pa_StL_0002	0.991	0.006	0.003	В	Α		
Pa_StL_0003	0.994	0.003	0.003	В	Α		
Pa_StL_0004	0.994	0.003	0.003	Α	Α		
Pa_StL_0005	0.992	0.005	0.003	Α	Α		
Pa_StL_0006	0.993	0.003	0.004	Α	Α		
Pa_StL_0150	0.993	0.004	0.003	Α	Α		
Pa_StL_0151	0.994	0.003	0.003	Α	Α		
Pa_StL_0153	0.990	0.008	0.002	Α	Α		
Pa_StL_0154	0.983	0.014	0.003	Α	Α		
Pa_StL_0007	0.993	0.004	0.002	Α	Α		
Pa_StL_0008	0.651	0.290	0.059	Α	Α		
Pa_StL_0009	0.988	0.009	0.003	Α	Α		
Pa_StL_0010	0.984	0.013	0.003	Α	Α		
Pa_StL_0011	0.993	0.005	0.002	Α	Α		
Pa_StL_0012	0.909	0.089	0.003	Α	Α		
Pa_StL_0013	0.994	0.004	0.002	Α	Α		
Pa_StL_0014	0.992	0.005	0.003	С	Α		
Pa_StL_0015	0.993	0.005	0.003	Α	Α		
Pa_StL_001 <i>7</i>	0.982	0.015	0.003	Α	Α		
Pa_StL_0018	0.993	0.004	0.002	Α	Α		
Pa_StL_0019	0.990	0.005	0.005	Α	Α		
Pa_StL_0020	0.993	0.005	0.002	С	Α		
Pa_StL_0021	0.993	0.004	0.003	В	Α		
Pa_StL_0030	0.992	0.006	0.002	В	Α		
Pa_StL_0031	0.991	0.006	0.003	Α	Α		
Pa_StL_0032	0.992	0.005	0.003	В	Α		
Pa_StL_0034	0.993	0.004	0.002	В	Α		
Pa_StL_0035	0.982	0.015	0.003	В	Α		
Pa_StL_0036	0.987	0.011	0.003	В	Α		
Pa_StL_0037	0.993	0.005	0.002	В	Α		
Pa_StL_0038	0.616	0.381	0.003	В	Α		
Pa_StL_0039	0.979	0.019	0.002	В	Α		
Pa_StL_0040	0.988	0.009	0.003	В	Α		
Pa_StL_0041	0.970	0.004	0.026	В	Α		
Pa_StL_0042	0.993	0.004	0.002	В	Α		
Pa_StL_0043	0.917	0.081	0.002	В	Α		
Pa_StL_0044	0.874	0.122	0.004	В	Α		
Pa_StL_0045	0.983	0.014	0.003	В	Α		

Ind.	M	icrosatellite da	ta	Chlorop	olast data
	Cluster A	Cluster B	Cluster C	HA	SG
Pa_StL_0047	0.993	0.005	0.002	В	Α
Pa_StL_0048	0.993	0.004	0.003	В	Α
Pa_StL_0049	0.993	0.004	0.003	В	Α
Pa_StL_0050	0.993	0.005	0.002	В	Α
Pa_StL_0051	0.992	0.004	0.005	В	Α
Pa_StL_0052	0.992	0.005	0.002	В	Α
Pa_StL_0053	0.993	0.004	0.003	В	Α
Pa_CLe_0041	0.994	0.003	0.003	В	Α
Pa_CLe_0042	0.994	0.004	0.002	Α	Α
Pa_CLe_0043	0.994	0.004	0.003	Α	Α
Pa_CLe_0044	0.993	0.005	0.002	Α	Α
Pa_CLe_0045	0.994	0.004	0.003	Α	Α
Pa_CLe_0046	0.992	0.005	0.002	Α	Α
Pa_CLe_0047	0.878	0.120	0.003	Α	Α
Pa_CLe_0048	0.994	0.003	0.003	Α	Α
Pa_CLe_0049	0.967	0.031	0.002	Α	Α
Pa_CLe_0050	0.992	0.006	0.003	Α	Α
Pa_CLe_0051	0.985	0.012	0.003	Α	Α
Pa_CLe_0052	0.991	0.006	0.003	Α	Α
Pa_CLe_0053	0.990	0.007	0.003	Α	Α
Pa_CLe_0054	0.994	0.004	0.003	Α	Α
Pa_CLe_0055	0.993	0.005	0.003	Α	Α
Pa_CLe_0056	0.994	0.002	0.003	Α	Α
Pa_CLe_0057	0.990	0.007	0.003	Α	Α
Pa_CLe_0058	0.994	0.003	0.003	Α	Α
Pa_CLe_0059	0.994	0.003	0.002	Α	Α
Pa_CLe_0060	0.994	0.003	0.003	Α	Α
Pa_CLe_0061	0.994	0.004	0.003	Α	Α
Pa_CLe_0062	0.965	0.031	0.004	Α	Α
Pa_CLe_0063	0.994	0.004	0.003	Α	Α
Pa_CLe_0064	0.987	0.008	0.004	Α	Α
Pa_CLe_0065	0.994	0.003	0.003	Α	Α
Pa_CLe_0066	0.994	0.002	0.003	Α	Α
Pa_CLe_0067	0.987	0.010	0.004	Α	Α
Pa_CLe_0068	0.994	0.003	0.003	Α	
APa_CLe_0069	0.987	0.010	0.003	Α	Α
Pa_CLe_0070	0.990	0.007	0.003	Α	Α
Pa_CLe_0071	0.966	0.004	0.030	Α	Α
Pa_CLe_0072	0.986	0.011	0.004	Α	Α
Pa_CLe_0073	0.994	0.004	0.002	Α	Α
Pa_Alt_0650	0.989	0.008	0.003	В	Α
Pa_Alt_0651	0.959	0.008	0.033	В	Α
Pa_Alt_0652	0.983	0.015	0.003	В	Α

Ind.	M	icrosatellite da	ta	Chloro	olast data
	Cluster A	Cluster B	Cluster C	HA	SG
Ag_PsP_0694	0.003	0.994	0.002	В	Α
Ag_PsP_0695	0.003	0.994	0.002	В	Α
Ag_PsP_0696	0.004	0.993	0.003	В	Α
Ag_PsP_0697	0.004	0.994	0.002	В	Α
Ag_PsP_0698	0.017	0.980	0.003	В	Α
Ag_PsP_0699	0.011	0.981	0.008	В	Α
Ag_PsP_0700	0.003	0.995	0.002	В	Α
Pa_LmP_1001	0.004	0.016	0.981	D	D
Pa_LmP_1002	0.003	0.002	0.995	D	D
Pa_LmP_1003	0.003	0.002	0.995	D	D
Pa_LmP_1004	0.003	0.003	0.994	D	D
Pa_LmP_1005	0.004	0.004	0.991	D	D
Pa_LmP_1006	0.002	0.002	0.995	D	D
Pa_LmP_1007	0.003	0.003	0.994	D	D
Pa_LmP_1008	0.003	0.003	0.994	D	D
Pa_LmP_1009	0.003	0.003	0.994	D	D
Pa_LmP_1010	0.003	0.008	0.989	D	D
Pa_LmP_1011	0.003	0.002	0.995	D	D
Pa_LmP_1012	0.003	0.002	0.995	D	D
Pa_LmP_1013	0.004	0.007	0.989	D	D
Pa_LgC_1014	0.002	0.002	0.995	D	D
Pa_LgC_1015	0.003	0.008	0.989	D	D
Pa_LgC_1016	0.002	0.002	0.995	D	D
Pa_LgC_1017	0.016	0.003	0.981	D	D
Pa_LgC_1018	0.003	0.002	0.994	D	D
Pa_LgC_1019	0.003	0.002	0.995	D	D
Pa_LgC_1020	0.003	0.002	0.995	D	D
Pa_LgC_1021	0.004	0.016	0.981	D	D
Pa_LgC_1022	0.003	0.002	0.995	D	D
Pa_LgC_1023	0.003	0.002	0.995	D	D
Pa_LgC_1024	0.003	0.003	0.994	D	D
Pa_LgC_1025	0.002	0.002	0.995	D	D
Pa_LgC_1026	0.003	0.002	0.995	D	D
Pa_LgC_1027	0.003	0.002	0.995	D	D
Pa_LgC_1028	0.003	0.002	0.995	D	D
Pa_LgC_1029	0.006	0.005	0.989	D	D
Pa_LgC_1030	0.006	0.005	0.989	D	D
Pa_LgC_1031	0.002	0.002	0.995	D	D
Pa_LgC_1032	0.003	0.003	0.994	D	D
Pa_LgC_1033	0.003	0.008	0.989	D	D