

Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	63 (2008)
Heft:	2
Artikel:	Neotypification of the name Lapiedra martinezii Lag. (Amaryllidaceae)
Autor:	Gonzalo Estébanez, Raúl / Cabezas, Francisco / Aedo, Carlos
DOI:	https://doi.org/10.5169/seals-879231

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Neotypification of the name *Lapiedra martinezii* Lag. (Amaryllidaceae)

Raúl Gonzalo Estébanez, Francisco Cabezas & Carlos Aedo

Abstract

GONZALO ESTÉBANEZ, R., F. CABEZAS & C. AEDO (2008). Neotypification of the name *Lapiedra martinezii* Lag. (Amaryllidaceae). *Candollea* 63: 205-207. In English, English and French abstracts.

In absence of original material, a neotype is designated for the name *Lapiedra martinezii* Lag. (Amaryllidaceae). The neotype is a specimen from SE Spain kept in the herbarium of the Royal Botanic Garden of Madrid (MA).

Key-words

AMARYLLIDACEAE – Lapiedra – Typification

Résumé

GONZALO ESTÉBANEZ, R., F. CABEZAS & C. AEDO (2008). Néotypification du nom *Lapiedra martinezii* Lag. (Amaryllidaceae). *Candollea* 63: 205-207. En anglais, résumés anglais et français.

En l'absence de matériel original, un néotype est désigné pour le nom *Lapiedra martinezii* Lag. (Amaryllidaceae). Ce néotype est un spécimen du SE de l'Espagne conservé dans l'herbier du Jardin Botanique Royal de Madrid (MA).

Address of the authors: Department of Biodiversity and Conservation. Real Jardín Botánico de Madrid, CSIC. Murillo Sq. 2, 28014, Madrid, Spain.
Email (RGE): rgonzalo@rjb.csic.es

Submitted on June 19, 2008. Accepted on July 31, 2008.

Introduction

Lapiedra Lag. is a monotypic genus of *Amaryllidaceae* described by LAGASCA (1816), comprising the western Mediterranean endemic species *L. martinezii* Lag. It has a wide disjunct area in S-SE Spain (ranging from the coast of Castellón to Málaga and including also some inland provinces, including Córdoba and Albacete) and an area in N Morocco, where it ranges from the coastal provinces between W Rif Mountains to the Gareb area and in Kert Ganc. This species grows in stony shrubby areas and in the fissures of calcareous rocks. Flowers usually bloom after early autumn rains, during September, and leaves fully develop after anthesis (MAIRE, 1959; JURY, 2002; AEDO, in press).

Three species were described and formerly included in this genus. *Lapiedra chilensis* Phil., distributed in South America, which was selected as the type species for the genus *Traubia* Moldenke by MOLDENKE (1963). It was lately adscribed by RAVENNA (1974) to *T. modesta* (Phil.) Ravenna. From S-SE Spain and NW Africa were also described *Lapiedra gracilis* Baker, currently synonymized to *Narcissus cavanillesi* Barra & G. López, and *L. placiana* Herb., lately identified as a heterotypical synonym of Lagasca's name (BAKER, 1888; MÜLLER DOBLIES & MÜLLER DOBLIES, 1978; AEDO, in press).

The main characters differentiating *Lapiedra* from the other genera of *Amaryllidaceae* are the leaves with a white line on the abaxial surface, the tepals slightly fused in the base, lanceolate, patens to erect-patens, white, with a green line with 3-4 nerves on the dorsal surface and anthers basifixated (BAKER, 1888; MEEROW & SNIJMAN, 1998).

While addressing the generic treatment of *Lapiedra* for *Flora Iberica*, we realized that there was no type designated for the name *L. martinezii*. The purpose of this paper is to contribute to the stability of nomenclature by the neotypification of *L. martinezii*.

Neotype of *Lapiedra martinezii* Lag.

In the protologue by LAGASCA (1816), two original locations were mentioned. The first is: "Hab. ad saxorum rimas subhumidas, in monte Arcis Saguntinae, prope Sanctuarium de la Fuen Santa juxta Algezares oppidum in Muriae Regno..." probably collected by Lagasca. Unfortunately, part of his personal herbarium was destroyed during his travel from Madrid to Gibraltar, thus some type specimens were lost (STAFLEU & COWAN, 1979). The second reference is "... atque non procul a Malacensi urbe legit acerrimus Naturae scrutator D. Felix Haenseler... Collitur in R.M.H". Haenseler was a pharmaceutical and naturalist, originally from Switzerland but brought up in Málaga, where he collected many specimens.

Lagasca and Haenseler also exchanged vouchers throughout Europe, and thus we can find specimens of their personal collections included in herbaria such as those of Boissier,

Willkomm, Webb, Agardh, Swartz or Smith (C, COI, FI, G, S). For instance, *Hemionitis pozoi* Lag. was typified from a voucher kept in Swartz's Herbarium in Stockholm (MORTON, 1959). Thus, we decided to ask for possible type material included in those major herbaria. We also checked three local Andalusian herbaria where historic collections by Lagasca and Haenseler could be kept (SALGUEIRO, 1998): GDA, MGC and SEV. All keepers kindly responded to our letters, but, unfortunately, no specimens of *Lapiedra martinezii* neither from Lagasca or Haenseler have been found.

In the absence of original material the following specimen is proposed as the neotype of *L. martinezii*, since it was collected very close to the locality mentioned by Lagasca. It also bears many diagnostic characters of this species, usually absent in most of herbarium specimens.

Lapiedra martinezii Lag., Elench. Pl.: 14. 1816.

Neotypus (here designated): **SPAIN. Murcia:** Cabo Tiñoso, Cala Cerrada, 37°32'31"N 1°9'35"W, 15 m, 9.X.2005, Aedo, C. 12305 (MA-731958) (Fig. 1).

Acknowledgements

We wish to thank the keepers of C, COI, G, GDA, FI, MGC, S, SEV herbaria for searching the herbarium material. This work was supported by the *Flora Iberica* project CGL2005-05471-C04-01, Ministerio de Educación y Ciencia, Spain.

References

- AEDO, C. (in press). Lapiedra. In: CASTROVIEJO, S. (ed.), *Fl. Iber.* 20.
- BAKER, J. G. (1888). *Handbook of the Amaryllidaceae, including the Alstroemerieae and Agaveae*. London.
- JURY, S. L. (2002). Lapiedra. In: VALDÉS, B., M. REJDALI, A. ACHHAL EL KADMIRI, S. L. JURY & J. M. MONTSERRAT (ed.), *Catalogue des plantes vasculaires du nord du Maroc* 2: 879. CSIC.
- LAGASCA, M. (1816). *Genera et species plantarum*. Madrid.
- MAIRE, R. (1959). Lapiedra. *Fl. Afrique N.* 6: 19. Lechevalier.
- MEEROW, A. W. & D. A. SNIJMAN (1998). Amaryllidaceae. In: KUBITZKI, K. (ed.), *The families and genera of vascular plants* 3: 83-110. Springer-Verlag.
- MOLDENKE, H. N. (1963). Amaryllid genera and species. *Pl. Life (Stanford)* 19: 54-55.
- MORTON, C. V. (1959). Sur la nomenclature de deux Fougères rares d'Espagne. *Bull. Soc. Bot. France* 106: 231-234.
- MÜLLER-DOBLIES, D. & U. MÜLLER-DOBLIES (1978). Studies on tribal systematics of Amaryllidaceae 1. The systematic position of Lapiedra Lag. *Lagascalia* 8: 13-23.
- RAVENNA, P. (1974). Contributions to South American Amaryllidaceae VI. *Pl. Life (Stanford)* 30: 29-79.
- SALGUEIRO, F. J. (1988). *Estudios sobre los herbarios históricos de la Universidad de Sevilla*. PhD thesis. Universidad de Sevilla.
- STAFLEU, F. A. & R. S. COWAN (1979). Taxonomic Literature. Volume II: H-Le. *Regnum Veg.* 98.




Fig. 1. – Neotype of *Lapiedra martinezii* Lag. [lower specimen, with the arrow].
[Aedo 12305, MA] [© Real Jardín Botánico. Reproduced with permission]

