

**Zeitschrift:** Candollea : journal international de botanique systématique = international journal of systematic botany  
**Herausgeber:** Conservatoire et Jardin botaniques de la Ville de Genève  
**Band:** 64 (2009)  
**Heft:** 1

**Artikel:** Leucas mathewiana Sunojk. (Lamiaceae) : a new species from India  
**Autor:** Sunojkumar, Purayidathkandy  
**DOI:** <https://doi.org/10.5169/seals-879189>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 10.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Leucas mathewiana Sunojk. (Lamiaceae), a new species from India

Purayidathkandy Sunojkumar

## Abstract

SUNOJKUMAR, P. (2009). *Leucas mathewiana* Sunojk. (Lamiaceae), a new species from India. *Candollea* 64: 45-48. In English, English and French abstracts.

*Leucas mathewiana* Sunojk., new species from India, is described and illustrated. Its relationships with *Leucas aspera* (Willd.) Link, *Leucas lavandulifolia* Sm., and *Leucas zeylanica* (L.) R. Br. are discussed.

## Key-words

LAMIACEAE – *Leucas* – Andhra Pradesh – India – Taxonomy

## Résumé

SUNOJKUMAR, P. (2009). *Leucas mathewiana* Sunojk. (Lamiaceae), une nouvelle espèce d'Inde. *Candollea* 64: 45-48. En anglais, résumés anglais et français.

*Leucas mathewiana* Sunojk., une nouvelle espèce d'Inde, est décrite et illustrée. Ses affinités avec *Leucas aspera* (Willd.) Link, *Leucas lavandulifolia* Sm. et *Leucas zeylanica* (L.) R. Br. sont discutées.

---

Address of the author: Department of Botany, University of Calicut, Malappuram, 673 635, Kerala, India.

Email: [drsunoj@gmail.com](mailto:drsunoj@gmail.com)

Submitted on March 20, 2008. Accepted on February 11, 2009.

Edited by P. Perret

The genus *Leucas* R. Br. (*Lamiaceae*) comprises about 100 species and is found in Southern to tropical Africa, Arabia and Asia (SEBALD, 1980). In India, the genus is represented by about 40 species, of which 21 are Indian endemics (SINGH, 2001). Majority of these are distributed in the Southern Peninsula. Three additional endemic species from this area were described recently by SUNOJKUMAR & MATHEW (2002) and SUNOJKUMAR (2005, 2008). During collection of all the hitherto described Indian *Leucas* for a taxonomic study, the author discovered a small population of *Leucas* at an open wasteland, near a deciduous forest in the Kalasamudram forest area of Andhra Pradesh state, South India. Fresh specimens were collected and studied. This, when compared with specimens and types of related taxa, has confirmed it to be another distinct species, which is described and illustrated below.

***Leucas mathewiana* Sunojk., spec. nova (Fig. 1)**

**Typus:** INDIA. Andhra Pradesh: Ananthapur Dist., Kalasamudram open waste place, ca. 250 m, 24.X.2001 (fl.), Sunojkumar, P. CU49664 (holo-: K; iso-: CALI).

*Leucadi asperae similis, foliis angustis (3-5.5 cm longis x 4-6 mm latis), inflorescentia terminali non-globulari, floribus minus laxe dispositis circulariter patentibus, calyce semi-obliquo dentibus aequalibus, corolla magna (16-18 mm longa) differt.*

Erect annual herb, 20-40 cm high, stem obtusely quadrangular and grooved, not much branched, pubescent; internodes 10-17 cm long. Leaves opposite decussate; petiole 3-5 mm long; lamina 3-5.5 cm long, 4-6 mm broad, approximately 8-9 times longer than broad, membraneous, narrowly linear-lanceolate, base long cuneate to angustate, tip obtuse, margin almost entire or remotely 2-3 serrate, lateral veins not prominent, sparsely pubescent on both side with extrorse hairs. Inflorescence a terminal solitary verticillaster, subtended by a pair of leaves, fewer numbers of flowers, arranged loosely, spreading horizontally. Bracteoles 4-7 mm, 0.05 mm broad at base, narrowly linear, spinulose tip, slightly bend toward flowers, hispid outside and margins. Flowers 10-25, pedicel 1-2 mm long. Calyx tubular, side  $\pm$  10 mm long, up to 14mm long at

seeding stage, slightly bent at middle, 10 nerved, prominent towards upper half, each nerve excurrent in a teeth, inside glabrous, outside fully hispid, hairs 1-2 mm long, scattered, mouth semi-oblique, rim annulate with thick veins; teeth 10, equal sized, 1-1.5 mm long, tip subulate, spinulose. Corolla white, 16-17 mm long; tube 6-8 mm long, included in calyx tube, outside hairy above, inside middle annulate with a ring of fleshy hairs; lower lip  $\pm$  10 mm long, middle lobe  $\pm$  7 mm wide, fan shaped, side lobes 2.5 mm broad at free end, base narrowed, margin slightly revolute; upper lip 6 mm long, concave, bent, white spreading hairs outside, margins ciliate, tip of upper cilia slightly dark brown coloured. Stamens included in the upper lip, light reddish brown, upper pair shorter than the lower; filaments hairy; anther lobe 0.9 mm long, 0.5 mm broad, confluent theca. Disc semi-cup shaped, four lobes equal. Carpel locules 1 mm high, top rounded. Style 11 mm long, bent towards upper side; stigma unequally bilobbed, lower lobe 0.8 mm long, upper 0.06 mm long. Nutlets 3.5-4 mm long, 1.5 mm broad, almost cylindrical, dark brownish, glabrous shining, basal attachment flat, oblong outside, apex rounded.


**Distribution.** – The new species is endemic to Andhra Pradesh state (South India), from a small population of less than 250-300 plants are known from a single locality near the Kalasamudram forest area in the Ananthapur District.

**Ecology.** – Rare; the new species is found in the open, rocky waste place, near a deciduous forest area at an altitude between 200 and 250 m. The plant is found growing among grasses.

**Etymology.** – The species is named after Dr Philip Mathew, Professor of Botany, University of Calicut, Kerala, India in appreciation of his contribution to plant taxonomy.

**Flowering and fruiting.** – Observed as flowering in September to October; fruiting from November to December.

**Notes.** – *Leucas mathewiana* is included in *Leucas* sect. *Plagiostoma* Benth., together with its close relatives *L. aspera* (Willd.) Link, *L. lavandulifolia* Sm., and *L. zeylanica* (L.) R. Br. Taxa included in this section are very similar in their appearance and can be distinguished by their floral (particularly calyx) characters. The new species is distinct from its relatives by the characters shown in Table 1.



**Fig. 1.** – *Leucas mathewiana* Sunojk. **A.** Habit; **B.** Bracteole; **C.** Flower; **D.** Corolla; **E.** Corolla split open; **F.** Stamens; **G.** Pistil; **H.** Flowering calyx; **I.** Seeding calyx; **J.** Basal disc; **K.** Calyx split open; **L.** Nutlets.

[Sunojkumar CU49664, K] [Drawn by the author]

**Table 1.** – Comparison between *Leucas mathewiana* Sunojk. and its relatives (*L. aspera* (Willd.) Link, *L. lavandulifolia* Sm., *L. zeylanica* (L.) R. Br.).

	<i>Leucas aspera</i>	<i>Leucas lavandulifolia</i>	<i>Leucas zeylanica</i>	<i>Leucas mathewiana</i>
<b>Inflorescence</b>	2-3 towards tip	2-3 towards tip	2-3 towards tip	solitary
<b>Inflorescence outline</b>	globular, compact	semi-globular compact	semi-globular compact	flowers loosely spreading
<b>Flowers number</b>	many, up to 75	many, up to 50	many, up to 50	less, 15-25
<b>Pediceal [mm]</b>	± 0.5	± 0.5	± 0.5	1-2
<b>Bracteoles</b>	equal to calyx	equal to calyx	equal to calyx	half of calyx
<b>Calyx length [mm]</b>	9-10	7-8	5-6	± 10
<b>Calyx nature and distribution of hairs outside</b>	dense hispid on upper half	dense pubescent on upper half	dense hispid on upper half	sparsely hispid along entire length
<b>Calyx mouth</b>	very oblique	very oblique	semi oblique	semi oblique
<b>Calyx teeth</b>	10, anterior slightly longer	7-10, anterior much longer	8, anterior slightly longer	10, equal sized
<b>Corolla size [mm]</b>	13-15	13-15	13-15	16-17
<b>Nutlets [mm]</b>	± 2.5 x ± 1	± 2.5 x ± 1.3	± 2.8 x ± 3	± 4 x ± 1.5

## Acknowledgements

I am grateful to Dr Philip Mathew, Professor and Dr A. K. Pradeep, Curator of Department of Botany, University of Calicut, Kerala for various helps; to Dr Oskar Sebald, Staatliches Museum für Naturkunde, Stuttgart, Germany for suggestions; and to Dr J. F. Veldcamp, Nationaal Herbarium Nederland, Leiden University, Leiden, Netherlands, for the Latin diagnose. I thank all the Directors and Curators of different herbaria (B, BLAT, BM, BP, BR, C, CAL, CALI, CANB, CGE, E, FRI, G, K, KFRI, LE, LINN, LIV, M, MH, P, SKU, TBGT, W, WELT) for giving permission to study the herbarium specimens or providing specimens on loan. Hospitality provided by the authorities of different herbaria, particularly Dr Alan Paton and Dr Gemma Bramley of Royal Botanic Gardens, Kew, U. K. during my visit there and the «Research grant 2004» awarded by the International Association for Plant Taxonomy, Vienna are gratefully acknowledged.

## References

- SEBALD, O. (1980). Die Gattung *Leucas* R. Br. (Labiatae) in Afrika und auf der Arabischen Halbinsel. *Stuttgarter Beitr. Naturk.*, A 341.
- SINGH, V. (2001). *Monograph on Indian Leucas R. Br. (Dronapushpi) Lamiaceae*. Scientific Publishers.
- SUNOJKUMAR, P. (2005). *Leucas sebaliana* Sunojk. (Lamiaceae) a new species from India. *Candollea* 60: 233-236.
- SUNOJKUMAR, P. (2008). *Leucas sivadasaniana*, a new species of Lamiaceae (*Leucas* sect. *Astrodon*) from Peninsular India. *Novon* 18: 247-250.
- SUNOJKUMAR, P. & P. MATHEW (2002). *Leucas beddomei* (Hook. f.) Sunojkumar & P. Mathew (Lamiaceae) a new status and name for *Leucas hirta* var. *beddomei* Hook. f. - a little known endemic from India. *Rheedea* 12: 169-174.