

| | |
|---------------------|---|
| Zeitschrift: | Candollea : journal international de botanique systématique = international journal of systematic botany |
| Herausgeber: | Conservatoire et Jardin botaniques de la Ville de Genève |
| Band: | 64 (2009) |
| Heft: | 1 |
| Artikel: | Begonia bullatifolia L. Kollmann and Begonia leopoldinensis L. Kollmann (Begoniaceae) : two new species from the Atlantic Forest in the State of Espírito Santo, Brazil |
| Autor: | Kollmann, Ludovic Jean Charles |
| DOI: | https://doi.org/10.5169/seals-879196 |

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Begonia bullatifolia L. Kollmann and Begonia leopoldinensis L. Kollmann (Begoniaceae), two new species from the Atlantic Forest in the State of Espírito Santo, Brazil

Ludovic Jean Charles Kollmann

Abstract

KOLLMANN, L. J. C. (2009). *Begonia bullatifolia* L. Kollmann and *Begonia leopoldinensis* L. Kollmann (Begoniaceae), two new species from the Atlantic Forest in the State of Espírito Santo, Brazil. *Candollea* 64: 117-122. In English, English and French abstracts.

This paper presents a contribution to the knowledge of the Brazilian Begoniaceae. Two new species, *Begonia bullatifolia* L. Kollmann and *Begonia leopoldinensis* L. Kollmann, are described and illustrated. Both are related to *Begonia santos-limae* Brade.

Key-words

BEGONIACEAE – Begonia – Brazil – State of Espírito Santo – Atlantic Forest – Taxonomy

Résumé

KOLLMANN, L. J. C. (2009). *Begonia bullatifolia* L. Kollmann et *Begonia leopoldinensis* L. Kollmann (Begoniaceae), deux nouvelles espèces de la Forêt Atlantique de l'Etat d'Espírito Santo, Brésil. *Candollea* 64: 117-122. En anglais, résumés anglais et français.

Ce travail présente une contribution à la connaissance des Begoniaceae brésiliennes. Deux nouvelles espèces, *Begonia bullatifolia* L. Kollmann et *Begonia leopoldinensis* L. Kollmann, sont décrites et illustrées. Les deux espèces sont proches de *Begonia santos-limae* Brade.

Address of the author: Museu de Biologia Prof. Mello Leitão, Av. José Ruschi 4, CEP 29650-000 Santa Teresa, ES, Brazil.

Email: Ludovic@limainfo.com.br

Submitted on July 3, 2008. Accepted on January 23, 2009.

Edited by P. Bungener

ISSN: 0373-2967

Candollea 64(1): 117-122 (2009)

© CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 2009

The genus *Begonia* L. is pantropical and comprises about 1400 species divided among 63 sections. More than 240 species, distributed in 13 sections, occur in Brazil (SMITH & al., 1986; DOORENBOS & al., 1998), mostly native to the Atlantic Forest (DUARTE, 1961). The Atlantic Forest is one of the most important World Hotspots (MITTERMEIER & al., 1999; SIMÕES & al., 2002) and estimated that is home to approximately 8000 endemic species of plants (MYERS & al., 2000).

Recent field works in the state of Espírito Santo contributed to the discovery of new species of *Begonia* (KOLLMANN, 2003, 2006, 2007, 2008; JACQUES & MAMEDE, 2004), illustrating the high biodiversity and the number of endemic species of this state.

Following fieldwork in the mountainous region of central Espírito Santo State, two species of *Begonia* were collected and described here as new species. These two species are part of sect. *Knesebeckia* (Klotzsch) A. DC. characterized by their rhizomatous habit, persistent stipules, stellate trichomes, a connective not extending beyond the anther, bifurcate placental branches, and ovules present on both faces of the placental branches.

1. *Begonia bullatifolia* L. Kollmann, spec. nova (Fig. 1)

Typus: BRAZIL. Edo Espírito Santo: Domingos Martins, R. Kautsky forest, 15.X.2006, Kollmann, L. 11058 (fl. fr.) [flower in cultivation 15.VII.2007] (holo-: MBML; iso-: RB).

Species haec Begoniae santoslimae Brade affinis, sed lamina bullatis nec peltatus, bracteis persistentibus et floribus pistillatis 6-tepalis differt.

Rhizomatous herb 10-15 cm tall (excluding inflorescence), sub-erect, hairy, with glandular trichomes, stellate trichomes erect and flattened stellate trichomes on abaxial face. Stems 0.8-1.2 cm diam., green-brown, with lenticels, internodes 0.4-0.7 × 0.8-1 cm, stellate trichomes red. Stipules persistent, 1.35-2 × 0.8-1.2 cm, ovate to oblong, papyraceous when dry, translucent, greenish, base red, nerves reddish, carinate near the apex, mucronate, adaxial face glabrate, abaxial face with stellate trichomes. Petioles 5.5-12.5 × 0.3-0.6 cm, cylindrical, with lenticels, stellate trichomes red. Leaf blades 7.5-10 × 9-13 cm, green, asymmetric, reniform to orbiculate, base cordate, adaxial face bullate, white when young due to the trichomes, adaxial face with white flattened stellate trichomes, abaxial face with red erect stellate trichomes; margins entire to repand, with hydathode; venation actinodromous, 7-9-veined, pilose, stomates single. Inflorescence 21-35 cm long, red-brown, thyrsiform, with lenticels, first internode 18.5-24 cm, second 3.5-4.5 cm, third 2 cm. Bracts of first node leaf like, 1.9-2.5 × 2.1-3.1 cm, all others 0.8-1.5 × 1-1.5 cm,

persistent, green, veins pinkish, papyraceous and brown when dry, obovate to orbiculate, apex retuse and lightly mucronate, both faces with glandular trichomes when young, abaxial face with stellate trichomes. *Staminate flowers* peduncles 0.8-1.4 cm long, white-pinkish, glandular trichomes; sepals 2 (outer tepals), 1.6-2.2 × 0.9-1.2 cm, white-pinkish, ovate, apex obtuse, adaxial face with glandular trichome; petals 2 (inner tepals), 1.2-1.5(-1.9) × 0.57-0.85 cm, white, obovate to spatulate, base cuneate, apex obtuse; stamens 26-37, yellow, filaments unequal, fused below into a column, glandular trichomes, anthers, obovate, rimose, connective shorter than the anthers. *Pistillate flowers* pedicels 1.5-2.2 cm long, pinkish; sepals 3, 1.3-2.2 × 0.6-1.2 cm, pinkish-white, ovate to obovate, apex obtuse, abaxial face with glandular trichomes, petals 3, 0.9-1.55 × 0.55-0.8 cm, white, obovate, apex obtuse; styles 3, greenish-yellow, united at base, bifurcate, spirally twisted, with bands of stigmatic papillae on the branches; ovary 3-locular, placentation axile, one placenta bifurcate per locule, ovules on both sides of placentae. *Capsules* 0.9-1.2 × 0.9 cm, glandular trichomes, basally dehiscent, wings 3, larger one 1-1.8 × 1.1-1.7 cm, smaller 0.7-1.6 × 0.6-1 mm, glandular trichomes. *Seeds* ca. 0.45 × 0.25 mm, cylindrical, oblong.

Paratypus. – BRAZIL. Edo Espírito Santo: Domingos Martins, R. Kautsky forest, 16.I.1995, Cintia Chamas 59 (fl.) (MBML).

Etymology. – The epithet refers to the bullate leaf blades.

Distribution and ecology. – *Begonia bullatifolia* grows in leaf litter and humus of the central mountains of the Atlantic Forest of Espírito Santo state, Brazil, at 750 m, in the Roberto Kautsky forest, Domingos Martins municipality. It is possible that this species came from another locality and was introduced to the Kautsky forest. Indeed, Roberto Kautsky often transplant species from the state of Espírito Santo.

Due to the apparent endemic distribution of *B. bullatifolia*, with an extent of occurrence estimated to be less than 10 km², it would seem prudent to include this species on the list of Critically Endangered species (CR) [B2ab(iii)] according to the IUCN (2001) criteria.

Relationships. – *Begonia bullatifolia* resembles *B. santos-limae* Brade, due to its rhizomatous stem, stipules persistent, stellate trichomes, thyrsiform inflorescence. The new species can be separated from it by the non peltate (vs. peltate) leaves with a bullate adaxial face (vs. flat), persistent inflorescence bracts (vs. caducous), and pistilate flower perianth of 6 tepals (vs. 5).




Fig. 1. – *Begonia bullatifolia* L. Kollmann. **A.** Habit; **B.** Stipule; **C.** Bracts; **D.** Sepal male flower; **E.** Petal male flower; **F.** Stamen; **G.** Sepals and petals female flower; **H.** Stigma, dorsal face (left), ventral face (right); **I.** Fruit; **J.** Seed.

[Kollmann 11058, MBML] [Drawn by the author]

2. *Begonia leopoldinensis* L. Kollmann, spec. nova (Fig. 2)

Typus: BRASIL. Edo Espírito Santo: Santa Leopoldina, Colina Verde (Morro do Agudo), Prop. Israel Elias Ramos, 300-500 m, 20°05'50"S 40°26'16"W, 16.V.2006 (fr.), *Magnago*, L. F. S. & al. 971 (MBML); 12.IX.2006 (fr.), *Demuner*, V. & al. 2821 (MBML); 13.III.2007 (fl.), *Demuner*, V. & al. 3142 (MBML); 14.III.2007 (fl. fr.), *Demuner*, V. & al. 3174 (MBML); 30.V.2007 (fl.), *Demuner*, V. & al. 4111 (MBML).

Species nova Begonia santoslimae Brade affinis, sed habitu minori, laminis late ovatis vel rotundatus-apicisque differt.

Rhizomatous herb, 15-34 cm tall (excluding inflorescence), prostrate to erect, hairy, glandulate trichomes and stellate trichomes erect. Stems 1-2 cm diam., brown, internodes 4.5-7.5 cm long. *Stipules* 1.7-2 × 1.4-1.8 cm, persistent, papyraceous when dry, ovate, apex obtuse to retuse, mucronate, adaxial face glabrate, abaxial face lightly crenate, pilose. *Petioles* 8-19 × 5.5-6 mm, red, cylindrical, pilose, stellate trichomes purple when young. *Leaf blades* 7.5-17.5 × 9-15 cm, peltate, asymmetric, widely ovate to orbiculate, base and apex rounded, adaxial face green, abaxial face green to purple, trichomes purplish, young leaf purplish due to the trichomes, margins entire with hydathodes; venation actinodromous, veins 8. *Inflorescence* reddish, thyrsiform, 40-54 cm long, with 3-10 nodes, stellate trichomes. *Bracts* 1.2-2 × 0.9-1.6 cm, of first node leaf-like, all others, deciduous, white with pinkish apex, ovate, apex carinate abaxially, mucronate, margins with glandulate trichomes. *Bracteoles* 0.4-1 × 0.4-0.5 cm, white with pinkish apex, deciduous, ovate to orbiculate, apex acute, mucronate, margins with glandulate trichome. *Staminate flowers* peduncles 1.1-1.5 cm long, glandulate trichomes; sepals 2 (outer tepals), 1.25-1.7 × 1.2-1.5 cm, pinkish-white, widely ovate, apex obtuse; petals 2 (inner tepals), pinkish-white, 1-1.15 × 0.6-0.75 cm, obovate, base cuneate, apex obtuse, stamens 21-27, yellow, filament 0.5-1 mm long, unequal, fused below into a column, anthers obovate, rimose, connective shorter than the anthers. *Pistilate flowers* pedicels ca. 1.7 cm long, pinkish, pilose, glandulate trichomes; tepals 5(-6), pinkish-white, 1-1.5 × 0.5-1.2 cm, unequal, ovate to obovate, apex obtuse, margins with glandulate trichome, styles 3, yellow, united at base, bifurcate, spirally twisted, with bands of filamentose stigmatic papillae on the branches; ovary 3-locular, placentation axile, one placenta per locule bifurcate, ovules on both sides of placentae. *Capsules* 0.8-1.2 × 0.75-1.1 cm, basally dehiscent, glandulate trichomes; wings 3, unequal, papyraceous, glandulate trichomes when young, larger 1.7-2.3 × 1-2 cm, lightly ascending, smaller, 1.5-1.9 × 0.5-0.9 cm. *Seeds* ca. 0.5 mm long, oblong, cylindrical.

Paratypi. – BRASIL. Edo Espírito Santo: Santa Leopoldina, Colina Verde (Morro do Agudo), Prop. Israel Elias Ramos, 300-500 m, 20°05'50"S 40°26'16"W, 16.V.2006 (fr.), *Magnago*, L. F. S. & al. 971 (MBML); 12.IX.2006 (fr.), *Demuner*, V. & al. 2821 (MBML); 13.III.2007 (fl.), *Demuner*, V. & al. 3142 (MBML); 14.III.2007 (fl. fr.), *Demuner*, V. & al. 3174 (MBML); 30.V.2007 (fl.), *Demuner*, V. & al. 4111 (MBML).

Etymology. – The specific epithet alludes to the type municipality, Santa Leopoldina, where the new species was found.

Distribution and ecology. – *Begonia leopoldinensis* was found growing in shaded places, in rock cracks, of rocky outcrops, in the dry Atlantic Forest at 300-500 m above sea level. In nature, this species is prostrate on rock but in cultivation the leaves are erect.

Due to the apparent endemic distribution of *B. leopoldinensis*, with an extent of occurrence estimated to be less than 10 km², it would appear prudent to include this species on the Critically Endangered species list (CR) [B2ab(ii)] according to the IUCN (2001) criteria.

Relationships. – *Begonia leopoldinensis* is similar to *B. santos-limae* Brade, in its rhizomatous stem, stellate trichomes, peltate leaf, and thyrsiform inflorescence. It can be distinguished from it by its smaller size, ca. 20 cm tall (vs. ca. 50 cm), petioles, 8-15 cm long (vs. 20-40 cm), large ovate to orbiculate leaf (vs. ovate), with a rounded apex (vs. obtuse).

Acknowledgments

The author wishes to acknowledge the staff of the Mello Leitão Biological Museum, especially Helio de Queiroz Boudet Fernandes, Director of the Museum and Curator of MBML Herbarium, for their cooperation, Roberto Kautsky and the reviewers for the comments.




Fig. 2. – *Begonia leopoldinensis* L. Kollmann. **A.** Habit; **B.** Stipule; **C.** Bracts; **D.** Bracteole; **E.** Sepal male flower; **F.** Petal male flower; **G.** Stamen; **H.** Tepals female flower; **I.** Stigma, ventral face (left), dorsal face (right); **J.** Fruit; **K.** Seed; **L.** Stellate trichome.

[Demuner, V. & al. 2511, MBML] [Drawn by the author]

References

- DOORENBOS, J., S. M. SOSEF & J. J. F. E. de WILDE (1998). The sections of Begonia. Studies in Begoniaceae VI. *Wageningen Agric. Univ. Pap.* 98(2).
- DUARTE, A. P. (1961). Considerações acerca do comportamento e dispersão de algumas espécies de Begônias do Estado da Guanabara. *Arch. Jard. Bot. Rio de Janeiro* 17: 57-105.
- IUCN (2001). *IUCN Red List of threatened species*. IUCN Species Survival Commission.
- JACQUES, E. L. & M. C. H. MAMEDE (2004). Novelties in Begonia (Begoniaceae) from the coastal forests of Brazil. *Brittonia* 56: 75-81.
- KOLLMANN, L. J. C. (2003). Begonia ruschii L. Kollmann (Begoniaceae), uma nova espécie da Floresta Atlântica do Espírito Santo, Brasil. *Bol. Mus. Biol. Prof. Mello-Leitão* 15: 29-33.
- KOLLMANN, L. J. C. (2006). Begonia novalombardiensis L. Kollmann (Begoniaceae), une nouvelle espèce de la forêt atlantique de l'Etat de l'Espirito Santo, Brésil. *Candollea* 61: 89-92.
- KOLLMANN, L. J. C. (2007). Begonia callosa L. Kollmann (Begoniaceae), a new species of Espírito Santo, Brazil. *Candollea* 62: 141-144.
- KOLLMANN, L. J. C. (2008). Duas novas espécies de Begonia (Begoniaceae), do Espírito Santo, Brasil. *Rodriguesia* 59: 259-263.
- MITTERMEIER, R. A., N. MYERS, & C. G. MITTERMEIER (1999). *Hotspots Earth's biologically richest and most endangered terrestrial ecoregions*. Conservation International.
- MYERS, N. R., A. MITTERMEIER, C. G. MITTERMEIER, G. A. B. FONSECA & J. KENT (2000). Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858.
- SIMÕES, L. L. & C. F. LINO (2002). *Sustentável Mata Atlântica: a exploração de seus recursos florestais*. Editora Senac.
- SMITH, L. B., D. C. WASSHAUSEN, J. GOLDING, & C. E. KAREGEANNES (1986). Begoniaceae. Part I: Illustrated key. Part II: Annotated Species List. *Smithsonian Contr. Bot.* 60.