

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany

Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève

Band: 64 (2009)

Heft: 2

Artikel: Pandanus nusbaumeri Callm. & L. Gaut. (Pandanaeae) : a new species from northern Madagascar

Autor: Callmander, Martin W. / Gautier, Laurent / Trigui, Sonia M.

DOI: <https://doi.org/10.5169/seals-879206>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Pandanus nusbaumeri Callm. & L. Gaut. (Pandanaaceae), a new species from northern Madagascar

Martin W. Callmander, Laurent Gautier & Sonia M. Trigui

Abstract

CALLMANDER, M. W., L. GAUTIER & S. M. TRIGUI (2009). *Pandanus nusbaumeri* Callm. & L. Gaut. (Pandanaaceae), a new species from northern Madagascar. *Candollea* 64: 213-218. In English, English and French abstracts.

Pandanus nusbaumeri Callm. & L. Gaut. (Pandanaaceae), a new species from the Montagne d'Ambre (northern Madagascar), is described and illustrated. It morphologically resembles three other species from Madagascar: *Pandanus insuetus* Huynh, *Pandanus perrieri* Martelli and *Pandanus sermolliana* Callm. & Buerki. It can, however, be easily distinguished from these three taxa on the basis of several morphological characters, including the dimensions of the syncarp and peduncle, the incompletely united drupes with the pileus divided by deep apical sinuses, forming acute pyramids equal in number to the carpels, each with a thick, slightly spinescent, sub-vertical, prominent stigma. This species appears to be endemic to the poorly-explored leeward, western slopes of the massif, which have tropophyllous semi-deciduous forests whereas the moist evergreen forest is found in most other parts of the Montagne d'Ambre.

Key-words

PANDANACEAE – *Pandanus* – Madagascar – Montagne d'Ambre – Taxonomy – IUCN Red List

Résumé

CALLMANDER, M. W., L. GAUTIER & S. M. TRIGUI (2009). *Pandanus nusbaumeri* Callm. & L. Gaut. (Pandanaaceae), une nouvelle espèce du Nord de Madagascar. *Candollea* 64: 213-218. En anglais, résumés anglais et français.

Pandanus nusbaumeri Callm. & L. Gaut. (Pandanaaceae), une nouvelle espèce de la Montagne d'Ambre (N Madagascar), est décrite et illustrée. Elle est proche morphologiquement de trois autres espèces à Madagascar: *Pandanus insuetus* Huynh, *Pandanus perrieri* Martelli and *Pandanus sermolliana* Callm. & Buerki. Elle peut en être cependant facilement différenciée morphologiquement par plusieurs caractères, comme la dimension du syncarpe et du pédoncule, ainsi que par les drupes incomplètement soudées possédant un pileus divisé par des sinus profonds formant des pyramides aigües en nombre égal à celui des carpelles, chacune étant surmontée d'un stigmate presque spinescent, épais, sub-vertical, proéminent. Cette espèce est probablement endémique du versant ouest du massif, moins visité des botanistes, peuplé de forêts tropophylles semi-décidues alors que la majeure partie de la Montagne d'Ambre est recouverte de forêt dense humide de montagne.

Addresses of the authors: MWC: Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri, 63166-0299, U.S.A. & Conservatoire et Jardin botaniques de la Ville de Genève, ch. de l'Impératrice 1, CP 60, 1292 Chambésy, Switzerland. Email: martin.callmander@mobot-mg.org

LG, SMT: Conservatoire et Jardin botaniques de la Ville de Genève, ch. de l'Impératrice 1, CP 60, 1292 Chambésy, Switzerland.

Submitted on May 4, 2009. Accepted on July 13, 2009.

Edited by P. Bungener

Introduction

Madagascar is one of the major centres of diversity of the genus *Pandanus* Parkinson (*Pandanaceae*) with ca. 80 species (CALLMANDER & al., 2007) of the ca. 600 to 700 known in the Paleotropics (STONE, 1974). In the process of preparing the volume on *Pandanaceae* for the “Flore de Madagascar et des Comores”, and as part of a comprehensive revision of all Malagasy *Pandanus*, a series of taxonomic treatments and notes have been published on the family (CALLMANDER, 2001; CALLMANDER & LAIVAO, 2002; CALLMANDER & al., 2001, 2003a, 2003b, 2008; LAIVAO & al., 2000, 2006, 2007). In this contribution, we describe a distinctive new species from Montagne d’Ambre in the far northern part of the island.

The Montagne d’Ambre is a vast volcanic massif with an oval form, covering 2840 km² (SAINT OURS, 1960) and culminating at 1475 m (Fig. 1). The first volcanic activity dates back to the Late Cretaceous (ca. 90 mya) with a series of eruptions in the Late Tertiary and the Quaternary (2 to 6 mya) (SAINT OURS, 1960), leading to the typical geomorphological shape of the massif we know today (BRENON, 1972). The Montagne d’Ambre is mainly covered with montane evergreen humid forest (at elevations > 1000 m) that is now geographically isolated from other areas with similar vegetation but might once have been linked to the more southern evergreen humid forest during the Quaternary (NUSBAUMER & al., in press). The recent discovery in the Daraina region (NE Madagascar) of several forest species that were until now considered as endemic to the Montagne d’Ambre supports this hypothesis (NUSBAUMER & al., in press). On the western, leeward side of the massif, a transition to trophophyllous forests is observed. These had not received much attention until the recent intensive botanical inventory led by a team from the Conservatoire et Jardin botaniques de la Ville de Genève (Switzerland) (CJB) in collaboration with the Département de Biologie et Ecologie Végétales of the Université d’Antananarivo (Madagascar) (DBEV).

Pandanus nusbaumeri Callm. & L. Gaut., *spec. nova* (Fig. 2)

Typus: MADAGASCAR. **Prov. Antsiranana:** Montagne d’Ambre, versants ouest, 12°35’00”S 49°08’22”E, 1170 m, 13.XI. 2007, fr., Gautier, L. & al. 5217 (holo-: G!; iso-: MO!, P!, TAN!).

Haec species P. insuetus Huynh, *P. perrieri* Martelli et *P. sermolliana* Callm. & Buerki *similis*, sed ab eis syncarpia pedunculique dimensionibus atque drupis incomplete connatis, pileo sinibus apicalibus profundis fisso, pyramides acutas carpellorum numerum aequantes quamque in stigma crassum leniter spinescens subverticalia prominens desinentem formantibus facile distinguitur.

Tree to 8 m tall, stem prickly, 12 cm diam. *Leaves* gradually attenuate in the apical part, 100-110 cm long, 4.5-5 cm wide in the median part, 5.5-6 cm wide near the sheath; dry leaves coriaceous; auricles lacking, leaf blade densely alveolate on the abaxial face, longitudinal and transverse veins visible on both surfaces; prickles brownish; marginal prickles beginning at 12-13 cm from the base and extending to the apex, antrorse, 2-3 mm in the lower third, 2-4(-6) mm apart, strong, to 2 mm in the mid third, (5-)-8-10(-14) mm apart, to 1 mm in the distal third, 2-3 mm apart; midrib armed, prickles small (< 1 mm), randomly disposed (2-20 mm apart) and slightly prominent in the basal half, then regularly disposed, spaced (2-4 mm apart) and prominent in the apical half; sheath 11-12 cm long, 6 cm wide at apex, 8-9 cm at base. *Infructescence* terminal, the solitary syncarp erect on a straight peduncle; syncarp 11-12 × 14-15 cm, sub-spherical; peduncle 17-18 cm long, 3-3.5 cm wide at apex, 1.8-2 cm in the middle, straight, trigonous, veins visible, first bract borne 5 cm from the base of syncarp, 8 to 9 bracts on entire peduncle. *Drupes* 12-15, 35-45 mm high, (3.5-)-5-6(-9) cm wide, 3.5-5.5 mm thick, (3-)-4(-6) angled; carpels (3-)-5-8(-14), incompletely united, free in the apical third, tapering to the base, pileus divided by apical sinuses 7-10 mm deep into acute pyramids as many as there are carpels; stigmas (3-)-5-8(-14) per drupe, 3-5 mm high, thick, slightly spinescent, sub-vertical, prolonged by a sinus 0.4-0.8 mm long towards the centre of the carpel, prominent at its centre; endocarp 15-25 mm long in the centre, shortened on both sides, 30-60 mm wide, 15-20 mm away from the stigmas; seed locule oblong, 10-12 × 6-7 mm, superior and inferior mesocarp thick and fibrous. *Male flowers* unknown.

Etymology. – The species is named in honour of Louis Nusbaumer, who was a member of the team that collected the type and has always demonstrated great enthusiasm for documenting difficult groups, such as *Pandanaceae*, thereby making an important contribution to a better understanding of diversity and distribution of the family in Madagascar, especially in the complex northern part of the island.

Distribution and ecology. – *Pandanus nusbaumeri* is only known from a single collection on the western slopes of the Montagne d’Ambre in trophophyllous semi-deciduous forest (Fig. 1).

Conservation status. – With only 1 collection known, an AOO of 9 km² (calculated following CALLMANDER & al., 2007) and only 1 subpopulation situated within the protected area of Montagne d’Ambre, *P. nusbaumeri* is assigned a preliminary status of “Vulnerable” (VU D2) (IUCN, 2008).

Notes. – *Pandanus nusbaumeri* is morphologically similar to three other members of the genus occurring in Madagascar: *P. insuetus* Huynh, *P. sermolliana* Callm. & Buerki and *P. perrieri* Martelli (Table 1). These four species all have in common a large infructescence and incompletely united drupes with each




Fig. 1. – Map of Northern Madagascar. **A.** Current Protected Areas in Northern Madagascar (hatched) and frame (enlarged in B); **B.** Site at which the type collection was made in the Montagne d'Ambre area, with remaining primary forest in grey.




Fig. 2. – *Pandanus nusbaumeri* Callm. & L. Gaut. **A.** Syncarp; **B.** Longitudinal section of a pluriloculate drupe; **C.** Leaf.
[A-C: Gautier, L. & al. 5217, G] [Drawing: C. Chatelain]

Table 1. – Principal morphological characters differentiating the new species, *Pandanus nusbaumeri* Callm. & L. Gaut., from *P. insuetus* Huynh, *P. perrieri* Martelli and *P. sermolliana* Callm. & Buerki.

	Leaf length [cm]	Syncarp dimension [cm]	Peduncle length [cm]	Apex of the drupe	Stigma (shape and place)
<i>Pandanus nusbaumeri</i> Callm. & L. Gaut.	100-110	14-15 × 11-12	17-18	apex divided into acute pyramids	slightly spinescent, sub-vertical, prominent at the center of each acute pyramids
<i>Pandanus insuetus</i> Huynh	170(-250)	18-20 in diam.	ca. 40	centre part of the apex with dense short ridges in longitudinal rows	deltoid, sub-vertical, gathered in a circle at the apex of the drupe 6-10 mm apart from another
<i>Pandanus perrieri</i> Martelli	175-250	18-20 in diam.	20-25	incompletely united with flat carpels apex	flat to deltoid, sub-horizontal or rarely sub-vertical and not raised
<i>Pandanus sermolliana</i> Callm. & Buerki	210-240	15-16 × 13-14	17-21	incompletely united with dome-like carpels apex	sub-vertical or rarely sub-horizontal, only slightly spinescent, and raised on an incompletely merged base

carpel bearing a sub-vertical stigma. The new species can, however, be easily distinguished from the three others by several significant morphological features, as summarized in Table 1. The new species differs from *P. insuetus* and *P. perrieri* by the size of the syncarp (14-15 × 11-12 cm) and peduncle (17-18 cm long) (vs. ca. 18-20 cm in diam. and peduncle ca. 40 cm long in *P. insuetus*; ca. 18-20 cm in diam. and peduncle 20-25 cm long in *P. perrieri*), drupes incompletely united, free in the upper third, pileus divided by apical sinuses 7-10 mm deep into acute pyramids equal in number to the carpels (Fig. 3A, B) (vs. drupes united except for the centre part of the apex that has dense short ridges in longitudinal rows (LAIVAO & al., 2006: 269, Fig. 1B) in *P. insuetus*; incompletely united but with a flat apex in *P. perrieri*), stigmas slightly spinescent, sub-vertical, prominent at the centre of each acute pyramids (vs. gathered in a circle at the apex of the drupe 6-10 mm apart one another in *P. insuetus*; sub-horizontal or rarely sub-vertical, flat or not raised in *P. perrieri*) (HUYNH, 2000; LAIVAO & al., 2006). *Pandanus nusbaumeri* closely resembles *P. sermolliana*, endemic to the Galoka-Kalabenono massifs (CALLMANDER & al., 2008) ca. 130 km south of Montagne d'Ambre. The new species shares the sub-spherical shape and size of syncarps (ca. 15-16 × 13-14 cm in *P. sermolliana*) and leaves that lack large auricles. However, *P. nusbaumeri* can be distinguished from *P. sermolliana* by its smaller leaves (100-110 cm long with a 11-12 cm sheath vs. leaves 210-240 cm with a 16-17 cm sheath in *P. sermolliana*),

drupes with an apex divided into acute pyramids (vs. drupes with dome-like pileus in *P. sermolliana*), stigmas (3-)5-8(-14) per drupe, prominent at the apex of an acute pyramid (vs. (1-)5(-7) per drupe, laterally disposed on the margin of a slightly concave plateau in *P. sermolliana*).

Acknowledgements

The authors thank the staff of the Parc Botanique et Zoologique de Tsimbazaza, Madagascar National Parks, formerly ANGAP (Association Nationale pour la gestion des Aires Protégées), and the Missouri Botanical Garden's (MBG) offices in Antananarivo and Antsirana, especially Lalao Andriamahefarivo and Jeremi Razafitsalama. We are grateful to Roy Gereau for preparing the Latin diagnosis, Pete Lowry for improving an earlier version of this manuscript and Cyrille Chatelain for the fine illustration. Financial support of MWC's participation was provided by grants from the U.S. National Science Foundation (0743355) and the Andrew W. Mellon Foundation. The Initiative Sud Experts Plantes, the Fonds Marc Birkigt and the Département de Botanique et Biologie végétale of the University of Geneva funded the inventory work, which was conducted under a collaboration between the CJB and the DBEV. We thank Prof. Charlotte Rajeriarison, Dr Roger Edmond, Patrick Ranirison, Solotiana D. Ramandimbiana and Mialy H. Razanajatovo from the DBEV.



Fig. 3. – *Pandanus nusbaumeri* Callm. & L. Gaut. **A.** Infructescence; **B.** Detail of the stigmas.
[A-B: Gautier, L. & al. 5217, G] [Photos: L. Gautier]

References

- BRENON, P. (1972). The Geology of Madagascar. In: BATTISTINI, R. & G. RICHARD-VINCENT (ed.), *Biogeography and Ecology in Madagascar*: 27-86. W. Junk. B. V. Publishers.
- CALLMANDER, M. W. (2001). *Pandanus* subg. *Martellidendron* (Pandanaceae) part II: revision of sect. *Martellidendron* Pic. Serm. in Madagascar. *Bot. J. Linn. Soc.* 137: 353-374.
- CALLMANDER, M. W. & M. O. LAIVAO (2002). Révision de *Pandanus* sect. *Dauphinensia* H. St. John (Pandanaceae) à Madagascar. *Bot. Helv.* 112: 47-67.
- CALLMANDER, M. W., S. WOHLHAUSER & M. O. LAIVAO (2001). Une nouvelle section du genre *Pandanus* (Pandanaceae) à Madagascar: *Pandanus* sect. *Tridentistigma*. *Adansonia* 23: 49-57.
- CALLMANDER, M. W., M. O. LAIVAO & S. WOHLHAUSER (2003a). Les *Pandanus* sect. *Acanthostyla* Martelli (Pandanaceae) d'altitude du Nord de Madagascar, avec description de deux nouvelles espèces. *Candollea* 58: 63-74.
- CALLMANDER, M. W., S. WOHLHAUSER & L. GAUTIER (2003b). Notes biogéographiques sur les Pandanaceae des forêts humides du Nord de Madagascar. *Candollea* 58: 351-367.
- CALLMANDER, M. W., G. E. SCHATZ, P. P. LOWRY II, M. O. LAIVAO, J. RAHARIMAMPIONONA, S. ANDRIAMBOLOLONERA, T. RAMINOSOA & T. CONSIGLIO (2007). Application of IUCN Red List criteria and assessment of Priority Areas for Plant Conservation in Madagascar: rare and threatened Pandanaceae indicate new sites in need of protection. *Oryx* 41: 168-176.
- CALLMANDER, M. W., S. BUERKI & S. WOHLHAUSER (2008). A new threatened species of Pandanaceae from northwestern Madagascar: *Pandanus sermolliana*. *Novon* 18: 421-424.
- HUYNH, K.-L. (2000). The genus *Pandanus* (Pandanaceae) in Madagascar (part 6). *Bot. Jahrb. Syst.* 122: 201-224.
- IUCN (2008). Categories and Criteria (version 3.1). IUCN, Gland, Switzerland.
- LAIVAO, M. O., M. W. CALLMANDER & S. WOHLHAUSER (2000). Une espèce nouvelle de *Pandanus* sect. *Martellidendron* (Pandanaceae) de la péninsule de Masoala, Madagascar. *Bot. Helv.* 110: 41-49.
- LAIVAO, M. O., M. W. CALLMANDER & S. BUERKI (2006). Sur les *Pandanus* (Pandanaceae) à stigmates saillants de la Côte Est de Madagascar. *Adansonia* 28: 267-285.
- LAIVAO, M. O., M. W. CALLMANDER & S. BUERKI (2007). Révision de *Pandanus* sect. *Foullioya* à Madagascar. *Adansonia* 29: 39-57.
- NUSBAUMER, N., P. RANIRISON, L. GAUTIER, C. CHATELAIN, P.-A. LOIZEAU & R. SPICHTER (in press). Loky-Manambato: point de rencontre des principales unités phytogéographiques de Madagascar. In: VAN DER BURGT, X., J. VAN DER MAESEN & J.-M. ONANA (ed.), *Proceedings of the XVIIIth AETFAT Congress, Yaoundé, Cameroun*. Royal Botanic Gardens, Kew
- SAINT OURS, J. DE (1960). *Etudes Géologiques dans l'extrême Nord de Madagascar et l'Archipel des Comores*. République Malgache. Ministère de L'Economie Nationale, Tananarive, Madagascar.
- STONE, B. C. (1974). Towards an improved infrageneric classification in *Pandanus* (Pandanaceae). *Bot. Jahrb. Syst.* 94: 459-540.