**Zeitschrift:** Contributions to Natural History: Scientific Papers from the Natural

History Museum Bern

**Herausgeber:** Naturhistorisches Museum Bern

**Band:** - (2008)

**Artikel:** African and Southeast Asian elements in the spider fauna of the

Western Ghats of India

Autor: Sudhikumar, Ambalaparambil Vasu / Maelfait, Jean-Pierre / Lens, Luc

**DOI:** https://doi.org/10.5169/seals-787070

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

European Arachnology 2008 (W. Nentwig, M. Entling & C. Kropf eds.), pp. 165–175. © Natural History Museum, Bern, 2010. ISSN 1660-9972 (Proceedings of the 24<sup>th</sup> European Congress of Arachnology, Bern, 25–29 August 2008).

# African and Southeast Asian elements in the spider fauna of the Western Ghats of India

AMBALAPARAMBIL VASU SUDHIKUMAR<sup>1</sup>, JEAN-PIERRE MAELFAIT †, LUC LENS<sup>2</sup>, FREDERIK HENDRICKX<sup>2</sup> & POTHALIL ANTONY SEBASTIAN<sup>3</sup>

- <sup>1</sup> Department of Zoology, Christ College, Irinjalakuda, Kerala, India. PIN: 680 125 (corresponding author: avsudhi@rediffmail.com)
- <sup>2</sup> Terrestrial Ecology Unit, Department of Biology, Ghent University, K.L. Ledeganckstraat 35 B–9000 Ghent, Belgium
- ³ Division of Arachnology, Dept. of Zoology, Sacred Heart College, Thevara, Cochin, Kerala, India −682013 (drpothalil@rediffmail.com)

#### **Abstract**

Although the Western Ghats are a biodiversity hotspot in the world, the studies conducted on the faunal affinity of this region are very few and there are no studies focused on spiders. Biota of the Western Ghats is the product of extreme climatic, ecological and biogeographical history. In this study, we tried to expose the faunal affinities of spiders distributed along this biodiversity hotspot based on their distribution throughout the world. A total of 270 species of 138 genera grouped in 39 families of spiders are so far reported from the Western Ghats of India. The biogeographical analysis revealed that a total of six genera of spiders are endemic to the Western Ghats, four genera showing affinities to the African region and 18 showing affinities to the Southeast Asian region. Another 20 genera are common to Africa, Western Ghats and Southeast Asian regions and 90 genera are cosmopolitan in distribution. These results agree with previous studies conducted on other faunal and floral elements of India, because India was part of the African continent until about 130 million years ago. It detached from there and later moved northwards and finally collided with Eurasia about 50 million years ago. In between, India was an isolated island for about 50 million years. This isolation might have lead to the creation of its unique fauna and flora. After the collision with Eurasia, the fauna of the Western Ghats underwent drastic changes due to faunal migrations between Southeast Asia and the Indian plate. There are different hypotheses explaining this migration of different animal groups through a continuous corridor of tropical evergreen forest. During this event, the Indian plate actually functioned as a "biotic ferry" which led to the spreading of African spiders to Southeast Asia. It is concluded that the spider fauna of the Western Ghats can be divided into an ancient African lineage, later Southeast Asian migrant and an endemic part. So a detailed phylogeographical approach is necessary to differentiate between these different time scale elements.

#### INTRODUCTION

One of the features of the biota of the Western Ghats is its strong affinity with African and Southeast Asian biota. There are several such cases of affinity in different taxa like mammals (Kurup 1974), birds (Ali 1935), freshwater fish (Jayaram 1974), amphibians (Das 2002), reptiles (Daniel 2002), insects (Singh 1974) and plants (Mani 1974a). Fauna and flora of the Western Ghats are an excep-

tionally interesting and challenging objective for biogeographers due to its antiquity, unique plate tectonic and palaeoclimatic history, and astounding array of physiognomic and habitat diversity (Ashton & Gunatilleke 1987). These factors have contributed to a considerable diversification of biota in this region (Morley 2000). Additionally, changes in the extents of habitats, especially of tropical evergreen forests in space and time, have resulted in striking endemism in almost all taxonomic groups, inviting much debate about processes underlying them (Swan 1993).

Dealing with complex palaeogeographical histories is a problem of major importance in biogeographical studies. Since the superposition of palaeogeographical events may produce multiple changes in species range distributions, highly complex patterns of animal and plant distributions are to be expected (Nelson & Platnick 1981; Myers & Giller 1988). As changes may interact and produce an apparent discordance between species distributions, the biogeographical relationships of areas are often difficult to trace back in time (McLennan & Brooks 2002; Brooks & Van Veller 2003). The Indian subcontinent exhibits such a complex biogeographical pattern, and several hypotheses based on different processes have been proposed to explain the presence of Southeast Asian and African faunal and floral elements in India (Mani 1974b). Some studies related this area as a sister area with the Southeast Asian region (Wallace 1876) or as sister area with the African region (Blanford 1876). Biogeographically the Indian region contains a complex assortment of barriers and dispersal routes varying in their nature as well as spatial scale to which multiple co-distributed taxonomic groups are unlikely to show a congruent response.

The spiders of the Western Ghats are a poorly worked out group compared to other parts of the country. With respect to its geographical, climatic and ecological features, the Western Ghats region harbours a rich amount of arachnids of which spiders have

a huge share. This area is also enriched with large forests and therefore possesses various assemblages of spiders. Due to high species endemism, the Western Ghats are listed among the 34 biodiversity hotspots of the world (Mittermeier et al. 2005). Due to the paucity of workers, much of the spider diversity in the Western Ghats has remained unexplored so far. As a result, the disappearance of many species remains undocumented. The present study provides the result of a preliminary analysis of the faunistic affinities of the spiders in the Western Ghats of India, which aims to demonstrate their great diversity and their high degree of endemism and to present a biogeographic hypothesis of the region.

#### **MATERIALS AND METHODS**

The Western Ghats occupy the western part of Peninsular India, between 8° and 20° North and between 73° to 77° East. The Western Ghats cover a practically unbroken relief dominating the west coast of the Indian peninsula for almost 1600 km from the river Tapti in the north to the tip of Peninsular India. Zoogeographically, the Western Ghats can be divided into three parts: a northern division, comprising the Deccan Trap area from the Tapti river down to 16° N, about the region of Goa; a central division, extending from 16° N southwards and including the Kudagu and Wayanad region of south Karnataka and Nilgiris; and a southern division, comprising the Anaimalai, Palani and Cardamom hills of Kerala (Bhimachar 1945). The Western Ghats are the main watershed in peninsular India from which all the principal rivers, namely Godavary, Krishna and Cauvery originate and flow east emptying into the Bay of Bengal. A large number of short perennial, torrential west flowing rivers also originate from it. The average height of the Ghats is less than 1500 m above sea level, but in the south it rises up to 2000 m and to exceptionally high peaks of 2500 m and above. Along its entire length, this hill range has only one discontinuity, the Palghat gap in Kerala, a more than 30 km wide gap, less than 100 m above sea level (Nair 1991).

The presence of spiders, which are reported from the Western Ghats, in other geographic regions of the world was collected from the literature. A checklist of spiders so far reported from the Western Ghats was prepared based on the world spider catalogue (Platnick 2008). Each species was classified according to its distribution in the African region and Southeast Asian region, and it was stated whether its distribution is endemic or cosmopolitan, *i. e.*, distribution in the tropics throughout the world.

#### **RESULTS**

A total of 270 species of 138 genera listed in 39 families of spiders are so far reported from the Western Ghats (Appendix 1). Among these 138 genera, six genera are endemic; Diplothele and Sasonichus of the family Barychelidae; Annandaliella, Haploclastus, Poecilotheria and Thrigmopoeus of the family Theraphosidae. A total of 91 species are endemic in the Western Ghats. A total of four genera shows affinities with the African region but there is no common species with distribution in the Western Ghats and Africa. A total of 18 genera shows affinities with the Southeast Asian region and a total of 68 species share a distribution in these two regions. Although there are species of another 20 genera showing a distribution in Africa, Southeast Asia and the Western Ghats, there is no common species between these regions. Among the 270 species of spiders reported from the Western Ghats, a total of 90 genera and 111 species are distributed in the tropics throughout the world.

#### **DISCUSSION**

India's peculiar and diverse biota has attracted the attention of researchers for more than a century. Earlier workers noticed close similarities with Africa's biota, which sparked a perpetual debate on the origins and biogeographical relationships of India's

flora and fauna. The question of whether the links with Southeast Asia or Africa are more important has been a central aspect of this discussion. It culminated in the refusal of Wallace's (1876) hypothesis that India's biota was essentially of Southeast Asian origin, by Blanford (1876), who in turn emphasized a strong African influence on India's fauna and flora.

As the present study revealed the occurrence of both African and Southeast Asian influence on the spider fauna of the Western Ghats, it has been related to the geographic history of India. The geographic history of India can be summarised as follows based on plate tectonics and continental drift. First, the Indian subcontinent detached from Africa – 130 million years ago (Krause et al. 1997), as part of the Madagascar-Seychelles-India block. Its long northward drift across the Tethys sea, with disconnection from Madagascar at – 88 million years ago (Storey et al. 1995) and the Seychelles at – 65 million years ago (Courtillot et al. 1988), ended only in the Palaeogene (Najman et al. 2001), after accretion to the Eurasian block. The first contact between both landmasses immediately enabled Southeast Asian animals and plants to invade the subcontinent (Briggs 1989), and lineages of old African origin, if they persisted on the drifting subcontinent, to disperse to Southeast Asia (Bossuyt & Milinkovitch 2001; Conti et al. 2002; Gower et al. 2002; Wilkinson et al. 2002). During this million years long process, the movement of the Indian subcontinent over the Reunion mantle plume at the Cretaceous-Tertiary transition (K-T transition) generated the notorious Deccan basalt floods (Courtillot et al. 1988) that almost wiped out old African lineage, which resulted in the formation of the Western Ghats. Upon impact with Asia in the Early Tertiary, the uplift of the Himalayan chain and subsequent Late Tertiary aridification further contributed to the progressive impoverishment of African elements from the Indian biota, with the exception of refugial areas in India (Raven & Axelrod 1974; Guleria 1992; Morley 2000). So the presence of four genera of spiders in the Western Ghats with African affinity can be explained by their common origin in former Gondwanaland and by vicariance.

Among the 138 genera of spiders reported from the Western Ghats, 20 genera share distribution in the African region and the Southeast Asian region along with the Western Ghats. As much of Southeast Asia was never part of old Africa, the presence of African forms in Southeast Asian region is intriguing. The main hypothesis for the presence of African elements in Southeast Asia is that fauna and flora associated with the drifting Indian plate were dispersed into Southeast Asia following accretion with Eurasia (Wilkinson et al. 2002). This hypothesis known as "out-of-India" hypothesis (McKenna 1973), has been proposed to explain the presence of African forms in Southeast Asia. The geophysical evidence for this scenario is well known (Briggs 2003a). Thus it is plausible that rafting peninsular India carried with it African forms to Asia. Mani (1974b) points out that with the physical contact of peninsular India with Asia, extensive interchange between Indian and Southeast Asian flora and fauna occurred. In all these studies, the drifting Indian landmass is perceived as a "biotic ferry" for old African groups. This biotic ferry model (McKenna 1973; Morley 2000; Briggs 2003b) became the standard explanation for the presence of African elements in Southeast Asia. Based on this model, a number of studies in different taxa like amphibians (Bossuyt & Milinkovitch 2001; Gower et al. 2002; Wilkinson et al. 2002) and plants (Conti et al. 2002) suggested that African elements colonized South and Southeast Asia as "out-of-India".

A total of 18 genera and 68 species are commonly present in the Western Ghats and Southeast Asia. Some taxonomic (Hora 1949; Bande & Prakash 1986; Bande 1992) and recent molecular (Krause & Maas 1990; Clyde et al. 2003) studies revealed the dispersion of organisms from the Southeast Asian region

into India, for which the term "out of Asia" hypothesis was coined by Hedges et al. (1993). It has been proposed that, during the Early Tertiary, Southeast Asian lineages reached India over temporary land connections almost immediately after the collision with Eurasia. Many paleontological studies also suggested that the latest Cretaceous-Paleocene Indian fauna and flora which survived extensive volcanic activities was almost completely replaced by the diverse and relatively advanced biota from Southeast Asia upon the India-Asia collision (Briggs 2003b).

Landmasses that have experienced a prolonged period of extensive isolation may lead to the evolution of their own fauna and flora. Such high level endemism is apparent on large islands or island groups, such as New Zealand (Hay et al. 1995), the Seychelles (Ruvinsky & Maxson 1996) and Madagascar (Vences et al. 2000). When long term isolation is followed by restoration of contact with other regions, the biotic uniqueness of an area may gradually fade due to floral and faunal interchange. Nevertheless, some previously isolated regions may incidentally retain inconspicuous remnants of a unique ancient biotic composition. During the journey of India from Africa to Asia it was in an isolated condition during 50 million years. When continents or islands are well isolated for such extended periods of time, evolution invariably produces endemic species. If the isolation continues long enough, a host of peculiar genera and families may appear. In this study a total of six genera and 91 species of spiders are found as endemic. The presence of some endemic genera in spiders of the Western Ghats, provides strong evidence that India underwent an extended isolation during its journey across the middle of the Indian Ocean. Recent molecular dating estimates in 14 species of amphibians (Bossuyt & Milinkovitch 2001) indicated that several lineages originated on the Indian subcontinent during its trans-Tethys drift. Remarkable suites of endemic species are characteristic of isolated land masses. This

is particularly true for India, where the biota is well known for its extraordinary high levels of endemism, with species frequently confined to minute distributional ranges (Myers et al. 2000).

Although fauna and flora of India originated in Africa, present similarity with African region is very low. This may be due to the faunal interchange with the Southeast Asian region and the evolution of its own biota during the period of isolation. These intruders from Southeast Asia not only led to the dilution of its original fauna but also dominated gradually as described in this study. So a detailed study using modern phylogenetic methodology is needed to differentiate between these faunal elements and to estimate the age of these different components in spiders of India, especially the Western Ghats region.

#### **ACKNOWLEDGEMENTS**

This study was supported by Special Research Fund (BOF) from Ghent University, Belgium to the first author.

#### REFERENCES

- Ali, S. 1935. The ornithology of Travancore and Cochin. *Journal of Bombay Natural History Society* 37: 814–843.
- Ashton, P.S. & Gunatilleke, C.V.S. 1987. New light on the plant biogeography of Ceylon. I. Historical plant geography. *Journal of Biogeography* 14: 249–285.
- Bande, M.B. 1992. The Palaeogene vegetation of peninsular India (megafossil evidence). *Palaeobotanist* 40: 275–284.
- Bande, M.B. & Prakash, U. 1986. The Tertiary flora of Southeast Asia with remarks on its palaeoenvironment and phytogeography of the Indo-Malayan region. *Revue of Paleobotany and Palynology* 49: 203–233.
- Bhimachar, B.S. 1945. Zoogeographical divisions of Western Ghats, as evidenced by the distribution of the hill stream fishes. *Current Science* 1: 12–16.
- Blanford, W.T. 1876. The African element in the fauna of India: a criticism of Mr.

- Wallace's views as expressed in the 'Geographical Distribution of Animals'. *Annals* and Magazine of Natural History, 4<sup>th</sup> Series 18: 277–294.
- Bossuyt, F. & Milinkovitch, M.C. 2001. Amphibians as indicators of Early Tertiary Out-of-India dispersal of vertebrates. *Science* 292: 92–95.
- Briggs, J.C. 1989. The historic biogeography of India: isolation or contact? *Systematic Zoology* 38: 322–332.
- Briggs, J.C. 2003a. Fishes and birds: Gondwana life rafts reconsidered. *Systematic Biology* 52: 548–553.
- Briggs, J.C. 2003b. The biogeographical and tectonic history of India. *Journal of Biogeography* 30: 381–388.
- Brooks, D.R. & Van Veller, M.G.P. 2003. Critique of parsimony analysis of endemicity as a method of historical biogeography. *Journal of Biogeography* 30: 819–825.
- Clyde, W.C., Khan, I.C.H & Gingerich, P.D. 2003. Stratigraphic response and mammalian dispersal during initial India Asia collision: evidence from the Ghazij Formation, Balochistan, Pakistan. *Geology* 31: 1097–1100.
- Conti, E., Eriksson, T., Schronenberger, J., Sytsma, K.J. & Baum, D.A.2002. Early Tertiary Out-of-India dispersal of Crypteroniaceae: evidence from phylogeny and molecular dating. *Evolution* 56: 1931–1942.
- Courtillot, V., Feraud, G., Maluski, H., Vandamme, D., Moreau, M.G. & Besse, J. 1988. Deccan flood basalts and the Cretaceous/ Tertiaty boundary. *Nature* 333: 843–846.
- Daniel, J.C. 2002. The book of Indian reptiles and amphibians. Oxford University Press, Mumbai.
- Das, I. 2002. An Introduction to the Amphibians and Reptiles of Tropical Asia. Natural History Publications (Borneo), Kota Kinabalu.
- Gower, D.J., Kupfer, A., Oommen, O.V., Himstedt, W., Nussbaum, R.A., Loader, S.P., Presswell, B., Muller, H., Krishna, S.B., Boistel, R. & Wilkinson, M. 2002. A molecular phylogeny of ichthyophiid caecil-

- ians (Amphibia: Gymnophiona: Ichthyophiidae): out of India or out of South East Asia? *Proceedings of Royal Society of London. B* 269: 1563–1569.
- Guleria, J.S. 1992. Neogene vegetation of peninsular India. *Paleobotanist* 40: 285–311.
- Hay, J.M., Ruvinsky, I., Hedges, S.B. & Maxson, L.R. 1995. Phylogenetic relationships of amphibian families inferred from DNA sequences of mitochondrial 12S and 16S ribosomal RNA genes. *Molecular Biology and Evolution* 12: 928–937.
- Hedges, S.B., Nussbaum, R.A. & Maxson, L.R. 1993. Caecilian phylogeny and biogeography inferred from mitochondrial DNA sequences of the 12S rRNA and 16S rRNA genes (Amphibia: Gymnophiona). *Herpetological Monographs* 7: 64–76.
- Hora, S.L. 1949. Satpura hypothesis of the distribution of Malayan fauna and flora of peninsular India. *Proceedings of the National Institute of Science, India* 15: 309–314.
- Jayaram, K.C. 1974. Ecology and distribution of freshwater fishes, amphibia and reptiles. Pp. 517–584. In: *Ecology and biogeography in India* (Mani, M.S., ed.). Dr. W. Junk B. V. Publishers, The Hague.
- Krause, D.W. & Maas, M.C. 1990. The biogeographic origins of the late Paleocene-early Eocene mammalian immigrants to the western interior of North America. Pp. 71–105. In: *Dawn of the age of mammals in the northern part of the Rocky Mountain interior, North America* (Bown, T.M. & Rose, K.D., eds.). Geological Society of America. Special Paper.
- Krause, D.W., Prasad, G.V.R., von Koenigswald, W., Sahni, A. & Grine, F.E. 1997. Cosmopolitanism among Late Cretaceous Gondwanan mammals. *Nature* 390: 504–507.
- Kurup, G.U. 1974. Mammals of Assam and the mammal-geography of India. Pp. 585–613. In: *Ecology and biogeography in India* (Mani, M.S., ed.). Dr. W. Junk B. V. Publishers, The Hague.
- Mani, M.S. 1974a. The flora. Pp. 159–177. In: *Ecology and biogeography in India* (Mani,

- M.S., ed.). Dr. W. Junk B. V. Publishers, The Hague.
- Mani, M.S. 1974b. Biogeographical evolution in India. Pp. 698–722. In: *Ecology and biogeography in India* (Mani, M.S., ed.). Dr W. Junk B. V. Publishers, The Hague.
- McKenna, M.C.C. 1973. Sweepstakes, filters, corridors, Noah's arks, and beached Viking funeral ships in palaeography. Pp. 291–304. In: *Implications of continental drift to the earth sciences* (Tarling, D.H. & Runcorn, S.K., eds.). Academic Press, London.
- McLennan, D.A. & Brooks, D.R. 2002. Complex histories of speciation and dispersal in communities: a re-analysis of some Australian bird data using BPA. *Journal of Biogeography* 29: 1055–1066.
- Mittermeier, R.A., Patricio, R.G., Hoffman, M., Pilgrim, J., Brooks, T., Mittermeier. C.G., Lamoreux, J. & Da Fonseca, G.A.B. 2005. Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions. *Conservation International*, Pp. 1–432.
- Morley, R.J. 2000. *Origin and evolution of tropical rain forests*. John Wiley and Sons, Chichester, U.K.
- Myers, A.A. & Giller, P.S. 1988. Analytical biogeography: an integrated approach to the study of animals and plant distributions. Chapman and Hall, London.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Da Fonseca, G.A.B. & Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Nair, S.C. 1991. *The Southern Western Ghats a biodiversity conservation plan*. New Delhi, INTACH.
- Najman, Y., Pringle, M., Godin, L. & Oliver, G. 2001. Dating of the oldest continental sediments from the Himalayan foreland basin. *Nature* 410: 194–197.
- Nelson, G. & Platnick, N.I. 1981. *Systematics and biogeography: cladistics and vicariance*. Columbia University Press, New York.
- Platnick, N.I. 2008. The world spider catalog, version 8.5. American Museum of Natural History, online at http://research.

- amnh.org/entomology/spiders/catalog/index.htm
- Raven, P.H. & Axelrod, D.I. 1974. Angiosperm biogeography and past continental movements. *Annals of Missouri Botanical Garden* 61: 539–673.
- Ruvinsky, I. & Maxson, L.A. 1996. Phylogenetic relationships among bufonoid frogs (Anura: Neobatrachia) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 5: 533–547.
- Singh, S. 1974. Some aspects of the ecology and geography of Diptera. Pp. 500–516. In: *Ecology and biogeography in India* (Mani, M.S., ed.). Dr. W. Junk B. V. Publishers, The Hague.
- Storey, M., Saunders, A.D., Duncan, R.A., Kelley, S.P. & Coffin, M.F. 1995. Timing of hotspot-related volcanism and the break-up of Madagascar and India. *Science* 267: 852–855.
- Swan, L.W. 1993. The Satpura hypotheses: a biogeographical challenge to geology. *Journal of Bombay Natural History Society* 90: 141–157.
- Vences, M., Glaw, F., Kosuch, J., Das, I. & Veith, M. 2000. Polyphyly of *Tomopterna* (Amphibia: Ranidae) based on sequences of the mitochondrial 16S and 12S rRNA genes, and ecological biogeography of Madagascan relict amphibian groups. Pp. 229–242. In: *Diversité et Endémisme à Madagascar* (Lourenço, W.R. & Goodman, S.M., eds.). Memories of Biogeographical Scociety, Paris.
- Wallace, A.R. 1876. The geographical distribution of animals with a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface. Harper and Brothers, New York.
- Wilkinson, J.A., Drewes, R.C. & Tatum, O.L. 2002. A molecular phylogenetic analysis of the family Rhacophoridae with an emphasis on the Asian and African genera. *Molecular Phylogenetics and Evolution* 24: 265–273.

#### **APPENDIX 1**

## Check list of spiders reported from the Western Ghats of India

- \* Endemic species.
- \*\* Species distributed in the Southeast Asian region.
- † Endemic genus.
- tt Genus with African affinity.
- +++ Genus with species distribution in the African region, the Southeast Asian region and the Western Ghats.
- ++++ Genus with species distribution in the Southeast Asian region.

#### FAMILY AGELENIDAE C.L. Koch, 1837

Agelena kariansholaensis\* Sugumaran et al. 2005

Agelena satmila\* Tikader, 1970

#### FAMILY ARANEIDAE Simon, 1895

Arachnura+++ angura\* Tikader, 1970

Arachnura scorpionoides\* Sugumaran et al., 2005

Araneus bilunifer Pocock, 1900

Araneus ellipticus (Tikader & Bal, 1981)

Araneus himalayaensis Tikader, 1975

Araneus nympha\*\* (Simon, 1889)

Argiope aemula (Walckenaer, 1842)

Argiope anasuja Thorell, 1887

Argiope catenulata (Doleschall, 1859)

Argiope pulchella Thorell, 1881

Chorizopes+++ bengalensis Tikader, 1975

Chorizopes calciope\*\* (Simon, 1895)

Cyclosa anaikattae\* Sugumaran et al., 2005

Cyclosa bifida (Doleschall, 1859)

Cyclosa confraga (Thorell, 1892)

Cyclosa hexatuberculata\* Tikader, 1982

Cyclosa insulana (Costa, 1834)

Cyclosa moonduensis\* Tikader, 1963

Cyclosa mulmeinensis (Thorell, 1887)

Cyclosa quinqueguttata (Thorell, 1883)

Cyclosa spirifera Simon, 1889

Cyrtarachne bengalensis\*\* Tikader, 1961

Cyrtophora bidenta\* Tikader, 1970

Cyrtophora cicatrosa\*\* (Stoliczka, 1869)

Cyrtophora citricola (Forskal, 1775)

Cyrtophora koronadalensis Barrion & Litsinger, 1995 Eriophora himalayaensis (Tikader, 1975)

Eriovixia+++ excelsa\*\* (Simon, 1889)

Eriovixia laglaizei\*\* (Simon, 1877)

Eriovixia poonaensis\* (Tikader & Bal, 1981)

Gasteracantha dalyi\*\* Pocock, 1900

Gasteracantha geminata (Fabricius, 1798)

Gasteracantha hasselti\* C.L. Koch, 1837

Gasteracantha kuhli Koch, 1837

Gasteracantha remifera Butler, 1873

Gea subarmata\*\* Thorell, 1890

Gibbaranea bituberculata\*\* (Walckenaer, 1802)

Heurodes++++ porculus\*\* (Simon, 1877)

Macracantha++++ arcuata\*\* (Fabricius, 1793)

Neoscona bengalensis Tikader & Bal, 1981

Neoscona molemensis\*\* Tikader & Bal, 1981

Neoscona mukerjei Tikader, 1980

Neoscona nautica\*\* (L. Koch, 1875)

Neoscona parambikulamensis\* Patel, 2003

Neoscona pavida\* (Simon, 1906)

Neoscona theisi (Walckenaer, 1842)

Neoscona vigilans (Blackwall, 1865)

Parawixia dehaani\*\* (Doleschall, 1859)

Parawixia mundanthuraiensis\* Sugumaran et al., 2005

Poltys+++ columnaris\*\* Thorell, 1890

Zygiella indica\* Tikader & Bal, 1980

#### FAMILY ATYPIDAE Thorell, 1870

Atypus sutherlandi Chennappaiya, 1935

#### FAMILY BARYCHELIDAE Simon, 1889

Sason††† robustum (O. P.-Cambridge, 1883) Sasonichus† sullivani\* Pocock, 1900

#### FAMILY CLUBIONIDAE Wagner, 1887

Clubiona coimbatorensis\* Sugumaran et al., 2005 Clubiona drassodes O. P.-Cambridge, 1874

#### FAMILY CORINNIDAE Karsch, 1880

Castianeira zetes Simon, 1897 Oedignatha††† carli Reimoser, 1934 Oedignatha microscutata Reimoser, 1934 Oedignatha scrobiculata Thorell, 1881

#### FAMILY CTENIDAE Keyserling, 1877

Acanthies++++ indicus\* Gravely, 1931 Ctenus cochinensis\* Gravely, 1931 Ctenus indicus\* Gravely, 1931

#### FAMILY DEINOPIDAE C.L. Koch, 1850

Deinopis goalparaensis\* Tikader & Malhotra, 1978

## FAMILY DICTYNIDAE O. P.-Cambridge, 1871

Dictyna umai\* Tikader, 1966

#### FAMILY DIPLURIDAE Simon, 1889

Indothele† mala\* Coyle, 1995

#### FAMILY ERESIDAE Koch, 1851

Stegodyphus pacificus Pocock, 1900 Stegodyphus sarasinorum\*\* Karsch, 1891 Stegodyphus tibialis (O. P.-Cambridge, 1869)

### FAMILY FILISTATIDAE Ausserer, 1867

Pritha poonaensis\* (Tikader, 1963)

#### FAMILY GNAPHOSIDAE Pocock, 1898

Drassodes carinivulvus\* Caporiacco, 1934 Gnaphosa kailana\* Tikader, 1966 Poecilochroa barmani\* Tikader, 1982 Setaphis subtilis\*\* (Simon, 1897) Zelotes ashae\* Tikader & Gajbe, 1976

#### FAMILY HAHNIIDAE Bertkau, 1878

Hahnia alini\* Tikader, 1964

#### FAMILY HERSILIIDAE Thorell, 1870

Hersilia+++ pectinata Thorell, 1895 Hersilia savignyi Lucas, 1836 Tama gravelyi \*\* Sinha, 1950

#### FAMILY LINYPHIIDAE Blackwall, 1859

Atypena simoni\* Jocqué, 1983 Lepthyphantes rudrai\*\* Tikader, 1970 Linyphia urbasae\*\* Tikader, 1970 Neriene sundaica\*\* (Simon, 1905)

#### FAMILY LYCOSIDAE Sundevall, 1833

Crocodilosa††† leucostigma (Simon, 1885) Evippa†† banarensis \* Tikader & Malhotra, 1980 Hippasa†† agelenoides\* (Simon, 1884) Hippasa greenalliae\* (Blackwall, 1867) Hippasa holomerae\* Thorell, 1895 Hippasa leucostigma\* Simon, 1885 Hippasa lycosina\* Pocock, 1900 Hippasa olivacea (Thorell, 1887)
Hippasa pisaurina\* Pocock, 1900
Lycosa barnesi\*\* Gravely, 1924
Lycosa bistriata\* Gravely, 1924
Lycosa carmichaeli Gravely, 1924
Lycosa madani Pocock, 1901
Lycosa tista\*\* Tikader, 1970
Pardosa atropalpis Gravely, 1924
Pardosa minuta\* Tikader & Malhotra, 1976
Pardosa oakleyi\* Gravely, 1924
Pardosa pseudoannulata\*\* (Bosenberg & Strand,

1906)
Pardosa sumatrana (Thorell 1890)

Pardosa sumatrana (Thorell, 1890) Wadicosa quadrifera (Gravely, 1924)

#### FAMILY MIMETIDAE Simon, 1881

Mimetus indicus\* Simon, 1906

#### FAMILY MITURGIDAE Simon, 1885

Cheiracanthium danieli Tikader, 1975 Cheiracanthium insulanum\*\* (Thorell, 1878) Cheiracanthium melanostomum (Thorell, 1895) Cheiracanthium triviale (Thorell, 1895)

#### FAMILY NEPHILIDAE Simon, 1894

Herennia++++ multipuncta\*\* (Doleschall, 1859) Nephila kuhlii\*\* (Doleschall, 1859) Nephila pilipes\*\* (Fabricius, 1793) Nephilengys malabarensis\* (Walckenaer, 1842)

## FAMILY OECOBIIDAE Blackwall, 1862

Oecobius putus O. P.-Cambridge, 1876

#### FAMILY OONOPIDAE Simon, 1890

Opopaea++ sponsa\* Brignoli, 1978

#### FAMILY OXYOPIDAE Thorell, 1870

Oxyopes ashae\* Gajbe, 1999
Oxyopes bharatae\* Gajbe, 1999
Oxyopes birmanicus\*\* Thorell, 1887
Oxyopes hindostanicus Pocock, 1901
Oxyopes javanus Thorell, 1887
Oxyopes lineatipes\*\* (C.L. Koch, 1847)
Oxyopes rukminiae\* Gajbe, 1999
Oxyopes sakuntalae Tikader, 1970
Oxyopes shweta Tikader, 1970
Oxyopes guadridentatus\*\* Thorell, 1895

Oxyopes sunandae Tikader, 1970 Oxyopes wroughtoni Pocock, 1901 Peucetia viridana (Stoliczka, 1869)

### FAMILY PHILODROMIDAE Thorell, 1870

Philodromus ashae\* Gajbe & Gajbe, 1999 Tibellus elongatus\* Tikader, 1960

#### FAMILY PHOLCIDAE C.L. Koch, 1851

Artema atlanta Walckenaer, 1837 Crossopriza lyoni (Blackwall, 1867) Pholcus phalangioides (Fuesslin, 1775) Uthina++++ atrigularis\*\* Simon, 1901

#### FAMILY PISAURIDAE Simon, 1890

Perenethis††† dentifasciata\*\* (O. P.-Cambridge, 1885)

Perenethis unifasciata\*\* (Doleschall, 1859) Pisaura mirabilis (Clerck, 1757) Polyboea vulpina Thorell, 1895 Thalassius+++ albocinctus (Doleschall, 1859)

## FAMILY PRODIDOMIDAE Simon, 1884

Zimiris indica\* Dyal, 1935

#### FAMILY PSECHRIDAE Simon, 1890

Fecenia++++ travancoria\* Pocock, 1899 Psechrus++++ torvus\*\* (Cambridge, 1869)

#### FAMILY SALTICIDAE Blackwall, 1841

Aelurillus improvisus\* Azarkina, 2002
Bavia+++ kairali \* Samson & Sebastian, 2004
Bianor angulosus\* (Karsch, 1879)
Brettus+++ anchorum \* Wanless, 1979
Hasarius adansoni (Audouin, 1826)
Hyllus+++ diardi (Walckenaer, 1837)
Hyllus semicupreus (Simon, 1885)
Menemerus bivittatus (Dufour, 1831)
Myrmarachne markaha\*\* Barrion & Litsinger,
1995

Myrmarachne orientales Tikader, 1973 Myrmarachne plataleoides (O. P.-Cambridge, 1869)

Phintella vittata (C.L. Koch, 1846)
Plexippus chandraseharani\* Samiayyan, 1995
Plexippus dharineae\* Samiayyan, 1995
Plexippus paykulli (Audouin, 1826)
Plexippus petersi (Karsch, 1878)

Portia+++ fimbriata\* (Doleschall, 1859)
Pseudicius daitaricus\* Prószyński, 1992
Rhene+++ danieli Tikader, 1973
Rhene flavigera\*\* (C.L. Koch, 1846)
Rhene rubrigera (Thorell, 1887)
Stenaelurillus lesserti\* Reimoser, 1934
Tamigalesus munnaricus\* Zabka, 1988
Telamonia+++ dimidiata\*\* (Simon, 1899)
Thiania++++ bhamoensis\*\* Thorell, 1887

#### FAMILY SCYTODIDAE Blackwall, 1864

Scytodes fusca Walckenaer, 1837 Scytodes thoracica (Latreille, 1802)

#### FAMILY SELENOPIDAE Simon, 1897

Selenops montigenus\*\* Simon, 1889 Selenops radiatus\*\* Latreille, 1819

#### FAMILY SPARASSIDAE Bertkau, 1872

Heteropoda hampsoni\*\* Pocock, 1901
Heteropoda lentula\*\* Pocock, 1901
Heteropoda leprosa\*\* Simon, 1884
Heteropoda lunula (Doleschall, 1857)
Heteropoda nicobarensis Tikader, 1977
Heteropoda nilgirina\* Pocock, 1901
Heteropoda phasma\*\* Simon, 1897
Heteropoda venatoria (Linnaeus, 1767)
Micrommata virescens\*\* (Clerck, 1757)
Olios hampsoni\*\* (Pocock, 1901)
Olios obesulus\*\* (Pocock, 1901)
Palystes++ flavidus\* Simon, 1897
Thelcticopis+++ maindroni\* Simon, 1906

## **FAMILY STENOCHILIDAE** Thorell, 1873 *Stenochilus*†††† *hobsoni*\*\* O. P.-Cambridge, 1870

#### FAMILY TETRAGNATHIDAE Menge, 1866

Dyschiriognatha++++ dentata\*\* Zhu & Wen, 1978
Leucauge bituberculata Baert, 1987
Leucauge celebesiana\*\* (Walckenaer, 1842)
Leucauge decorata (Blackwall, 1864)
Leucauge dorsotuberculata Tikader, 1982
Leucauge pondae Tikader, 1970
Leucauge subgemmea Bösenberg & Strand, 1906
Leucauge tessellata (Thorell, 1887)
Opadometa++++ fastigata\*\* (Simon, 1877)
Orsinome+++ marmorea\*\* Pocock, 1901

Tetragnatha andamanensis Tikader, 1977
Tetragnatha ceylonica\*\* O. P.-Cambridge, 1869
Tetragnatha cochinensis Gravely, 1921
Tetragnatha fletcheri\*\* Gravely, 1921
Tetragnatha javana\*\* (Thorell, 1890)
Tetragnatha mandibulata\*\* Walckenaer, 1842
Tetragnatha maxillosa\*\* Thorell, 1895
Tetragnatha nitens\*\* (Audouin, 1826)
Tetragnatha siruvaniensis Sugumaran et al., 2005
Tetragnatha sutherlandi Gravely, 1921
Tetragnatha vermiformis Emerton, 1884
Tetragnatha viridorufa Gravely, 1921
Tylorida††† culta\*\* (O. P.-Cambridge, 1869)
Tylorida ventralis\*\* (Thorell, 1877)

#### FAMILY THERAPHOSIDAE Thorell, 1870

Anandaliella† travancorica\* Hirst, 1909
Chilobrachys††† fimbriatus\*\* Pocock, 1899
Haploclastus† kayi\* Gravely, 1915
Haploclastus nilgirinus\* Pocock, 1899
Ischnocolus decoratus\* Tikader, 1977
Plesiophrictus††† bhori\* Gravely, 1915
Plesiophrictus raja\* Gravely, 1915
Poecilotheria† regalis\* Pocock, 1899
Poecilotheria rufilata\* Pocock, 1899
Poecilotheria striata\* Pocock, 1895
Thrigmopoeus† parambikulamensis\* Sanjay & Daniel, 2002

#### FAMILY THERIDIIDAE Sundevall, 1833

Achaearanea diglipuriensis Tikader, 1977 Achaearanea durgae Tikader, 1970 Achaearanea mundula\*\* (L. Koch, 1872) Achaearanea triangularis\* (Patel, 2003) Argyrodes ambalikae Tikader, 1970 Argyrodes flavescens O. P.-Cambridge, 1880 Argyrodes gazedes\* Tikader, 1970 Argyrodes gazingensis\* Tikader, 1970 Argyrodes xiphias Thorell, 1873 Ariannes flagellum (Doleschall, 1857) Chrysso argyrodiformis (Yaginuma, 1952) Chrysso isumbo Barrion & Litsinger, 1995 Chrysso nigra (O. P.-Cambridge, 1880) Coleosoma†††† floridanum\*\* Banks, 1900 Faiditus xiphias Thorell, 1887 Phycosoma martinae (Roberts, 1983) Theridion incertum O. P.-Cambridge, 1885

Theridion manjithar\* Tikader, 1970 Theridion otsospotum Barrion & Litsinger, 1995 Theridula angula Tikader, 1970 Steatoda alboclathrata\* (Simon, 1897)

#### FAMILY THOMISIDAE Sundevall, 1833

Camaricus+++ formosus\* Thorell, 1887
Camaricus khandalaensis\* Tikader, 1980
Misumena decorata Tikader, 1980
Misumena silveryi\* Tikader, 1965
Misumenops andamanensis Tikader, 1980
Oxytate virens (Thorell, 1891)
Ozyptila amkhasensis\* Tikader, 1980
Pistius bhadurii\* Basu, 1965
Strigoplus+++ netravathi\* Tikader, 1963
Thomisus andamanensis Tikader, 1980
Thomisus beautifularis\* Basu, 1965
Thomisus lobosus Tikader, 1965
Thomisus pugilis Stoliczka, 1869
Xysticus himalayaensis Tikader & Biswas, 1974

#### FAMILY ULOBORIDAE Thorell, 1869

Miagrammopes extensus Simon, 1889 Philoponella hilaris\* (Simon, 1906) Uloborus coimbatoriensis Sugumaran et al., 2005 Uloborus danolius\*\* Tikader, 1969 Uloborus krishnae\* Tikader, 1970 Zosis geniculata\*\* (Olivier, 1789)

#### FAMILY ZODARIIDAE Thorell, 1881

Asceua+++ cingulata\* (Simon, 1905) Cryptothele collina\*\* Pocock, 1901 Cryptothele sundaica\* Thorell, 1890 Hermippus++ arjuna\* (Gravely, 1921) Storena arakuensis\* Patel & Reddy, 1989