

Zeitschrift: Contributions to Natural History : Scientific Papers from the Natural History Museum Bern

Herausgeber: Naturhistorisches Museum Bern

Band: - (2009)

Heft: 12/1

Artikel: Taxonomic review of some Drassodes species from Greece and other East Mediterranean countries (Araneae: Gnaphosidae)

Autor: Chatzopoulou, Eli / Chatzaki, Maria

DOI: <https://doi.org/10.5169/seals-786972>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Taxonomic review of some *Drassodes* species from Greece and other East Mediterranean countries (Araneae: Gnaphosidae)

Eli Chatzopoulou & Maria Chatzaki

ABSTRACT

Contrib. Nat. Hist. 12: 349–359.

This paper summarizes the results of a study on the Drassodinae collections of the National Museum of Natural History of Paris, France and of the Natural History Museum of the University of Crete, Greece. *Drassodes alexandrinus* (O.P. CAMBRIDGE, 1874) is proposed as a new synonym of *D. lutescens* (C.L. KOCH, 1839), based on an extensive examination of specimens from insular and continental Greece and from adjacent East Mediterranean countries. Further synonymizations are proposed: *D. carinatus* STRAND, 1906 is placed in the synonymy of *Haplodrassus dalmatensis* (L. KOCH, 1866), and *D. oreinos* CHATZAKI, 2002 is placed in the synonymy of *D. albicans* (SIMON, 1878). The above results change the catalogue of the *Drassodes* species recorded on Crete and reduce the percentage of endemism in the area.

Introduction

The genus *Drassodes* is one of the most diverse genera of the family Gnaphosidae worldwide. In Greece it is the second species richest genus, together with *Haplodrassus* (10 species) and following *Zelotes* (28 species). Quite a number of species earlier described as belonging to this genus were either misidentified or transferred to other genera that were created later as subgroups of this large genus (for example *Haplodrassus*, *Leptodrassus*, *Leptodrassex* etc.). Because of the simplicity of the genitalic characters, the distinction of species becomes difficult, especially in the case of very closely related species. Very often taxonomists have created new species based on very small differences between specimens, mainly because most of the time they did not have a high number of vouchers, hence intraspecific variation could not be detected. In a recent revision on the Gnaphosidae of Crete, Chatzaki & al. (2002a, b) came up with 3 new combinations (transfers to other genera), 3 synonymizations




Fig. 1. Map of sampling localities in Greece.

and 1 new species of *Drassodes* from Crete, resulting in a total number of only 4 species on the island. Levy (2004) defined 12 new synonyms within this genus, thus reducing the total number of the existing species in the East Mediterranean area as well.

In this paper some of the *Drassodes* species occurring in the area of Greece and neighboring regions are reviewed, based on the close examination of specimens from insular and continental Greece and from adjacent East Mediterranean countries. The examination of comparative material at the National Museum of Natural History of Paris (MNHN) led to the proposal of further synonymizations.

Materials and Methods

Material comes from multiple expeditions by members of the Natural History Museum of the University of Crete (NHMC) using either pitfall trapping or hand collecting. The islands reported here are shown in Fig. 1. Material from




Fig. 2. Sketch of *Drassodes* genitalia with abbreviations measured for morphometric analyses. – ♂: Embolic origin (EO), Tegulum length (Te), Cymbium length (Cy). – ♀: Septum width, maximum (MS) and minimum (mS), Lateral pouches length (LP), Spermathecae length (SL).

adjacent countries was loaned by several colleagues: material from Bulgaria was loaned by Mr. Dolanský, material from Syria was loaned by Dr. Řezáč and material from Israel was loaned by the late Dr. Gershon Levy.

The following morphometric measurements were needed for the taxonomic analysis (see Fig. 2): ♂ – Embolic origin (EO), Tegulum length (Te), Cymbium length (Cy), Pedipalpal tibia length measured laterally (Ti). ♀ – Septum width, maximum (MS) and minimum (mS), Lateral pouches length (LP), Spermathecae length (SL).

All drawings were made by the last author.

Only relevant primary and secondary citations were considered in the lists of synonymies for each species. For a complete list, compare Platnick (2009).


Results and Discussion

Drassodes lutescens (C.L. KOCH, 1839) (Figs. 3–13)

Drassus alexandrinus O. P.-CAMBRIDGE, 1874; O. P.-Cambridge (1874): 393, pl. 51, fig. 18; **new synonymy**.

Drassodes alexandrinus (O. P.-CAMBRIDGE, 1874); Levy (2004): 16, figs. 40–45.

Material examined: GREECE: Kriti: Moni Vrondisiou (16/4/99–09/6/99, NHMC81.2.10.15 – 2 ♂ 1 ♀); Panagia Almyri (26/1–02/7/00, NHMC81.2.1565.6 – 1 ♂); Gramvousa (26/4–26/6/96, NHMC81.2.10.11 – 1 ♂ 1 ♀); Elafonisi (30/12/96–14/3/97, NHMC81.2.321.4 – 2 ♂ 2 ♀); Achendriás (08/6–04/8/99, NHMC81.2.910.19 – 1 ♂ 4 ♀); Mt. Youchtas (06/6–02/7/96, NHMC81.2.1762.8 – 10 ♂ 4 ♀). Gavdos: Fanari (27/7–10/11/96, NHMC81.2.129.2 – 7 ♂); Vatsiana (14/6–24/8/97, NHMC81.2.474.7 – 4 ♀); Lavrakas sanddunes (26/7–09/11/96, NHMC81.2.132.8 – 1 ♂). Kythira: Mermygaris (10/4/96, – 1 ♂). Antikythira: Potamos, 700 m W (27/3–05/8/01, NHMC81.2.2406 – 3 ♀). Kea: Koundouros (21/4/06, NHMC81.2.8487 – 1 ♀). Naxos: Moutsouna (01/5/06–02/7/06, NHMC81.2.9209 – 2 ♂ 3 ♀). Santorini: Profitis Ileias (17/5–24/8/03, – 44 ♂ 7 ♀). Samos: Psili Ammos (02/5/06–06/7/06, NHMC81.2.9207 – 1 ♀). Rodos: Prasonisi (13/5–08/7/06, NHMC81.2.8467.1 – 5 ♂ 2 ♀). Kos: Kefalos (12/5–23/8/01, NHMC81.2.2223 – 6 ♀). Karpathos: Pyles–Volada (26/6–09/9/01, NHMC81.2.1772 – 1 ♂ 1 ♀). Nisyros: Moni Evangelistrias, road of Palaiokastro (01/5–01/6/05, NHMC81.2.8403.15 – 1 ♀); Nikia to Avlaki – phrygana maquis (01/5–05/6/05, NHMC81.2.8405.19 – 7 ♂); Nikia to Avlaki – phrygana (01/5–05/6/05, NHMC81.2.8406.22 – 1 ♀); Gyali islet (30/4–05/6/05, NHMC81.2.8401.21 – 1 ♀); Kandelioussa islet (03/5–06/6/05, NHMC81.2.8408.10 – 5 ♂ 2 ♀); Pacheia (or Passas) islet (03/5–06/6/05, NHMC81.2.8409.9 – 7 ♂); Pergousa islet (05/5–06/6/05, NHMC81.2.8410.10 – 10 ♂ 1 ♀). Kalymnos: near Emporeios (06/4–09/6/05, NHMC81.2.8419.16 – 3 ♂ 1 ♀); Opposite Agia Aikaterini monastery (06/4–09/6/05, NHMC81.2.8420.16 – 37 ♂ 3 ♀); Stimenia, at the end of the road (06/4–09/6/05, NHMC81.2.8422.11 – 19 ♂ 2 ♀); Pserimos islet, NW part – phrygana (01/4–07/6/05, NHMC81.2.8411.17 – 8 ♂); Pserimos islet, SE part – old cultivations near the beach, phrygana (02/4–07/6/05, NHMC81.2.8412.23 – 1 ♂); Pserimos islet, cultivations (02/4–07/6/05, NHMC81.2.8413.17 – 1 ♂); Plati islet – *Juniperus macrocarpa* (03/4–08/6/05, NHMC81.2.8415.13 – 14 ♂ 5 ♀); Plati islet – *Juniperus oxycedrus* (31/3–08/6/05, NHMC81.2.8416.12 – 8 ♂); Kalavros islet (04/4–10/6/05, NHMC81.2.8424.20 – 72 ♂ 4 ♀); Agia Kyriaki islet (04/4–



Figs. 3–6. *Drassodes lutescens* – male palp. 3. Crete, Elafonisi (left palp), 4. Gavdos islet (left palp), 5. Rodos (left palp), 6. Naxos (right palp). Scale bars = 0.1 cm.

10/6/05, NHMC81.2.8425.14 – 21 ♂ 3 ♀); Telendos islet (05/4–10/6/05, NHMC81.2.8426.16 – 5 ♂ 3 ♀). Astypalaia: Chora to Maltezana, shortly after the narrowest part of the islet (24/4–13/6/05, NHMC81.2.8431.13 – 5 ♂ 3 ♀); Stream after Agios Ioannis church (27/4–13/6/05, NHMC81.2.8437.13 – 3 ♂); Koutsomytis islet (24/4–11/6/05, NHMC81.2.8433.11 – 3 ♂); Kounoupoi islet (24/4–11/6/05, NHMC81.2.8434.13 – 1 ♂); Ofidousa islet (24/4–12/6/05, NHMC81.2.8435.19 – 1 ♂ 1 ♀); Pontikousa islet (26/4–12/6/05, NHMC81.2.8436.10 – 1 ♂).

Remarks: The similarity between *D. lutescens* and *D. alexandrinus* was already stressed by Levy (2004). He illustrated both species and representatives of both sexes with great detail (figs. 33–39 and 40–45 respectively). The main differences between the two species are as follows: in males of *D. alexandrinus* the tibia/cymbium ratio is smaller, the tibial apophysis (if present) is tiny and the filamentous part of the embolus is placed at the lower third of tegulum (in *D. lutescens* it is found at the upper third of tegulum). In females the lateral caps (pouches in Chatzaki & al. 2002b) are larger, enclose the two spermathecae (lower and higher) and reach the epigastric furrow, and the median septum is narrower than that of *D. lutescens*.

Levy also reported that, according to the drawings of Chatzaki & al. (2002b; 613, figs. 23–26), *D. lutescens* from Crete was misidentified and should be placed under the name of *D. alexandrinus*. Based on this remark Bosmans & Chatzaki (2005) doubted the identity of the Greek records of *D. lutescens*.

mS/MS	Loc.	LP/SL	Loc.	EO/Te	Loc.	Cy/Ti	Loc.
0.20	CRT	0.33	SM	0.20	CRT	0.71	GVD
0.20	CRT	0.35	NS	0.25	GVD	0.75	ISR
0.20	CRT	0.44	GVD	0.25	CRT	0.75	CRT
0.20	CRT	0.45	CRT	0.31	ISR	0.76	ISR
0.25	GVD	0.47	KRP	0.33	ISR	0.91	CRT
0.25	CRT	0.50	CRT	0.34	GVD	0.91	NS
0.29	CRT	0.50	CRT	0.35	NS	0.96	GVD
0.29	CRT	0.53	CRT	0.38	CRT	1.08	CRT
0.29	CRT	0.57	NS	0.40	CRT	1.09	CRT
0.33	CRT	0.57	CRT	0.40	CRT	1.10	CRT
0.33	CRT	0.59	CRT	0.41	CRT	1.12	GVDP
0.33	CRT	0.62	NS	0.42	CRT	1.13	CRT
0.33	GVDP	0.63	CRT	0.43	YALI	1.13	CRT
0.33	GVDP	0.63	KEA	0.43	CRT	1.13	CRT
0.33	CRT	0.67	CRT	0.43	CRT	1.15	CRT
0.33	CRT	0.67	CRT	0.44	NX	1.15	CRT
0.40	CRT	0.67	GVDP	0.44	CRT	1.15	CRT
0.40	CRT	0.67	CRT	0.45	CRT	1.16	CRT
0.40	CRT	0.67	CRT	0.47	NS	1.19	CRT
0.43	NS	0.67	NS	0.47	CRT	1.20	CRT
0.43	CRT	0.67	CRT	0.48	GVDP	1.21	CRT
0.50	NS	0.67	CRT	0.48	CRT	1.23	GVD
0.50	YALI	0.67	CRT	0.48	CRT	1.25	CRT
0.50	RD	0.67	RD	0.50	GVD	1.28	YALI
0.50	RD	0.69	SYR	0.50	CRT	1.29	CRT
0.50	CRT	0.71	CRT	0.50	CRT	1.29	CRT
0.50	ISR	0.73	CRT	0.50	CRT	1.33	GVDP
0.50	CRT	0.73	CRT	0.50	CRT	1.33	CRT
0.50	CRT	0.75	RD	0.52	CRT	1.33	CRT
0.60	CRT	0.77	CRT	0.53	CRT	1.34	CRT
0.60	CRT	0.77	GVD	0.56	CRT	1.34	NX
0.60	CRT	0.77	GVD	0.57	GVDP	1.35	NS
0.60	CRT	0.79	CRT	0.58	GVD	1.37	YALI
0.60	CRT	0.79	NX	0.67	YALI	1.43	GVD
0.67	KEA	0.80	CRT				
0.73	SYR	0.83	CRT				
0.75	KRP	0.83	CRT				
0.75	NX	0.83	ISR				
0.80	NS	0.85	CRT				
1.00	CRT	0.86	CRT				
1.00	CRT	0.86	CRT				
1.00	SM	0.91	CRT				
0.43	CRT	0.92	CRT				


EB/Te: 0.44 (0.2–0.67)

Cy/Ti: 1.14 (0.71–1.43)

mS/MS: 0.44 (0.2–1)

SL/LP: 0.67 (0.33–0.92)

Tab. 1. Ratios of morphometric measurements of 34 male and 43 female specimens of *D. lutescens*. The average and range of each ratio are also shown. The following abbreviations are used: a) Embolic origin: EO; Tegulum length: Te; Cymbium length: Cy; Tibia length: Ti; Maximum septum width: MS; Minimum septum width: mS; Lateral pouches length: LP; Spermathecae length: SL, to indicate the morphometric characters measured and b) Greece: CRT – Crete, GVD – Gavdos, GVDP – Gavdopoula, KRP – Karpathos, NS – Nisyros, RD – Rodos, SM – Samos, YALI – Yali, NX – Naxos, KEA – Kea; Israel: ISR; Syria: SYR, to indicate the localities.




Figs. 7–11. *Drassodes lutescens* – male tibial apophysis. 7. Crete, Elafonisi (right palp), 8. Gavdos islet (left palp), 9. Naxos (right palp), 10. Crete, Mt. Youchtas (left palp), 11. Rodos (left palp). Scale bars = 0.1 cm.

and pointed out the need of a re-examination of this material. Vouchers from many Greek islands and from adjacent countries were re-examined or examined for the first time in detail, in order to give an answer to this problem.

Very few individuals from the Greek region strictly follow the distinctive characters of *D. alexandrinus* as illustrated by Levy, except for the dentition of the chelicerae, which is identical in all individuals examined and follows the dentition of *D. alexandrinus*. A great variation of all other characters of the male and female genital organs produces an actual gradient among specimens, and is the reason why the two species are considered to be synonymous.

Male pedipalp (Figs. 3–11, Tab. 1): cymbium/tibia ratio ranges from 0.71 to 1.43, but in most individuals the cymbium is longer than the tibia (27 out of 34 individuals). The size of the tibial apophysis ranges from tiny to quite apparent (Figs. 7–11). However, this distinction does not constantly follow the respective placement of the embolus (tiny in the specimens in which the embolus originates at the lower part of the tegulum and larger in the specimens in which the embolus originates at the higher part of the tegulum). The embolus of most individuals examined is placed at the middle third of the tegulum with few exceptions (5 out of 34 male individuals) in which the embolus is placed at the lower third of the tegulum. As already reported, the shape of



Figs. 12–23. *Drassodes lutescens* – epigyne. 12. Gavdos islet, 13. Crete, Achendrias, 14. Crete, Gramvousa, 15. Crete, Elafonisi, 16. Antikythira, 17. Naxos, 18. Kea, 19. Rodos, 20. Samos, 21. Bulgaria, 22. Syria, 23. Israel. Scale bars = 0.1 cm.

the sperm duct inside the bulb is variable, deviating from a simple upright U-shaped duct (Figs. 4–5) to more complicated sigmoid, laterally directed loops leading to a longer embolus originating from a more basal part of the tegulum (Figs. 3, 6). Shape and size of the membranous lamella are not constant.

Epigyne (Figs. 12–23, Tab. 1): sides of median septum range from almost parallel and wide apart ($mS/MS = 1$, Figs. 15, 17, 22) to conical, being very close at the upper part of the epigyne and very broad at the basal part ($mS/MS = 0,2$, Figs. 16, 18). This feature leads to a variety of shapes. The lateral pouches almost touch the epigastric furrow and/or the lower edges of the median septum in most of cases examined. Their upper part is always situated below the upper rim of the median septum or hardly reaches this height (most extreme in the specimen from Israel, Fig. 23, and the specimen from Naxos,

Fig. 24. *Drassodes lutescens* – vulva.
Astypalaia. Scale bar = 0.1 cm.




Fig. 17, and the specimen from Samos, Fig. 20). The variation presented by the specimens examined is in fact due to the degree of sclerotization of these two structures and the height of the two compartments they form (LP), relative to the height of the two spermathecae of the vulva (SL). At one extreme (*D. lutescens* according to Levy (2004; fig. 38), and Fig. 22 in the present study) the two compartments do not enclose the upper spermathecae ($LP/SL = 0,33$), while at the other extreme (*D. alexandrinus* according to Levy (2004; fig. 44) and Fig. 23 in the present study) the upper spermathecae are fully enclosed in them ($LP/SL = 0,92$). As in males, within the Greek populations there are all the intermediate relative positions of these two extreme forms.

Vulva (Fig. 24): structures of all specimens examined are not different between extreme epigynal forms. The only variation observed is in the relative sizes of the two sets of spermathecae, upper and lower, as illustrated in Figs. 12–23.

In any case, even with the strictest morphological criteria, Greek populations cannot be attributed to one of the two species, as females resemble more *D. lutescens* (lateral pouches covering only part of the upper spermathecae) and males resemble more *D. alexandrinus* (embolus originating from basal half of tegulum). In the morphometric measurements (Tab. 1) the gradient recorded is irrelevant to the geographical location in which specimens were found. In view of the difficulty to clearly separate the two species, the existence of intermediate forms in any of the distinctive characters and the fact that the structures of the genital organs in both sexes are basically the same, it is proposed that *D. alexandrinus* and *D. lutescens* are in reality one species which keeps the name of the earlier created *D. lutescens*.

Distribution: Mediterranean countries, South Urals, Pakistan, Afghanistan, Oman.

***Haplodrassus dalmatensis* (L. KOCH, 1866)**

Drassodes carinatus STRAND, 1906; Strand (1906): 609; **new synonymy**.

Material examined: MNHN, AR 9471, 1 ♀, Ethiopia.

Remarks: The examination of a voucher specimen of the female of *D. carinatus* STRAND, 1906 from the Natural History Museum of Paris (MNHN, AR 9471, Ethiopia, 1 ♂) fully conform to the characters of *H. dalmatensis* without any doubt. Since this voucher comes from a topotypic locality (the type was not available to the authors) a new synonymy is proposed.

***Drassodes albicans* (SIMON, 1878)**

Drassodes oreinos CHATZAKI, 2002; Chatzaki & al. (2002b): 615, figs. 29–31, 38–39, 71; **new synonymy**.

Material examined: Crete, several localities above 1200 m, including type (Chatzaki & al. 2002b) Spain, Sierra Nevada, Veleta (MNHN, AR 9356, 3 ♀); without any locality (MNHN, AR 9362, 4 ♀).

Remarks: *Drassodes albicans* was described by Simon (1878) from southwestern areas of Europe (Spain, France and Corsica). It was further recorded from three areas in Greece by Strand (1917): from the islands Kos and Kriti (=Crete) and from continental Greece (Stereia Ellada, Fokida, Mt. Korax). The description was based on female individuals only. The genitalic characters of this species are identical to those of *D. oreinos* Chatzaki, 2002 (Chatzaki & al. 2002b; 615, figs. 29–31, 38–39, 71), a species recorded only from the high mountains of Crete. Because of this clear preference to mountain habitats, the record of *D. oreinos* from Kos (Strand 1917) – where high elevations are not present – is doubtful. A closer examination of all old specimens is needed in order to decide if there are misidentifications among the older records or not.

Acknowledgements

The study is partly included in the program PYTHAGORAS I "Multilevel approach of biodiversity on four island groups in the East Aegean" and it is co-funded by the European Union (75%) and by the Operational program EPE-AEK II (Education and Initial Vocational Training) of the Ministry of National Education and Religious Affairs. We are deeply grateful to the late Dr. Levy,

and to Dr. Dolanský, and Dr. Řezáč for providing comparative material for this study, as well as to Dr. Rollard for hosting the last author at the Natural History Museum of Paris in April 2003 and for providing comparative material too and any other help and advice. This visit was funded by the European Union and the French government (program COLPARYST).

References

Bosmans, R. & Chatzaki, M. (2005): A catalogue of the spiders of Greece. A critical review of all spider species cited from Greece with their localities. — Arachnological contributions. Newsletter of the Belgian Arachnological Society 20(2, suppl.): 1–124.

Chatzaki, M., Thaler, K. & Mylonas, M. (2002a): Ground spiders (Gnaphosidae; Araneae) of Crete (Greece). Taxonomy and distribution. I. — Revue suisse de Zoologie 109(3): 559–601.

Chatzaki, M., Thaler, K. & Mylonas, M. (2002b): Ground spiders (Gnaphosidae, Araneae) of Crete and adjacent areas of Greece. Taxonomy and distribution. II. — Revue suisse de Zoologie 109(3): 603–633.

Levy, G. (2004): Spiders of the genera *Drassodes* and *Haplodrassus* (Araneae, Gnaphosidae) from Israel. — Israel Journal of Zoology 50: 1–37.

Strand, E. (1917): Arachnologica varia XIX–XX. — Archiv für Naturgeschichte 82: 158–167.

Addresses of the authors:

Eli Chatzopoulou
Natural History Museum of Crete
University of Crete
Irakleio, Greece

Maria Chatzaki (Corresponding author)
Department of Molecular Biology and Genetics
Democritus University of Thrace
University campus, Dragana
GR-68100 Alexandroupolis, Greece

E-mail: maria.chatzaki@gmail.com
mchatzak@mbg.duth.gr

