

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 27 (1953)

Artikel: Generalization of a geometrical theorem of Euler.
Autor: Truesdell, C.
DOI: <https://doi.org/10.5169/seals-21896>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Generalization of a geometrical theorem of Euler

By C. TRUESDELL

Graduate Institute for Applied Mathematics, Indiana University

In his first derivation of the equation of continuity in hydrodynamics¹⁾ Euler calculated the change in area of an infinitesimal triangle and the change in volume of an infinitesimal tetrahedron when their vertices are carried through an infinitesimal motion in time dt . He discards terms $O(dt^2)$ in the displacements, but his subsequent calculation of area and volume is algebraically exact. For the volume change ΔV he obtains

$$\frac{\Delta V}{V} = I_1 dt + I_2 dt^2 + I_3 dt^3 , \quad (1)$$

where I_1, I_2, I_3 are the sum of the one-, two-, and three-rowed principal minors of the velocity gradient matrix and for the area change an analogous formula. Since terms $O(dt^2)$ have been neglected in the displacements, the formula (1) is in general false. Of course, the leading term is all that is required to derive the continuity equation, and the leading term is correct. Euler's subsequent derivations of the continuity equation neglect terms $O(dt^2)$ consistently, as previously had d'Alembert's in special cases, and it is a trivial matter nowadays to construct a rigorous proof making no outward use of infinitesimals at all.

Now usually when an error is found in the work of Euler, after a lapse of time sufficient for it to become remarked habitually in the histories, Euler is found to have been right after all, if once rightly understood. The present error is noticed here for the first time; and, to break with custom, so is the proper reinterpretation. While (1) is indeed false in Euclidean kinematics, it turns out to represent a correct result in the theory of affine motions.

Let x be a vector in n -space, and consider the transformations

$$\bar{x} = (At + I)x , \quad (2)$$

where A is an $n \times n$ matrix whose entries are elements of a field, I is

¹⁾ *Principia motus fluidorum*, Novi Comment. Acad. Sci. Petrop. 6 (1756—1757) 271—311 (1761). This paper was written c. 1752.

the unit matrix, and t is an indeterminate. If we wish to, we may choose to interpret (2) as a continuous family of homogeneous strains.

Introduce the Euclidean definition of volume. That is, define the volume of a parallelopiped whose $n + 1$ vertices are 0 and $x_i, i = 1, 2, \dots, n$, by the formula

$$V \equiv \det x_i . \quad (3)$$

Then follows

$$\begin{aligned} \bar{V} &= \det \bar{x}_i = \det [(At + I)x_i] , \\ &= \det(At + I) \det x_i , \\ &= V \det(At + I) . \end{aligned} \quad (4)$$

By the expansion of the secular determinant follows

$$\frac{\bar{V}}{V} = \sum_{i=0}^n I_i t^i ,$$

where I_i = the sum of the i -rowed principal of A , or, equivalently, the i th elementary symmetric function of the proper numbers of A . It is this result which Euler proved by explicit calculation in the cases $n = 2, 3$. Thus Euler is the discoverer of the secular expansion. From the manner in which he obtained the I_i , it is clear though not actually proved that they are indeed invariants.

From (5) follows

$$\frac{1}{r! V} \frac{d^r \bar{V}}{dt^r} \Big|_{t=0} = \begin{cases} I_r, r = 1, 2, \dots, n, \\ 0, r \geq n+1, \end{cases} \quad (6)$$

yielding a geometrical interpretation for the principal invariants I_r of an arbitrary matrix.

By means of a known theorem²⁾, when t and the entries of A are complex numbers from (6) and (5) we get

$$J_r = \frac{(-1)^r}{(r-1)!} \frac{d^r \log \bar{V}}{dt^r} \Big|_{t=0} , \quad (7)$$

$$\log \frac{\bar{V}}{V} = \sum_{r=1}^{\infty} \frac{(-1)^{r-1} J_r}{r} t^r , \quad (8)$$

where J_r = the sum of the r th powers of the proper numbers of A . It is not difficult to show that the reciprocal of the radius of convergence of (8) is the greatest among the absolute values of the proper numbers of A . When A is Hermitian, several corollaries regarding the signs of the successive derivatives of $\log V$ may be read off from (7).

(Received February 13, 1953)

²⁾ E. g. § 132 of *Burnside and Panton, Theory of Equations*, 2nd ed., Dublin (1886).