

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 27 (1953)

Artikel: On admissibility of sequences and a theorem of Pólya.
Autor: Buck, R. Creighton
DOI: <https://doi.org/10.5169/seals-21887>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On admissibility of sequences and a theorem of Pólya

R. CREIGHTON BUCK

1. Introduction. Let K be the space of entire functions which obey the growth conditions $f(z) = O(e^{Az})$, $f(iy) = O(e^{c|y|})$ for some $A < \infty$ and $c < \pi$. By a theorem of Carlson [7] any such function is completely determined by its values at the positive integers. A sequence $\{a_n\}$ of complex numbers is said to be admissible for the sequence of functionals T_n and the function space C if there exists $f \in C$ such that $T_n(f) = a_n$ for $n = 0, 1, \dots$. [3] For the functionals $T_n(f) = f(n)$ and $C = K$, admissibility is a delicate property; if one term of an admissible sequence is altered the result is inadmissible. More generally, if two sequences agree except for a non-void set of indices of density zero, only one can be admissible. A necessary and sufficient condition for admissibility in this case has been given. [Buck 2, Theorem 2.3.] The present paper deals with the closely related problem of admissibility for the functionals

$T_n^*(f) = \Delta^n f(0)$ and the class K . Since $T_n^* = (-1)^n \sum_{k=0}^n \binom{n}{k} (-1)^k T_k$ and $T_n = \sum_{k=0}^n \binom{n}{k} T_k^*$, a sequence $\{a_n\}$ is admissible for $\{T_n\}$ if and only if

the sequence $b_n = \Delta^n a_0$ is admissible for $\{T_n^*\}$. In replacing $\{T_n\}$ by $\{T_n^*\}$ much is gained. Admissibility no longer depends as much on the precise structure of a sequence but rather on matters of size and angular distribution. For this reason, it is much easier to discuss many questions relative to $\{T_n\}$ admissibility in terms of $\{T_n^*\}$. This approach has been used with success in the characterization problem for integral-valued entire functions. [Buck, 5] In the present paper, we discuss several other applications. In particular, Theorem 2 answers a number of questions raised by the "even difference" theorem of Agnew and Fuchs. The last section, which is somewhat independent, contains a new and extremely brief proof of the classical theorem of Pólya on functions of zero type.

2. Admissibility $\{T_n^*\}$. The first theorem gives a convenient necessary and sufficient condition for the $\{T_n^*\}$ admissibility of a sequence $\{b_n\}$.

Since the proof follows closely the pattern of that for the corresponding theorem for $\{T_n\}$, we omit most of the details. (See [2].)

Theorem 1. Given a complex sequence $\{b_n\}$, let $g(z) = \sum_0^\infty b_n z^n$, Then:

(i) $\{b_n\}$ is admissible $\{T_n^*\}$ if and only if g is analytic at zero and can be continued to the interval $-1 \leq x \leq 0$.

(ii) when $\{b_n\}$ is admissible, the interpolating function f for which $\Delta^n f(0) = b_n$ is given by

$$f(z) = (ML) - \sum_0^\infty \binom{z}{n} b_n \quad (2.1)$$

where $\binom{z}{n} = z(z-1)\cdots(z-n+1)/n!$ and ML denotes Mittag-Leffler summability.

(iii) $g(z)$ is entire if and only if $f(z)$ is of zero type

(iv) if $g(z)$ is a polynomial of degree N , so is $f(z)$, and conversely.

Proof. If $f \in K$ then $g(z) = \sum \Delta^n f(0) z^n$ is given by

$$g(z) = \frac{1}{2\pi i} \int_\Gamma \Phi(w) [1 - (e^w - 1)z]^{-1} dw \quad (2.2)$$

where $\Phi(w)$ is the Borel transform of f and Γ encloses the indicator set $D(f)$ of f . [4] [2]. Let E be the image of the boundary of $D(f)$ under the map $w \rightarrow (e^w - 1)^{-1}$. $g(z)$ is then analytic at zero and may be continued via (2.2) to the component of the complement of E which contains zero; this set in particular contains the interval $[-1, 0]$. If f is of zero type, $D(f)$ is the origin, and g is entire. Conversely, let $g(z) = \sum b_n z^n$ be analytic on $[-1, 0]$ and consider the function $f(z)$ defined by

$$f(z) = \frac{1}{2\pi i} \int_\Gamma t^{-1} g(t) [(1+t)/t]^z dt$$

where Γ is now a path enclosing the interval $[-1, 0]$. Calculation shows that $f \in K$ and that $\Delta^n f(0) = b_n$. Moreover, if g is entire, f is of zero type. Statement (ii) follows from a known theorem concerning the expansion of functions into Newton series [Buck 4, Theorem 4.3] and implies (iv) immediately. As an illustration, all "small" sequences are admissible $\{T_n^*\}$. [18, p. 52, Thm. 10].

Corollary 1. If $\limsup |b_n|^{1/n} < 1$, $\{b_n\}$ is admissible $\{T_n^*\}$.

Any theorem connecting the presence of singular points of a power series with its coefficients may be used to yield characterization theorems for sequences $\{b_n\}$ and in turn for $\{a_n\}$. At this point we insert a

generalization of the familiar theorem concerning power series with positive coefficients. [15, p. 215].

Lemma. Let $\limsup |c_n|^{1/n} = 1/R$. Let $S_n(z) = \sum_0^n c_k z^k$ and suppose that there is a sequence of points z_j approaching a point $\beta = Re^{i\theta}$ from outside the circle $|z| = R$ such that for each j , $S_n(z_j)$ approaches the point at infinity in an angle of opening less than π . Then, β is a singular point for $f(z) = \sum c_n z^n$.

If $f(z)$ is regular at β , it is regular in a circular neighborhood N of β and $(ML) - \lim S_n(z) = f(z)$ for all z in N . But, N contains a point z_j , and since Mittag-Leffler summability is totally regular, $(ML) - \lim S_n(z_j) = \infty$.

We note that Borel summability could have been used in place of Mittag-Leffler, if N is slightly modified; also, the same method yields an analogous result for Dirichlet series and for Laplace transforms. In the classical case, $c_n \geq 0$ so that $\lim S_n(x) = +\infty$ for all $x > R$.

Corollary 2. Let $\{c_n\}$ be a complex sequence with $\limsup |c_n|^{1/n} = 1$, and obeying the condition described in the lemma, with $R = 1$. Then, the sequence $b_n = (-1)^n c_n$ is not admissible $\{T_n^*\}$.

Corollary 3. If $\limsup |b_n|^{1/n} \geq 1$ but $(-1)^n b_n \geq 0$, $\{b_n\}$ is not admissible $\{T_n^*\}$.

For $b_n = \Delta^n a_0$, the oscillation conditions $(-1)^n b_n \geq 0$ and $(-1)^n a_n \geq 0$ are closely related; in fact, the latter implies the former. In particular, we obtain again the following theorem for $\{T_n\}$ admissibility [2, Theorem 4.1].

Corollary 4. If $a_0 \neq 0$ and $(-1)^n a_n \geq 0$, then there is no function $f \in K$ such that $f(n) = a_n$ for $n = 0, 1, \dots$.

Similarly, Corollary 2 could be turned into a somewhat complicated theorem concerned with admissibility $\{T_n\}$. Some growth condition in Corollary 2 is needed since $b_n = (-1)^n 2^{-n}$ is an oscillating sequence, achieved by the function 2^{-z} . The condition given says essentially that infinitely many of the terms b_n are "large", for example, bounded away from zero.

3. Vanishing differences. Agnew [1] proved that if $\{a_n\}$ is a bounded sequence such that $\Delta^n a_0 = 0$ for $n = 0, 2, 4, \dots$, then $a_n = 0$ for all $n = 0, 1, \dots$. Pollard [11] gave a different proof of this and assuming that $a_n = O(n^r)$ proved that $a_n = f(n)$ where $f(z)$ is a polynomial.

Fuchs [6] approached the problem from a different direction and proved the theorem with a weakened assumption on the set of n for which $\Delta^n a_0 = 0$, which in fact was shown to be best possible. We consider the effect of relaxing both this condition and the growth restriction on $\{a_n\}$. We use a simple relation connecting the functions $g(z) = \sum_0^\infty b_k z^k$ and $F(z) = \sum_0^\infty a_n z^n$, namely

$$g(z) = (1 + z)^{-1} F(z/(1 + z)). \quad (3.1)$$

This is easily established, assuming that F is analytic at the origin. (See also [11].)

Theorem 2. Let $\limsup |a_n|^{1/n} \leq 1$ and let $\Delta^n a_0 = 0$ for $n \in A$, a set of integers of density d . If $d > \frac{1}{3}$, then the series $\sum_0^\infty \binom{z}{n} \Delta^n a_0$ converges for all z to a function $f(z)$ of exponential type not exceeding $\log(1 + 2 \cos \pi d)$ such that $f(n) = a_n$ for $n = 0, 1, 2, \dots$. In particular, if $d \geq \frac{1}{2}$, f is of type zero. The value $\frac{1}{3}$ as a lower bound for d is best possible.

Proof. $F(t) = \sum_0^\infty a_n t^n$ is regular for $|t| < 1$ so that by (3.1) $g(z) = \sum_0^\infty \Delta^n a_0 z^n$ is regular in the half plane $x > -\frac{1}{2}$. Let the radius of convergence of this series be R . By Pólya's density theorem for power series [12], every arc of $|z| = R$ of opening $2\pi(1 - d)$ contains a singularity of $g(z)$. Combining these, we see that if $d > \frac{1}{3}$ then $R > 1$, and by Corollary 1, $b_n = \Delta^n a_0$ is admissible $\{T_n^*\}$; it then follows that $\{a_n\}$ is admissible $\{T_n\}$ and is therefore the sequence $\{f(n)\}$ for a unique function $f \in K$. When $d \geq \frac{1}{2}$, R is infinite, $g(z)$ is entire and by Theorem 1, f is of zero type. If $\frac{1}{3} < d < \frac{1}{2}$, $R \geq (2 \cos \pi d)^{-1}$ and further calculation shows that the function $\Phi(w)$ of (2.2) is regular at least for $|w| > \log(1 + R^{-1})$. The type of f does not exceed this value, and in particular is less than $\log 2$, so that the Newton series (2.1) is in fact convergent to $f(z)$ [18, p. 52, Thm. 10]. That the number $1/3$ cannot be improved follows from the fact that the sequence $\{b_n\}$ defined by $\sum_0^\infty b_n z^n = (z - 1)/(z + 1)(z^3 - 1)$ is not admissible $\{T_n^*\}$ so that the corresponding sequence $\{a_n\}$ has vanishing differences of density $\frac{1}{3}$, obeys the growth condition $a_n = o(1)$, but is not admissible $\{T_n\}$.

The effect of the weakened growth condition $\limsup |a_n|^{1/n} \leq 1$ is striking; in contrast with the Agnew-Fuchs result, d may exceed $\frac{1}{2}$ and

may in fact be 1, without $f(z)$ being a polynomial. Witness for example $f(z) = \sum_0^{\infty} \binom{z}{n} (1/n!)$. In this connection, the following more detailed information may be of interest. As we have seen, the region of regularity of $g(z)$ restricts the rate of growth of $f(z)$, and when $g(z)$ is entire, $f(z)$ is of growth at most order 1, type 0. In this case, the rate of growth of g might be expected to impose further restrictions.

Theorem 3. Let $g(z) = \sum_0^{\infty} \Delta^n f(0) z^n$, where $f \in K$. If g is analytic in $|z| < R$ and $R > 1$, f is (at most) of order 1 type $\log(1 + R^{-1})$; if g is entire, and of infinite order, f is of order 1 type 0; if g is of finite order ρ , f is of order $\rho/(1 + \rho)$.

Since the first two statements have already been discussed, we prove only the last. From (3.1) $F(z) = \sum f(n) z^n = (1 - z)^{-1} g(z/(1 - z))$. If $\zeta = 1/(1 - z)$ this may be written as $\zeta g(\zeta - 1)$ which is of order ρ as a function of ζ . By a theorem of Whittaker and Wilson [17] f is of order $\rho/(1 + \rho)$.

For the specific example, $f(z) = \sum \binom{z}{n} (1/n!)$ g is of infinite order and f of zero type. In contrast, $\sum \binom{z}{n} (1/n^2!)$ is of order $\frac{1}{2}$ with d again 1.

4. The Theorem of Pólya. The theorem in question is the following [13]:

Theorem 4. Let $f(z)$ be of order 1 type 0 and suppose that $f(n) = O(1)$ for $n = 0, 1, -1, 2, -2, \dots$. Then, f is constant.

Many proofs of this have been given since it was first proposed. (See Szego [14], Tschakaloff [16], Paley and Wiener [10, p. 81], Levinson [9, p. 127], Korevaar [8]). The following proof is new and has the virtue of extreme simplicity, involving no interpolation series or delicate growth estimates. We make the initial observation, as in [8], that nothing is lost by the assumption that $\sum_{-\infty}^{\infty} |f(n)| < \infty$. Since f is of zero type, $g(z)$ is entire and is given by (2.2). Expanding the kernel $[1 - (e^w - 1)z]^{-1}$, we have

$$\begin{aligned} g(z) &= \frac{1}{2\pi i} \int \Phi(w) (1+z)^{-1} \sum_0^{\infty} e^{nw} [z/(1+z)]^n dw \\ &= (1+z^{-1}) \sum_0^{\infty} f(n) [z/(1+z)]^n \end{aligned}$$

valid for $|z| < |1+z|$. (This is also another verification of (3.1).) From our assumption on $\sum |f(n)|$, $g(z)$ is bounded in the half plane $x > -\frac{1}{2}$. Expanding the kernel in the opposite fashion,

$$\begin{aligned}
g(z) &= \frac{1}{2\pi i} \int \Phi(w) \left(-\frac{1}{z}\right) \sum_0^{\infty} e^{-(n+1)w} [(z+1)/z]^n dw \\
&= \left(-\frac{1}{z}\right) \sum_0^{\infty} f(-n-1) [(1+z)/z]^n
\end{aligned}$$

valid for $|1+z| < |z|$. Again, $g(z)$ is bounded in the half plane $x < -\frac{1}{2}$. Combining these, g and hence f is constant.

REFERENCES

- [1] *R. P. Agnew*, On sequences with vanishing even or odd differences, *Amer. J. Math.* 66 (1944), 339—340.
- [2] *R. C. Buck*, A class of entire functions, *Duke math. J.* 13 (1946), 541—559.
- [3] — Interpolation and uniqueness of entire functions, *Proc. Nat. Acad. Sci.* 33 (1947), 288—292.
- [4] — Interpolation series, *Trans. Amer. Math. Soc.* 64 (1948), 283—298.
- [5] — Integral valued entire functions, *Duke math. J.* 15 (1948), 879—891.
- [6] *W. H. J. Fuchs*, On the closure of $\{e^{-ttdv}\}$, *Proc. Camb. Phil. Soc.* 42 (1946), 91—105.
- [7] *G. H. Hardy*, On two theorems of F. Carlson and S. Wigert, *Acta Math.* 42 (1920), 327—339.
- [8] *J. Korevaar*, A simple proof of a theorem of Pólya, *Simon Stevin*, 26 (1948—49), 72—80.
- [9] *N. Levinson*, Gap and Density Theorems, *Amer. Colloq. Pub.* 26, New York, 1940.
- [10] *R. E. A. C. Paley* and *N. Wiener*, *Fourier Transforms in the Complex Plane*, *Amer. Coll. Pub.* 19, New York, 1934.
- [11] *H. Pollard*, Sequences with vanishing even differences, *Duke math. J.* 12 (1945), 303—304.
- [12] *G. Pólya*, Über Lücken und Singularitäten von Potenzreihen, *Math. Z.* 29 (1929), 549—640.
- [13] — Aufgabe 105, *J.-Ber. Deutsch. Math.-Verein*, 40 (1931), 80.
- [14] *G. Szegő*, Lösung der Aufgabe 105, *J.-Ber. Deutsch. Math.-Verein*, 43 (1934), 10—11.
- [15] *Titchmarsh*, *Theory of Functions*, Oxford, 1932.
- [16] *L. Tschakaloff*, Zweite Lösung der Aufgabe 105, *J.-Ber. Deutsch. Math.-Verein*, 43 (1934), 11—13.
- [17] *J. M. Whittaker* and *R. Wilson*, Fabry's theorem and the isolated singularity of finite exponential order, *J. London Math. Soc.* 14 (1939), 202—208.
- [18] *J. M. Whittaker*, *Interpolatory Function Theory*, Cambridge 1935.

Received 14th July 1952.