Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	30 (1956)
Artikel:	A property of bounded analytic functions.
Autor:	Eggleston, H.G.
DOI:	https://doi.org/10.5169/seals-23907

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# A property of bounded analytic functions

H. G. Eggleston

A well known theorem of Fatou [4] asserts that if f(z) is a bounded regular function defined in |z| < 1, then the radial limit  $\lim_{r \to 1} f(re^{i\theta})$ where r tends to unity through real values less than one exists for all  $\theta$ except for a set of at most zero measure,  $0 \le \theta < 2\pi$ . We denote the

set of all points at which this limit exists by F(f), see [3]. For any point  $\zeta$  of |z| = 1 the cluster set of f(z) at  $\zeta$ ,  $C(f; \zeta)$ , is defined to be the set of all complex numbers w such that there exists a

sequence  $\{z_n\}$  with the three properties

 $|z_n| < 1$   $n = 1, 2, \ldots;$   $z_n \rightarrow \zeta;$   $f(z_n) \rightarrow w$ .

The set of points  $\zeta$  for which  $C(f; \zeta)$  is a single point will be denoted by G(f).

It is trivial that  $F(f) \supset G(f)$ . The object of this note is to show that the set of points in F(f) and not in G(f) is a set of the first category. This result has a similar appearance to the theorem of E. F. Collingwood [3], concerning the sets of points S(f) and I(f).<sup>1</sup>) But both the content and method of proof are different from those of Collingwood.

The connection of this result with Fatou's theorem is that it enables us to disprove a natural conjecture. Since sets of zero measure and sets of first category can to some extent be regarded as interchangeable<sup>2</sup>) one might suppose that the set F(f) would contain all of |z| = 1except a set of first category. In fact not only is this false but it is even true that F(f) itself need only be a set of first category. As concrete examples of such functions f(z) there are,

(A) a Blaschke product

$$f(z) = \prod_{\nu=1}^{\infty} \frac{z e^{-i\theta_{\nu}} - |z_{\nu}|}{1 - z \bar{z}_{\nu}},$$

where  $|z_{\nu}| < 1$ , amp  $z_{\nu} = \theta_{\nu}$ ,  $\sum_{\nu=1}^{\infty} (1 - |z_{\nu}|) < \infty$  and the points  $z_{\nu}$  are

<sup>2</sup>) See [8], p. 77.

<sup>&</sup>lt;sup>1</sup>) See [3], p. 177, for definitions.

so distributed that the closure of their union contains the whole circumference |z| = 1.

(B) a certain function g(z) which maps |z| < 1 onto a bounded simply connected domain whose prime-ends are all of diameter greater than some positive number  $\delta$ . The existence of such functions has been established by F. Frankl [5]. g(z), unlike the Blaschke product of (A), is univalent.

For both the function f(z) of (A) and g(z) of (B) and for every  $\zeta$  of |z| = 1, the cluster set at  $\zeta$  does not reduce to a single point. This is obvious for the function g(z) since the cluster set of g(z) at  $\zeta$  is a prime end of the domain onto which g(z) maps |z| < 1. For the function f(z) of (A), the radial limit exists for almost all  $\theta$  and is of modulus unity (see [7], p. 196). But for any  $\zeta$  there is a subsequence of zeros of f(z) that tends to  $\zeta$ . Thus for almost all  $\zeta$  of |z| = 1,  $C(f; \zeta)$  has diameter greater than or equal to unity. But the set of  $\zeta$  for which the diameter of  $C(f; \zeta)$  is greater than or equal to unity is a closed set, and thus contains the whole of |z| = 1.

We shall refer to a point that belongs to the set F(f) as a Fatou point and a point that belongs to G(f) as a convergence point. We prove next the theorem :

**Theorem.** If a non-constant, regular and bounded function f(z) defined in |z| < 1 is such that for a certain arc  $\gamma$  of |z| = 1, the set F(f) is of second category on every subarc of  $\gamma$ , then all points of  $\gamma$  except at most those of a set of first category belong to G(f).

It is sufficient to show that, given any positive number  $\varepsilon$ , and any subarc  $\gamma_1$  of  $\gamma$ , there is a subarc of  $\gamma_1$  say  $\gamma_2$  such that if  $\zeta$  belongs to  $\gamma_2$ , then the diameter of  $C(f; \zeta)$  is less than  $\varepsilon$ . For suppose that this had been established then it would follow that the set of points  $\zeta$  on |z| = 1 for which the diameter of  $C(f; \zeta)$  is greater than or equal to  $\varepsilon$  would not contain any interval of |z| = 1 in  $\gamma$  and, since this set is clearly closed, it would be non-dense in  $\gamma$ . We allow  $\varepsilon$  to assume a sequence of positive values decreasing to zero and we conclude that the set of points  $\zeta$  for which the diameter of  $C(f; \zeta)$  is positive is a set of first category in  $\gamma$ . This is the statement of the theorem.

Suppose then that  $\gamma_1$  is a given subarc of  $\gamma$  which we shall assume without real loss of generality to be a small arc of |z| = 1. Denote by  $D_1$  the part of |z| < 1 bounded by  $\gamma_1$  and by the linear segment joining the endpoints of  $\gamma$ , and denote by  $E_1 = f(D_1)$  the set of values assumed by f(z) as z varies in  $D_1$ .

If  $E_1$  is of diameter less than  $\varepsilon$  then any subarc of  $\gamma_1$  (strictly interior to  $\gamma_1$ ) will serve as the arc  $\gamma_2$ .

Otherwise select a point  $p_1$  of  $E_1$  and let  $K_1$  and  $L_1$  denote respectively the subsets of  $D_1$  for which

$$|f(z) - p_1| = \epsilon/2$$
 ,  $|f(z) - p_1| = \epsilon/4$  .

Since the values taken by f(z) at the set of points z where f'(z) = 0 form an enumerable set, we can select  $p_1$  such that for every point of  $K_1$  and  $L_1$ ,  $f'(z) \neq 0$ . We also choose  $p_1$  so that its distance from the component of the complement of  $E_1$  which contains the point at infinity, is less than  $\varepsilon/8$ .

Under the above circumstances each of  $K_1$  and  $L_1$  consists of the intersection of |z| < 1 with an at most enumerable set of arcs whose end points belong to |z| = 1.3)

Let  $k_1(r)$  denote the union of all the arcs of  $K_1$  that are completely contained in the annulus,  $1 - r \leq |z| \leq 1$ , and define  $l_1(r)$  similarly. For any arc  $\alpha$  of  $K_1$  let  $p(\alpha)$  be the projection of  $\alpha$  in 0 < |z| < 1from the point z = 0 onto the circumference |z| = 1 by means of the radii arg z =constant.  $p(\alpha)$  is a subinterval of  $\gamma$  which may be open or closed or semi-open. Further write

$$h_1(r) = \bigcup_{\alpha \in k_1(r)} p(\alpha) ; \qquad m_1(r) = \bigcup_{\alpha \in l_1(r)} p(\alpha)$$

Let  $r_i$  be a sequence of positive numbers decreasing to zero and write

$$H_1 = \bigcap_{i=1}^{\infty} h_1(r_i) ; \qquad M_1 = \bigcap_{i=1}^{\infty} m_1(r_i) .$$

Now if  $\zeta \in H_1$  the radius vector joining z = 0 to  $z = \zeta$  intersects  $K_1$  in a sequence of points whose closure contains the point  $\zeta$ ; for if this were not the case, there would be a segment of the radius say  $\zeta_1 \zeta$  disjoint from the set  $K_1$ . Choose the integer *i* so large that  $r_i$  is less than the distance of  $\zeta$  from  $\zeta_1$ . Since  $\zeta \in H_1 \subset h_1(r_i)$  there is an arc  $\alpha$  of  $k_1(r_i)$  that intersects the radius joining z = 0 to  $z = \zeta$ . Moreover by the definition of  $k_1(r_i)$ ,  $\alpha$  must meet segment  $\zeta \zeta_1$  and this is a contradiction which establishes the statement above.

It follows that if  $\zeta$  belongs to  $H_1 \cap M_1$  it does not belong to F(f) i.e.

$$H_1 \cap M_1 \cap F(f) = 0 \quad . \tag{1}$$

Next consider the set  $h_1(r_i)$ . This is the union of an at most enumerable

<sup>&</sup>lt;sup>3</sup>) This follows because  $f'(z) \neq 0$  for  $z \in K_1$  or  $z \in L_1$  and by applications of Koebe's lemma; see [4], [2], p. 96. A similar argument is used in [9], p. 247.

sequence of intervals  $p(\alpha)$  of  $\gamma_1$ . If  $h_1(r_i)$  is dense in  $\gamma_1$ , so is the set obtained from  $h_1(r_i)$  by replacing  $p(\alpha)$  by the same arc without its end points; i. e. if  $h_1(r_i)$  is dense in  $\gamma_1$  it contains an open subset of  $\gamma_1$  also dense in  $\gamma_1$ . Thus finally if each  $h_1(r_i)$ ,  $i = 1, 2, \ldots$  is dense in  $\gamma_1$ , then

$$H_1 = \bigcap_{i=1}^{-} h_1(r_i)$$

contains a  $G_{\delta}$  set dense in  $\gamma_1$ . By a similar argument applied to  $M_1$  we see that either one of the sets  $h_1(r_i)$ ,  $m_1(r_i)$  is not dense in the whole of  $\gamma$  or the set N, the union of the complements of  $H_1$  and  $M_1$  in |z| = 1is of first category on  $\gamma_1$ . But (1) implies  $F(f) \subset N$  and F(f) is by hypothesis of second category; thus this situation cannot arise and at least one of the sets  $h_1(r_i)$ ,  $m_1(r_i)$ ,  $i = 1, 2, \ldots$  is not dense in the whole of  $\gamma_1$ .

Suppose then that the set  $h_1(r_i)$  contains no point of the closed subarc  $\beta$  of  $\gamma_1$ . Let  $\beta_1$  be the arc which is the middle third of the arc  $\beta$ . Denote the circle  $|z| = 1 - r_i$  by T and suppose that  $\beta_1$  is the arc  $\{z \mid z = e^{i\theta}, 0 \leq \theta_1 \leq \theta \leq \theta_2 < 2\pi$ . Let X be the set of points

$$\{z \mid 1 - r_i \leqslant |z| \leqslant 1, \theta_1 \leqslant amp \, z \leqslant \theta_2\}$$

By the definition of  $\beta_1$  any arc of  $K_1$  which meets X must also meet T, for if this were not so, the arc of  $K_1$  concerned would belong to  $k_1(r_i)$ and some point of  $\beta_1$  would belong to  $h_1(r_i)$ . If there were infinitely many such arcs then there would be a point say  $z_0$  of T such that every neighbourhood of  $z_0$  contains points of infinitely many arcs of  $K_1$ . But  $K_1$  is closed and thus  $z_0 \in K_1$ . But this implies that  $f'(z_0) = 0$  and thus there cannot be infinitely many arcs of  $K_1$  with points in every neighbourhood of  $z_0$ . Thus only a finite number of arcs of  $K_1$  meet X.

Hence there is a point of  $\beta_1$ , say  $\zeta_1$ , and a positive number  $\delta$  such that every point of  $K_1$  is distant at least  $\delta$  from  $\zeta_1$ . Select two points  $\zeta', \zeta''$  of |z| = 1 whose distances from  $\zeta_1$  are less than  $\delta$ . Denote the small arc of |z| = 1 whose end points are  $\zeta' \zeta''$  by  $\gamma(1)$  and let D(1) be the part of |z| < 1 bounded by  $\gamma(1)$  and the linear segment joining  $\zeta'$  to  $\zeta''$ .

Since D(1) does not meet  $K_1$  there are two possibilities; (i) the set of values E(1) taken by F(z) in D(1) is contained in the circle whose centre is  $p_1$  and whose radius is  $\varepsilon/2$ , or (ii) the set of values E(1) is exterior to this circle.

If it had been one of the sets  $m_1(r_i)$  that was not dense in the whole of  $\gamma_1$  then we should have been led to similar conclusions except that the  $\epsilon/2$  would be replaced by  $\epsilon/4$ . We can cover both cases without needless repetition by retaining the  $\epsilon/2$  in (i) and replacing it by  $\epsilon/4$  in (ii). If (i) holds we have an arc  $\gamma(1)$  which is a subarc of  $\gamma_1$  and has all the properties that we required of the arc  $\gamma_2$ .

If (ii) holds we repeat the argument. That is to say we find a point  $p_2$  of E(1), an arc  $\gamma(2)$  contained in  $\gamma(1)$  and corresponding sets D(2) and E(2) such that either

(i) E(2) is contained in the circle centre  $p_2$  and radius  $\varepsilon/2$  or

(ii) E(2) is contained in the exterior of the circle centre  $p_2$  and radius  $\epsilon/4$ .

If (i) holds  $\gamma(2)$  will serve as the arc  $\gamma_2$ . Otherwise we repeat the argument.

Since every pair of points of the sequence  $p_1, p_2, \ldots$  are at a distance of at least  $\varepsilon/4$  apart and since all these points are contained in the bounded set of values taken by F(z) in |z| < 1, only a finite number of repetitions of the argument are possible. We are eventually led to case (i) and to an arc with the required properties.

This completes the proof of the theorem.

Remark. To see that the Blaschke product f(z) of (A) and the function g(z) of (B) are such that their Fatou sets F(f) and F(g) are of first category we have only to observe that to any set of second category on |z| = 1, there always corresponds an arc such that the set is of second category at each point of the arc<sup>4</sup>). Now if F(f) (or F(g)) was of second category we could find an interval of |z| = 1 as described above. In this interval the set G(f) (or G(g)) would, by the theorem, be of second category. However this is not so since we know that G(f) and G(g) are void.

## REFERENCES

- S. Banach, Théorème sur les ensembles de première catégorie. Fund. Math. 16 (1930) 395.
- [2] E. F. Collingwood and M. L. Cartwright, Boundary theorems for functions meromorphic in the unit circle. Acta Math. 87 (1952) 83-146.
- [3] E. F. Collingwood, On the linear and angular cluster sets of functions meromorphic in the unit circle. Acta Math. 91 (1954) 165-185.
- [4] P. Fatou, Séries trigonométriques et séries de Taylor. Acta Math. 30 (1906).
- [5] F. Frankl, Zur Primendentheorie. Rec. Math. Moscow 38 (1931) 66-69.
- [6] P. Koebe, Abhandlung zur Theorie der konformen Abbildung I. J. reine angew. Math. 146 (1915) 177-225.
- [7] R. Nevanlinna, Eindeutige analytische Funktionen. Berlin 1936.
- [8] W. Sierpinski, Hypothèse du continu. Warsaw 1934.
- [9] S. Stoilow, Les propriétés topologiques des fonctions analytiques d'une variable. Ann. Inst. Henri Poincaré 2 (1932) 233-266.

(Received Decembre 7, 1954.)

<sup>4</sup>) See [1]. See also Bagemihl and Seidel, Proc. Nat. Acad. Sci. 89 (1953) 1068-1075.