

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 31 (1956-1957)

**Artikel:** On the singularities of DIRICHLET series.  
**Autor:** Tanaka, Chuji  
**DOI:** <https://doi.org/10.5169/seals-515703>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On the singularities of DIRICHLET series

by CHUJI TANAKA, Tokyo

**1. Introduction.** Let us put

$$F(s) = \sum_{n=1}^{\infty} a_n \exp(-\lambda_n s) \quad (s = \sigma + it, \quad 0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_n \rightarrow +\infty). \quad (1.1)$$

The object of this note is to establish the following theorems :

**Theorem 1.** *Let (1.1) have the finite simple convergence-abscissa  $\sigma_s$ . Then there exists a sequence  $\{\varepsilon_n\}$  ( $\varepsilon_n = \pm 1$ ) such that*

$$\sum_{n=1}^{\infty} \varepsilon_n a_n \exp(-\lambda_n s)$$

*has  $\sigma = \sigma_s$  as the natural boundary.*

**Theorem 2.** *Let (1.1) have the finite simple convergence-abscissa  $\sigma_s$ . Then there exists a new DIRICHLET series  $\sum_{n=1}^{\infty} b_n \exp(-\lambda_n s)$  having  $\sigma = \sigma_s$  as the natural boundary such that*

$$|b_n| = |a_n| \quad (n = 1, 2, \dots), \quad \lim_{n \rightarrow \infty} |\arg(b_n) - \arg(a_n)| = 0 \quad (i)$$

or

$$\arg(b_n) = \arg(a_n) \quad (n = 1, 2, \dots), \quad \lim_{n \rightarrow \infty} |b_n/a_n| = 1. \quad (ii)$$

Under the additional conditions  $\lim_{n \rightarrow \infty} \log n / \lambda_n = 0$ , these two theorems have been proved by O. SZASZ ([1], p. 107) and the author ([2], p. 308) respectively. The method of its proofs is based upon A. OSTROWSKI's criterion of singularities.

**2. Lemmas.** To establish these theorems, we need some lemmas.

**Lemma 1.** (A. OSTROWSKI [3, 4], pp. 12–16.) *Let (1.1) have the simple convergence-abscissa  $\sigma_s = 0$ . For  $s = 0$  to be the singular point of (1.1), it is necessary and sufficient that we have*

$$\overline{\lim}_{m \rightarrow \infty} |O_m(\sigma, \omega)|^{1/m} \geq 1 \quad (m : \text{positive integer}),$$

where

$$\sigma (> 0), \quad \omega (0 < \omega < 1) : \text{fixed constants}, \quad (i)$$

$$O_m(\sigma, \omega) = \sum_{m/\sigma \cdot (1-\omega) \leq \lambda_n \leq m/\sigma \cdot (1+\omega)} a_n (\lambda_n e \sigma / m)^m \exp(-\lambda_n \sigma). \quad (ii)$$

**Lemma 2.** *The simple convergence-abscissa  $\sigma_s$  of (1.1) is determined by*

$$\sigma_s = \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} a_n \right| ,$$

where

$$[x]: \text{ GAUSS's symbol,} \quad (\text{i})$$

$$f(x) = 1/m \cdot [m(x - m)] , \quad m = [x] . \quad (\text{ii})$$

**Proof.** By T. KOJIMA's theorem [5],  $\sigma_s$  is given by

$$\begin{aligned} \sigma_s &= \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x] \leq \lambda_n < x} a_n \right| \\ &= \overline{\lim}_{x \rightarrow +\infty} 1/[x] \cdot \log \left| \sum_{[x] \leq \lambda_n < x} a_n \right| . \end{aligned}$$

Hence, for any given  $\varepsilon (> 0)$ ,

$$\left| \sum_{[x] \leq \lambda_n < x} a_n \right| < \exp \{[x](\sigma_s + \varepsilon)\} \quad \text{for } [x] > K(\varepsilon) .$$

Therefore

$$\begin{aligned} \left| \sum_{[x]+f(x) \leq \lambda_n < x} a_n \right| &= \left| \sum_{[x] \leq \lambda_n < x} - \sum_{[x] \leq \lambda_n < x+f(x)} a_n \right| \\ &< 2 \exp \{[x](\sigma_s + \varepsilon)\} \quad \text{for } [x] > K(\varepsilon) , \end{aligned}$$

so that

$$\sigma_s^* = \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} a_n \right| \leq \sigma_s + \varepsilon .$$

Letting  $\varepsilon \rightarrow 0$ ,

$$\sigma_s^* \leq \sigma_s . \quad (2.1)$$

Putting

$$\begin{cases} x = m + y/m & (m = [x], \quad 0 < y < m) , \\ [x] + f(x) = m + [y]/m , \end{cases}$$

we have easily

$$\sigma_s^* = \overline{\lim}_{\substack{m \rightarrow +\infty \\ 0 < y < m}} 1/m \cdot \log \left| \sum_{m+[y]/m \leq \lambda_n < m+y/m} a_n \right| \quad (m: \text{ positive integer}).$$

Hence, for any given  $\varepsilon (> 0)$ ,

$$\left| \sum_{\substack{m+[y]/m \leq \lambda_n < m+y/m \\ 0 < y < m}} a_n \right| < \exp \{m(\sigma_s^* + \varepsilon)\} \quad \text{for } m > K^*(\varepsilon) .$$

Accordingly

$$\begin{aligned} &\left| \sum_{[x] \leq \lambda_n < x} a_n \right| \\ &= \left| \sum_{m \leq \lambda_n < m+1/m} + \sum_{m+1/m \leq \lambda_n < m+2/m} + \cdots + \sum_{m+[y]/m \leq \lambda_n < m+y/m} \right| \\ &\leq ([y] + 1) \exp \{m(\sigma_s^* + \varepsilon)\} \leq m \exp \{m(\sigma_s^* + \varepsilon)\} \quad \text{for } m > K^*(\varepsilon) , \end{aligned}$$

so that

$$\sigma_s = \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x] \leq \lambda_n < x} a_n \right| \leq \sigma_s^* + \varepsilon .$$

Letting  $\varepsilon \rightarrow 0$ ,

$$\sigma_s \leq \sigma_s^* . \quad (2.2)$$

Combining (2.1) with (2.2), lemma 2 is completely established.

**Lemma 3.** Put

$$O(x, t) = \sum_{[x] + f(x) \leq \lambda_n < x} a_n (\lambda_n e/[x])^{[x]} \exp(-\lambda_n(1+it)) .$$

If

$$\left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| \geq \left| \sum_{[x] + f(x) \leq \lambda_n < X} a_n \right|$$

for every  $X$  such that  $[x] + f(x) < X \leq x$ , then

$$|O(x, t)| \geq \left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| \cdot \{1 - (|t| + K(x))/[x]\} ,$$

where  $K(x) = O(1)$ .

**Proof.** Put

$$\begin{cases} x = m + y/m \quad (m = [x], \quad 0 < y < m) , \\ [x] + f(x) = m + [y]/m . \end{cases}$$

Let us denote by  $\{\lambda_{m,i}\}$  ( $i = 1, 2, \dots, r$ ) the exponents  $\lambda'_n$ s contained in  $m + [y]/m \leq \lambda_n < m + y/m$  and by  $\{a_{m,i}\}$  the coefficient corresponding to  $\lambda_{m,i}$ . Setting

$$S_k = \sum_{i=1}^k a_{m,i} \quad (1 \leq k \leq r), \quad f(\lambda) = \lambda^m \exp(-\lambda(1+it)) ,$$

then by ABEL's transformation

$$\begin{aligned} O(x, t) &= (e/m)^m \cdot \sum_{i=1}^r a_{m,i} f(\lambda_{m,i}) \\ &= (e/m)^m \cdot \{S_r f(\lambda_{m,r}) + \sum_{i=1}^{r-1} S_i (f(\lambda_{m,i}) - f(\lambda_{m,i+1}))\} . \end{aligned}$$

Since

$$|S_r| \geq |S_i| \quad (i = 1, 2, \dots, r-1) ,$$

we have

$$|O(x, t)| \leq (e/m)^m \cdot |S_r| \cdot \{ |f(\lambda_{m,r})| - \sum_{i=1}^{r-1} |f(\lambda_{m,i}) - f(\lambda_{m,i+1})| \} . \quad (2.3)$$

On the other hand,  $|f(\lambda)| = \lambda^m \exp(-\lambda)$  is monotone-decreasing for  $m \leq \lambda$ . Hence, for  $m \leq \alpha < \beta < m+1$ ,

$$|f(\alpha) - f(\beta)| \leq |f(\alpha)| - |f(\beta)| + |f(\beta)|(\beta - \alpha)|t| ,$$

so that

$$|f(\alpha) - f(\beta)| < \int_{\alpha}^{\beta} z^m \exp(-z) \cdot (|t| + 1 - m/z) dz . \quad (2.4)$$

Therefore, by (2.4)

$$\begin{aligned} \sum_{i=1}^{r-1} |f(\lambda_{m,i}) - f(\lambda_{m,i+1})| &< \int_{\lambda_{m,1}}^{\lambda_{m,r}} z^m \exp(-z) \cdot (|t| + 1 - m/z) dz \\ &< (|t| + 1) \cdot \int_{m+[y]/m}^{m+([y]+1)/m} z^m \exp(-z) dz . \end{aligned}$$

Hence, by (2.3) and  $[y] + 1 \leq m$

$$\begin{aligned} & |O(x,t)| \\ & \geq |S_r| (e/m)^m \{(m + ([y] + 1)/m)^m \cdot \exp(-(m + ([y] + 1)/m)) \\ & \quad - (|t| + 1) \cdot \int_{m+[y]/m}^{m+([y]+1)/m} z^m \exp(-z) dz\} \\ & > |S_r| \{(1 + ([y] + 1)/m^2)^m \cdot \exp(-([y] + 1)/m) - (|t| + 1) \cdot 1/m\} \\ & > |S_r| \{\exp(-0(1/m)) - (|t| + 1) \cdot 1/m\} = |S_r| \cdot \{1 - (|t| + 0(1))/m\} , \end{aligned}$$

which proves lemma 3.

**Lemma 4.** *Let (1.1) have the simple convergence-abscissa  $\sigma_s = 0$ . Then there exists a sequence  $\{x_\nu\}$  ( $[x_\nu] \uparrow +\infty$ ) independent of  $t$  ( $-\infty < t < +\infty$ ) such that*

$$\left. \begin{aligned} (a) \quad [x_\nu](1 + \omega) &< [x_{\nu+1}](1 - \omega) \quad (\nu = 1, 2, \dots, 0 < \omega < 1) , \\ (b) \quad \lim_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| &= 0 , \\ (c) \quad \liminf_{\nu \rightarrow \infty} |O(x_\nu, t)|^{1/[x_\nu]} &\leq 1 \quad \text{for arbitrary } t (-\infty < t < +\infty) . \end{aligned} \right\}$$

**Proof.** By lemma 2, we can find a sequence  $\{y_\nu\}$  ( $[y_\nu] \uparrow +\infty$ ) such that

$$\left. \begin{aligned} [y_\nu](1 + \omega) &< [y_{\nu+1}](1 - \omega) \quad (\nu = 1, 2, \dots, 0 < \omega < 1) , \\ \lim_{\nu \rightarrow \infty} 1/y_\nu \cdot \log \left| \sum_{[y_\nu] + f(y_\nu) \leq \lambda_n < y_\nu} a_n \right| &= 0 . \end{aligned} \right\} \quad (2.5)$$

Put

$$\max_{[y_\nu] + f(y_\nu) \leq x \leq y_\nu} \left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| = \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| ,$$

where  $[x_\nu] = [y_\nu]$ ,  $f(x_\nu) = f(y_\nu)$ . Since

$$\begin{aligned} 0 = \lim_{\nu \rightarrow \infty} 1/y \cdot \log \left| \sum_{[y_\nu] + f(y_\nu) \leq \lambda_n < y_\nu} a_n \right| &\leq \overline{\lim}_{\nu \rightarrow \infty} 1/x \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| \\ &\leq \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| = 0 , \end{aligned}$$

by (2.5) and (2.6), selecting a suitable subsequence, if necessary, we have easily

$$\left. \begin{aligned} & [x_\nu](1 + \omega) < [x_{\nu+1}](1 - \omega) \quad (\nu = 1, 2, \dots, 0 < \omega < 1) , \\ & \lim_{\nu \rightarrow \infty} 1/x_\nu \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| = 0 , \\ & \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| \geq \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < X} a_n \right| \\ & \text{for every } X \text{ such that } [x_\nu] + f(x_\nu) < X \leq x_\nu . \end{aligned} \right\} \quad (2.7)$$

On account of (2.7) and lemma 3

$$\left. \begin{aligned} & [x_\nu](1 + \omega) < [x_{\nu+1}](1 - \omega) \quad (\nu = 1, 2, \dots, 0 < \omega < 1) , \\ & \lim_{\nu \rightarrow \infty} 1/x_\nu \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| = 0 , \\ & \lim_{\nu \rightarrow \infty} |O(x_\nu, t)|^{1/[x_\nu]} \geq \lim_{\nu \rightarrow \infty} \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right|^{1/[x_\nu]} = 1 \\ & \text{for arbitrary } t \ (-\infty < t < +\infty) , \end{aligned} \right\}$$

which is to be proved.

**Lemma 5.** *Let (1.1) have the simple convergence-abscisse  $\sigma_s = 0$ . Then, for any sequence  $\{x_\nu\}$  ( $[x_\nu] \uparrow +\infty$ ),  $\sum_{\nu=1}^{+\infty} u_\nu(s)$  converges absolutely for  $\sigma > 0$ , where*

$$u_\nu(s) = \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \exp(-\lambda_n s) .$$

**Proof.** By lemma 2

$$0 = \overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| = \overline{\lim}_{x \rightarrow +\infty} 1/[x] \cdot \log \left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| .$$

Hence, for any given  $\varepsilon (> 0)$

$$\left| \sum_{[x] + f(x) \leq \lambda_n < x} a_n \right| < \exp(\varepsilon[x]) \quad \text{for } [x] > X(\varepsilon) . \quad (2.8)$$

Using the same notations as in lemma 3, by ABEL's transformation

$$\begin{aligned} u_\nu(s) &= \sum_{i=1}^{r_\nu} a_{m_\nu, i} \exp(-\lambda_{m_\nu, i} s) \\ &= S_{r_\nu} \exp(-\lambda_{m_\nu, r_\nu} s) \\ &\quad + \sum_{i=1}^{r_\nu-1} S_i \{ \exp(-\lambda_{m_\nu, i} s) - \exp(-\lambda_{m_\nu, i+1} s) \} , \end{aligned}$$

where

$$x_\nu = m_\nu + y_\nu/m_\nu \quad (m_\nu = [x_\nu] , \quad 0 < y_\nu < m_\nu) ,$$

$$\text{Since} \quad [x_\nu] + f(x_\nu) = m_\nu + [y_\nu]/m_\nu .$$

$$|\exp(-\alpha s) - \exp(-\beta s)| \leq |s|/\sigma \cdot \{\exp(-\alpha\sigma) - \exp(-\beta\sigma)\}$$

for  $\alpha < \beta$ ,  $\sigma > 0$ , by (2.8)

$$\begin{aligned} |u_\nu(s)| &< |s|/\sigma \cdot \exp(\varepsilon[x_\nu]) \cdot \{\exp(-\lambda_{m_\nu, r_\nu}\sigma) \\ &+ \sum_{i=1}^{r_\nu-1} \exp(-\lambda_{m_\nu, i}\sigma) - \exp(-\lambda_{m_\nu, i+1}\sigma)\} \\ &\leq |s|/\sigma \cdot \exp([x_\nu](\varepsilon - \sigma)) \quad \text{for } \sigma > 0, \quad [x_\nu] > X(\varepsilon) . \end{aligned}$$

Therefore, putting  $\varepsilon = \sigma/2$

$$|u_\nu(s)| \leq |s|/\sigma \cdot \exp(-\sigma/2 \cdot [x_\nu]) ,$$

so that  $\sum_{\nu=1}^{+\infty} u_\nu(s)$  is absolutely convergent for  $\sigma > 0$ . q. e. d.

**Lemma 6.** Let (1.1) have the simple convergence-abscissa  $\sigma_s = 0$ .

(A) Let us put

$$\sum_{\nu=1}^{+\infty} \delta(m_\nu) u_\nu(s) = \sum_{n=1}^{+\infty} \varepsilon_n a_n \exp(-\lambda_n s) , \quad (2.9)$$

where

$$(a) \lim_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| = 0 \quad ([x_\nu] = m_\nu \uparrow +\infty)$$

$$(b) \lim_{x \rightarrow +\infty} 1/x \cdot \log |\delta(x)| = 0 ,$$

$$(c) u_\nu(s) = \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \exp(-\lambda_n s)$$

$$(d) \begin{cases} \varepsilon_n = \delta(m_\nu) & \text{for } \lambda_n \in \{I_\nu\} \\ \varepsilon_n = 0 & \text{for } \lambda_n \notin \{I_\nu\} \end{cases} \quad (I_\nu : [x_\nu] + f(x_\nu) \leq x < x_\nu)$$

Then (2.9) has the simple convergence-abscissa  $\sigma = 0$ .

(B) Set

$$f_0(s) + \sum_{\nu=1}^{+\infty} \delta(m_\nu) u_\nu(s) = \sum_{n=1}^{+\infty} \varepsilon'_n a_n \exp(-\lambda_n s) , \quad (2.10)$$

where

$$(e) f_0(s) = \sum_{\lambda_n \notin \{I_\nu\}} a_n \exp(-\lambda_n s)$$

$$(f) \begin{cases} \varepsilon'_n = \delta(m_\nu) & \text{for } \lambda_n \in \{I_\nu\} \\ \varepsilon'_n = 1 & \text{for } \lambda_n \notin \{I_\nu\} \end{cases} .$$

Then (2.10) has also the simple convergence-abscissa  $\sigma = 0$ .

---

<sup>1)</sup> By lemma 1, there exists a sequence  $\{x_\nu\}$  satisfying (a).

**Proof.** (A) By lemma 2

$$\overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} a_n \right| = 0 . \quad (2.11)$$

Taking account of (a), (b), (c) and (d), we get

$$\begin{aligned} \overline{\lim}_{\substack{x \rightarrow +\infty \\ [x] \neq m_\nu}} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon_n a_n \right| &= -\infty , \\ \overline{\lim}_{\substack{x \rightarrow +\infty \\ [x] = m_\nu \\ x_\nu \leq x < m_\nu + 1}} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon_n a_n \right| &= \overline{\lim}_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \{ |\delta(m_\nu)| \cdot \left| \sum_{m_\nu + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| \} , \\ &= 0 , \\ \overline{\lim}_{\substack{x \rightarrow +\infty \\ [x] = m_\nu \\ m_\nu + f(x_\nu) < x < x_\nu}} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon_n a_n \right| &= \overline{\lim}_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \{ |\delta(m_\nu)| \cdot \left| \sum_{[x_\nu + f(x_\nu)] + f(x) \leq \lambda_n < x} a_n \right| \} , \\ &\leq 0 \quad (\text{by (2.11)}) , \\ \overline{\lim}_{\substack{x \rightarrow +\infty \\ [x] = m_\nu \\ m_\nu \leq x \leq m_\nu + f(x_\nu)}} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon_n a_n \right| &= -\infty , \end{aligned}$$

so that

$$\overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon_n a_n \right| = 0 ,$$

which proves that the simple convergence-abscissa of (2.9) is  $\sigma = 0$ .

(B) Since  $f_0(s) = f(s) - \sum_{\nu=1}^{+\infty} u_\nu(s)$ , by lemma 5  $f_0(s)$  is simply convergent at least for  $\sigma > 0$ . On account of (A),  $\sum_{\nu=1}^{+\infty} \delta(m_\nu) \cdot u_\nu(s) = \sum_{n=1}^{+\infty} \varepsilon_n a_n \exp(-\lambda_n s)$  is simply convergent exactly for  $\sigma > 0$ , so that (2.10) is simply convergent at least for  $\sigma > 0$ . In other words, by lemma 2

$$\overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon'_n a_n \right| \leq 0 .$$

On the other hand, by (a), (b), and (f)

$$\begin{aligned} \overline{\lim}_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} \varepsilon'_n a_n \right| &= \overline{\lim}_{\nu \rightarrow +\infty} 1/x_\nu \cdot \log \{ |\delta(m_\nu)| \cdot \left| \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \right| \} . \\ &= 0 . \end{aligned}$$

Therefore

$$\overline{\lim}_{x \rightarrow +\infty} 1/x \cdot \log \left| \sum_{[x]+f(x) \leq \lambda_n < x} \varepsilon'_n a_n \right| = 0 ,$$

which proves that (2.10) has also the simple convergence-abscissa  $\sigma = 0$ . q. e. d.

### 3. Proofs of theorems

**Proof of theorem 1.** Without any loss of generality, we can assume  $\sigma_s = 0$ . Let us denote by  $\{x_\nu\}$  the sequence of lemma 4. Then, by lemma 5, we can put

$$\sum_{\nu=1}^{+\infty} u_\nu(s) = f_1(s) + f_2(s) + f_3(s) \dots + f_n(s) + \dots ,$$

where

$$(i) \quad u_\nu(s) = \sum_{[x_\nu] + f(x_\nu) \leq \lambda_n < x_\nu} a_n \exp(-\lambda_n s) ,$$

(ii) each  $f_n(s)$  ( $n = 1, 2, \dots$ ) contains infinite number of  $\{u_\nu(s)\}$ .

For any given sequence  $\{\delta_n\}$  ( $\delta_n = \pm 1$ ), set

$$\begin{aligned} f(s : \{\delta_n\}) &= f_0(s) + \delta_1 f_1(s) + \dots + \delta_n f_n(s) + \dots \\ &= f_0(s) + \sum_{\nu=1}^{+\infty} \alpha_\nu u_\nu(s) , \\ &= \sum_{n=1}^{+\infty} \varepsilon'_n s_n \exp(-\lambda_n s) , \end{aligned} \tag{3.1}$$

where

$$(i) \quad f_0(s) = \sum_{\lambda_n \in \{I_\nu\}} a_n \exp(-\lambda_n s) \quad (I_\nu : [x_\nu] + f(x_\nu) \leq x < x_\nu) ,$$

$$(ii) \quad \alpha_\nu = \pm 1 ,$$

$$\begin{aligned} (iii) \quad \varepsilon'_n &= \alpha_\nu \quad \text{for } \lambda_n \in \{I_\nu\} , \\ \varepsilon'_n &= 1 \quad \text{for } \lambda_n \notin \{I_\nu\} . \end{aligned}$$

Then, by lemma 6 (B), (3.1) has the simple convergence-abscissa  $\sigma = 0$ .

For  $\{\delta_n\} \neq \{\delta'_n\}$  ( $\delta_n, \delta'_n = \pm 1$ ), we have

$$\begin{aligned} f(s : \{\delta_n\}) - f(s : \{\delta'_n\}) &= \sum_{n=1}^{+\infty} \delta''_n f_n(s) \\ &= \sum_{\nu=1}^{+\infty} \pm 2 \cdot u_\nu^*(s) \\ &= \sum_{n=1}^{+\infty} \varepsilon_n a_n \exp(-\lambda_n s) , \end{aligned} \tag{3.2}$$

where

$$(i) \quad \delta''_n = \delta_n - \delta'_n \quad (\text{Since } \{\delta_n\} \neq \{\delta'_n\}, \text{ there exists at least one } n \text{ such that } \delta''_n = \pm 2) ,$$

$$(ii) \quad u_\nu^*(r) = \sum_{[x'_\nu] + f(x'_\nu) \leq \lambda_n < x'_\nu} a_n \exp(-\lambda_n s) \quad (\{x'_\nu\} : \text{subsequence of } \{x_\nu\}) ,$$

$$(iii) \quad \begin{aligned} \varepsilon_n &= \pm 2 \quad \text{for } \lambda_n \in \{I'_\nu\} \quad (I'_\nu : [x'_\nu] + f(x'_\nu) \leq x < x'_\nu) \\ \varepsilon_n &= 0 \quad \text{for } \lambda_n \notin \{I'_\nu\} . \end{aligned}$$

On account of lemma 4 (b) and lemma 6 (A), (3.2) has also the simple convergence-abscissa  $\sigma = 0$ .

(3.2) has  $\sigma = 0$  as the natural boundary. To prove this, by lemma 1, it is sufficient to show that

$$\overline{\lim}_{m \rightarrow +\infty} |O_m(1, \omega; t)|^{1/m} \geq 1 \quad \text{for arbitrary } t \quad (-\infty < t < +\infty), \quad (3.3)$$

where

$$O_m(1, \omega; t) = \sum_{m(1-\omega) \leq \lambda_n \leq m(1+\omega)} \varepsilon_n a_n (\lambda_n e/m)^m \cdot \exp(-\lambda_n(1+it)) .$$

Since  $O_{[x', \nu]}(1, \omega; t) = \pm 2O(x'_\nu, t)$  by lemma 4 (a), we have by lemma 4 (c)

$$\begin{aligned} \overline{\lim}_{m \rightarrow \infty} |O_m(1, \omega; t)|^{1/m} &\geq \overline{\lim}_{\nu \rightarrow +\infty} |O_{[x', \nu]}(1, \omega; t)|^{1/[x', \nu]} \\ &= \overline{\lim}_{\nu \rightarrow +\infty} |O(x'_\nu, t)|^{1/[x', \nu]} \\ &\geq \overline{\lim}_{\nu \rightarrow +\infty} |O(x_\nu, t)|^{1/[x_\nu]} \geq 1 , \end{aligned}$$

which proves (3.3).

Let us denote by  $E(\{\delta_n\})$  the set of regular points on  $\sigma = 0$  of (3.1), which is evidently an open set. Then we have

$$E(\{\delta_n\}) \cap E(\{\delta'_n\}) = 0 \quad \text{for} \quad \{\delta_n\} \not\equiv \{\delta'_n\} . \quad (3.4)$$

In fact, if  $s_0 \in E(\{\delta_n\}) \cap E(\{\delta'_n\}) \neq 0$ , (3.2), would be regular at  $s = s_0$ , which is impossible by (3.3). If  $E(\{\delta_n\}) \neq 0$  for all  $\{\delta_n\}$ , by (3.4) the set of all functions  $\{f(s : \{\delta_n\})\}$  is at most enumerable, which contradicts the power of continuum of the set of all  $\{\delta_n\}$ . Hence  $E(\{\delta_n\}) = 0$  for at least one  $\{\delta_n\}$ , which shows that  $f(s : \{\delta_n\})$  has  $\sigma = 0$  as the natural boundary.

(3.1) is evidently of the form  $\sum_{n=1}^{+\infty} \varepsilon'_n a_n \exp(-\lambda_n s)$  ( $\varepsilon'_n = \pm 1$ ). q. e. d.

**Proof of theorem 2.** Without any loss of generality, we can assume  $\sigma_s = 0$ . Let us denote by  $\{x_\nu\}$  the sequence of lemma 4. Let us put

$$f(s, \theta, \alpha) = f_0(s) + f_r(s, \theta, \alpha) = \sum_{n=1}^{\infty} \varepsilon'_n a_n \exp(-\lambda_n s) , \quad (3.5)$$

where

$$(i) \quad f_0(s) = \sum_{\lambda_n \in \{I_\nu\}} a_n \exp(-\lambda_n s) \quad (I_\nu : [x_\nu] + f(x_\nu) \leq x < x_\nu) ,$$

$$(ii) \quad f_1(s, \theta, \alpha) = \sum_{\nu=1}^{+\infty} \exp(\alpha \theta / [x_\nu]) \cdot u_\nu(s) ,$$

$$u_\nu(s) = \sum_{\lambda_n \in I_\nu} a_n \exp(-\lambda_n s) ,$$

$\theta$ : a real constant,  $\alpha$ : a parameter ( $= \sqrt{-1}$  or 1),

- (iii)  $\varepsilon'_n = \exp(\alpha\theta/[x_\nu]) \quad \text{for } \lambda_n \in \{I_\nu\} ,$   
 $\varepsilon'_n = 1 \quad \text{for } \lambda_n \notin \{I_\nu\} .$

Then, by lemma 6 (B), (3.5) has the simple convergence-abscissa  $\sigma = 0$ . Put

$$\begin{aligned} & f(s, \theta, \alpha) - f(s, \theta', \alpha) \\ &= \sum_{\nu=1}^{+\infty} \{\exp(\alpha\theta/[x_\nu]) - \exp(\alpha\theta'/[x_\nu])\} \cdot u_\nu(s) \\ &= \sum_{n=1}^{+\infty} \varepsilon_n a_n \exp(-\lambda_n s) \quad \text{for } \theta \neq \theta' , \end{aligned} \tag{3.6}$$

where

$$\begin{aligned} \varepsilon_n &= 0 \quad \text{for } \lambda_n \notin \{I_\nu\} , \\ \varepsilon_n &= \exp(\alpha\theta/[x_\nu]) - \exp(\alpha\theta'/[x_\nu]) \\ &= \alpha(\theta - \theta')/[x_\nu] + O(1/x_\nu^2) \quad \text{for } \lambda_n \in \{I_\nu\} . \end{aligned}$$

Then, by lemma 4 (b) and lemma 6 (A), (3.6) has also the simple convergence-abscissa  $\sigma = 0$ .

(3.6) has  $\sigma = 0$  as the natural boundary. To establish this fact, by lemma 1, it is sufficient to show that

$$\overline{\lim}_{m \rightarrow \infty} |O_m(1, \omega, t)|^{1/m} \geq 1 \quad \text{for arbitrary } t \quad (-\infty < t < +\infty) . \tag{3.7}$$

Since

$$O_{[x_\nu]}(1, \omega, t) = \{\alpha(\theta - \theta')/[x_\nu] + O(1/x_\nu^2)\} \cdot O(x_\nu, t) ,$$

by lemma 4 (c)

$$\begin{aligned} \overline{\lim}_{m \rightarrow \infty} |O_m(1, \omega, t)|^{1/m} &\geq \overline{\lim}_{\nu \rightarrow \infty} |O_{[x_\nu]}(1, \omega, t)|^{1/[x_\nu]} \\ &= \overline{\lim}_{\nu \rightarrow \infty} |O(x_\nu, t)|^{1/[x_\nu]} \\ &\geq \underline{\lim}_{\nu \rightarrow \infty} |O(x_\nu, t)|^{1/[x_\nu]} \geq 1 , \end{aligned}$$

from which follows (3.7).

Let us denote by  $E(\theta, \alpha)$  the set of regular points on  $\sigma = 0$  of (3.5), which is an open set. Then we have

$$E(\theta, \alpha) \cap E(\theta', \alpha) = 0 \quad \text{for } \theta \neq \theta' . \tag{3.8}$$

In fact, if  $s_0 \in E(\theta, \alpha) \cap E(\theta', \alpha) \neq 0$  (3.6), would be regular at  $s = s_0$ , which is impossible by (3.7). If  $E(\theta, \alpha) \neq 0$  for all  $\theta$  ( $0 < \theta < \gamma$ ,  $\gamma$ : a fixed constant), by (3.8) the set of all functions  $\{f(s, \theta, \alpha)\}$  is at most enumerable, which

contradicts the power of continuum of the set of  $\theta$ . Hence  $E(\theta, \alpha) = 0$  for at least one  $\theta$ , which shows that  $f(s, \theta, \alpha)$  has  $\sigma = 0$  as the natural boundary. If  $\alpha = \sqrt{-1}$  (or = 1), (i) (or (ii)) of theorem 2 is established. q. e. d.

## REFERENCES

- [1] O. SZASZ, *Über Singularitäten von Potenzreihen und Dirichletschen Reihen am Rande des Konvergenzbereiches*. Math. Ann. 85 (1922).
- [2] C. TANAKA, *Note on Dirichlet series (IV). On the singularities of Dirichlet series*. Proc. Amer. Math. Soc. 4 (1953).
- [3] A. OSTROWSKI, *Über das Hadamardsche Singularitätskriterium in der Theorie der Taylorschen und Dirichletschen Reihe*. Sitzungsber. Berl. Math. Ges. Bd. 27 (1928).
- [4] V. BERNSTEIN, *Leçons sur les progrès récents de la théorie des séries de Dirichlet*. Paris 1933.
- [5] T. KOJIMA, *On the convergence-abscissa of general Dirichlet series*, Tôhoku Math. Journ. 6 (1914-15).

(Received May 9, 1956)