

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 32 (1957-1958)

**Artikel:** On subharmonic functions and differential geometry in the large.  
**Autor:** Huber, Alfred  
**DOI:** <https://doi.org/10.5169/seals-25335>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On subharmonic functions and differential geometry in the large \*)

by ALFRED HUBER, Basle and Zurich

## 1. Introduction

We consider an *open, two-dimensional RIEMANNIAN manifold*  $M$  whose metric is defined by a positive definite quadratic form

$$ds^2 = E(\xi, \eta) d\xi^2 + 2F(\xi, \eta) d\xi d\eta + G(\xi, \eta) d\eta^2, \quad (1.1)$$

$\xi$  and  $\eta$  denoting local parameters. If  $E$ ,  $F$  and  $G$  are sufficiently regular, then it is possible to introduce (local) *isothermic parameters*, i. e. there exists a coordinate transformation  $x = x(\xi, \eta)$ ,  $y = y(\xi, \eta)$  such that  $E = G > 0$ ,  $F = 0$  in the  $(x, y)$ -parameter system. Then we can write

$$ds^2 = e^{2u(x, y)} (dx^2 + dy^2) = e^{2u(z)} |dz|^2, \quad (1.2)$$

putting  $z = x + iy$ . Such a transformation always exists, for example, when  $E$ ,  $F$  and  $G$  are of class  $C^3$ , and in this case the corresponding function  $u$  is also of class  $C^3$  (cf. A. WINTNER [34, p. 687]).

By the Theorema egregium the GAUSSIAN curvature  $K$  can be calculated from the  $E$ ,  $F$ ,  $G$  and their partial derivatives up to the second order. In the isothermic parameter system (1.2) one obtains the particularly simple expression

$$K = -e^{-2u} \Delta u \quad (\Delta \equiv \partial^2/\partial x^2 + \partial^2/\partial y^2). \quad (1.3)$$

Hence, letting  $dA = e^{2u} dx dy$  denote the area element on  $M$  we have

$$K dA = -\Delta u dx dy. \quad (1.4)$$

Furthermore, one finds after some calculation (using e. g. [6, p. 175]) the following expression for the *geodesic curvature*  $k$  of a curve on  $M$

$$k = e^{-u} \left( k_e + \frac{\partial u}{\partial n} \right). \quad (1.5)$$

Here  $k_e$  denotes the euclidean curvature of the corresponding curve  $z = z(t)$  in the  $z$ -plane with the convention that  $\text{sign } k_e \equiv \text{sign } \frac{d}{dt} \left[ \arg \frac{dz}{dt} \right]$ , and  $n$

---

\*) This research was supported by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command under contract No. AF 18 (600)-573 and carried out at the University of Maryland.

designates the normal to  $z(t)$  in the direction  $\arg\left(-i \frac{dz}{dt}\right)$ . (1.2) and (1.5) imply

$$kds = \left(k_e + \frac{\partial u}{\partial n}\right) |dz|. \quad (1.6)$$

In general, isothermic parameters can only be introduced in the small. In order to be able to treat problems pertaining to differential geometry in the large we have to consider the RIEMANN surface  $S$  which is determined by the conformal structure of  $M$ . (For a detailed discussion of this step the reader is referred to [31, pp. 2–5]. At this point we have to introduce the additional condition that  $M$  is *orientable*. However, if this should not be the case we simply replace  $M$  by an orientable, two-sheeted covering surface [33, p. 61]). The local uniformizers are then defined as functions which map a portion of  $M$  conformally onto a region in the  $z$ -plane. Hence their real and imaginary parts form a set of local isothermic parameters. Conversely, if  $x$  and  $y$  are local isothermic parameters, then either  $x + iy$  or  $y + ix$  constitutes a local uniformizer.

We thus are led to conceive of  $M$  as a RIEMANN surface on which a *conformal metric*

$$ds = e^{u(z)} |dz| \quad (z = \text{local uniformizer}) \quad (1.7)$$

has been introduced. Thereby a change of uniformizers  $z = \varphi(\zeta)$  implies the transformation

$$\tilde{u}(\zeta) = u(\varphi(\zeta)) + \log|\varphi'(\zeta)|, \quad (1.8)$$

due to the conformal invariance of  $ds = e^{u(z)} |dz| = e^{\tilde{u}(\zeta)} |d\zeta|$ .

We shall mainly be concerned with the relation between the surface integral of the GAUSSIAN curvature (curvatura integra) and the topological and conformal structures of complete, open, two-dimensional RIEMANNIAN manifolds. (According to the definition of H. HOPF and W. RINOW [17] the manifold  $M$  is called *complete* if every divergent path on  $M$  has infinite length. A path  $s$  is said to be *divergent* (or to *tend to the ideal boundary* of  $M$ ) if (1)  $s$  is the topological image  $p = p(t)$  of the half-open interval  $0 \leq t < 1$ , (2) given an arbitrary subcompact  $K$  of  $M$  there always exists a number  $t'(K) < 1$  such that  $p(t)$  lies outside  $K$  for  $t > t'$ ).

The present article originated from a suggestion of Professor H. HOPF. He drew our attention to the connection between differential geometry and potential theory which is revealed by relations (1.3) and (1.4). For example, the function  $u(x, y)$  is subharmonic in a certain  $(x, y)$ -parameter region if and only if  $K \leq 0$  in the corresponding domain on  $M$ . (This fact had already been used by E. F. BECKENBACH and T. RADÓ [3] in their proof of the isoperi-

metric inequality on surfaces of negative curvature.) Analogously,  $u$  is superharmonic if and only if  $K \geq 0$ . Furthermore, (1.4) discloses an even deeper connection: The surface integral of  $K$ , considered as a set function, is essentially the measure associated with  $u$  (i. e. the mass distribution of density  $\Delta u/2\pi$ , which appears when  $u$  is represented as a sum of a logarithmic potential and a harmonic function). Consequently, results of differential geometry in the large involving the curvatura integra, such as those due to S. COHN-VOSSEN [10], F. FIALA [12], CH. BLANC and F. FIALA [5] (see H. HOPF [16] for further references), have a potentialtheoretical meaning. It is therefore natural to apply functiontheoretical methods to this field in the hope that not only other (and eventually simpler) proofs of known results will be found, but also theorems which are new in both their differentialgeometrical and potentialtheoretical aspects. From this viewpoint [18], [19] and the present paper have been written.

Our geometrical results are contained in sections 4, 5 and 6. We consider manifolds which are given in the form (1.7), assuming merely that  $u$  can be represented as a difference of subharmonic functions<sup>1)</sup>,

$$u(z) = u_1(z) - u_2(z) \quad (z = \text{local uniformizer}). \quad (1.9)$$

Of course, neither the functions  $u_1$  and  $u_2$  nor their associated measures  $\mu_1$  and  $\mu_2$  are uniquely determined. However, the difference  $\mu = \mu_1 - \mu_2$  does not depend on the choice of decomposition or uniformizer. Consequently,  $\mu$  is defined as a measure on  $S$ . Let  $\mu = \mu^+ - \mu^-$  denote the JORDAN decomposition [30, p. 11] of  $\mu$ , which can be characterized by the property that

$$\mu^+(e) \leq \mu_1(e) \quad \text{and} \quad \mu^-(e) \leq \mu_2(e)$$

for all BOREL sets  $e$  and any representation  $\mu = \mu_1 - \mu_2$  of  $\mu$  as a difference of positive measures. Further, let  $C^+ = 2\pi\mu^-(S)$  and  $C^- = 2\pi\mu^+(S)$ . The difference  $C = C^+ - C^- = -2\pi\mu(S)$ , defined whenever  $C^+$  and  $C^-$  are not both infinite, will be called the *curvatura integra* of the metric. This terminology is justified, since for sufficiently regular  $u$  we have indeed, by (1.4),

$$C^+ = \iint_S \Delta^- u dx dy = \iint_M K^+ dA ,$$

$$C^- = \iint_S \Delta^+ u dx dy = \iint_M K^- dA$$

and

$$C = - \iint_S \Delta u dx dy = \iint_M K dA ,$$

---

<sup>1)</sup> Such a representation always exists, for example, when  $u$  is of class  $C^2$ . For the definition and properties of subharmonic functions the reader may consult the book of T. RADÓ [25].

where  $\Delta^+u = \max[\Delta u, 0]$ ,  $\Delta^-u = \max[-\Delta u, 0]$ ,  $K^+ = \max[K, 0]$  and  $K^- = \max[-K, 0]$ .

From the viewpoint of potential theory (1.9) is the most natural condition to impose on  $u$ . (We mention that such a metric has already been considered by A. BEURLING [4]. He restricted himself to the case of negative curvature, i. e. subharmonic  $u$ .) The following remark may illustrate that this generality is also useful for differentialgeometrical purposes: The theory presented here applies to all RIEMANNIAN manifolds whose coefficients  $E$ ,  $F$  and  $G$  in (1.1) are of class  $C^1$  and which possess a continuous GAUSSIAN curvature in the sense of H. WEYL [32, pp. 43–44]. This is a consequence of results due to S. S. CHERN, P. HARTMAN and A. WINTNER [9] who demonstrated that under these hypotheses isothermic parameters can be introduced, the corresponding function  $u$  being of class  $C^1$ . Furthermore, one deduces easily from [9] that  $u$  is representable in the form (1.9) and that the curvatura integra  $C$  (in the above definition) is equal to the surface integral of the GAUSSIAN curvature.

An interesting special case of the metric (1.7) is given by the modulus of an analytic differential (cf. R. NEVANLINNA [24, p. 103]),  $ds = |dw| = |\varphi(z)| |dz|$ . Thereby we allow  $dw$  to be multiple-valued as long as  $|dw|$  is single-valued. Furthermore, we admit isolated singularities  $a_k$  in whose neighborhoods  $\varphi$  is representable in the form  $\varphi(z) = (z - a_k)^{p_k} \Psi(z)$ , where  $p_k$  denotes an arbitrary real number and  $\Psi$  is a function regular at  $a_k$  ( $k = 1, 2, 3, \dots$ ). The  $p_k$ 's are conformal invariants, and we have  $C = -2\pi \sum p_k$ , this quantity being defined whenever  $C^+ = 2\pi \sum \max(-p_k, 0)$  and  $C^- = 2\pi \sum \max(p_k, 0)$  are not both infinite.

Throughout section 4 we suppose that  $S$  is finitely connected. Theorem 10 states that  $C \leq 2\pi\chi$  for any complete metric (1.7) whose curvatura integra exists,  $\chi$  denoting the EULER-POINCARÉ characteristic of  $S$ . This result has already been proved by S. COHN-VOSSEN (Satz 6, p. 79 in [10]) under more restricted regularity conditions. (He admitted manifolds whose metrics were defined by positive definite quadratic forms (1.1), the coefficients  $E$ ,  $F$  and  $G$  being of class  $C^2$ . In this case the GAUSSIAN curvature is defined and continuous. Hence, by a previous remark, our theorems can be applied.) Our proof is different from the one given by COHN-VOSSEN, although the central idea of this author has rather been transformed than altogether eliminated. We believe that our reasoning is simpler, at least if one disregards the complications needed for getting rid of unnatural regularity assumptions. It is function-theoretical. No use is made of the theorem of H. HOPF and W. RINOW [17] which states that on a (sufficiently regular) complete manifold any two points can be joined by a geodesic whose length is equal to their distance.

The remainder of section 4 is devoted to sufficient conditions for equality,

$C = 2\pi\chi$ . One of these results (Theorem 11) implies a statement due to S. COHN-VOSSEN (Satz 7, p. 79 in [10]).

In section 5 infinitely connected, complete manifolds are investigated. Finally, in section 6 theorems of F. FIALA [12] and of CH. BLANC and F. FIALA [5] are extended.

Theorems 10 to 14, as formulated, apply only to orientable manifolds. However, it is easy to consider also the non-orientable case. The completeness of the metric is not destroyed if we pass to a two-sheeted, orientable covering manifold. Furthermore, in this process *curvatura integra*, EULER-POINCARÉ characteristic, total area and length of closed curves are all multiplied by 2. Consequently, an application of the above mentioned theorems to the covering manifold yields immediately the corresponding results in the non-orientable case. Hence in these theorems the metric (1.7) need not necessarily be defined on a RIEMANN surface. It is sufficient to suppose that  $S$  is a generalized RIEMANN surface in the sense that  $S$  is defined like a RIEMANN surface (cf. R. NEVANLINNA [24, p. 53]), but that both directly and indirectly conformal neighborhood relations are admitted.

In section 2 some theorems on conformal metrics defined in doubly connected, *schlicht* regions are demonstrated. These results are needed for subsequent applications, but they are also of interest by themselves. In particular statements concerning integrals of moduli of analytic functions along certain curves are implied. Such integrals have been the object of previous investigations – we mention the work of L. FEJÉR and F. RIESZ [11], R. M. GABRIEL [13], F. CARLSON [8], M. RIESZ [29] and B. ANDERSSON [1] – but, to our knowledge, problems of the type treated here have not been considered.

Section 3 contains a special result (Theorem 7) whose possible generalizations<sup>2)</sup> might warrant further investigation.

The reader is assumed to be familiar with some properties of subharmonic functions (cf. T. RADÓ [25]), in particular the theory of F. RIESZ [27].

We express our sincere gratitude to Professor H. HOPF for suggesting the problem. We are very much indebted to Professor A. PFLUGER for encouragement.

## 2. Some theorems on conformal metrics defined in *schlicht* annular regions

In the following let  $\Omega$  be a doubly connected region in the  $z$ -plane which does not contain the point at infinity in its interior. We denote by  $\Gamma$  and  $\Delta$  the outer and inner boundaries of  $\Omega$ , respectively. (We make no regularity

---

<sup>2)</sup> Such as the subharmonic analogue or the extension from the plane to more general RIEMANN surfaces. See also the remarks to Theorem 7.

assumptions about  $\Gamma$  and  $\Delta$ . In particular we allow  $\Gamma$  to consist of only the point at infinity.) Let  $\Omega_0$  designate the interior region of  $\Gamma$ .

Let  $u(z)$  be a function defined in  $\Omega$  and locally representable as a difference of two subharmonic functions

$$u(z) = u_1(z) - u_2(z) . \quad (2.1)$$

$u(z)$  can assume the values  $+\infty$  ( $u_1$  finite,  $u_2 = -\infty$ ) and  $-\infty$  ( $u_1 = -\infty$ ,  $u_2$  finite). It may be left undefined at those points where  $u_1 = u_2 = -\infty$ . This point set is of no concern to us since it is a set of measure zero with respect to all occurring integrations.

In a well known way (F. RIESZ [27]) measures  $\mu_1(e)$  and  $\mu_2(e)$  are associated with the functions  $u_1(z)$  and  $u_2(z)$ , respectively. We define

$$\mu(e) = \mu_1(e) - \mu_2(e) . \quad (2.2)$$

If, for a given function  $u(z)$ , there exists one decomposition of the form (2.1), then there are an infinite number. Of course, the corresponding measures  $\mu_1$  and  $\mu_2$  depend on the choice of decomposition. However, the difference  $\mu$  is the same for every such representation. Furthermore, for every prescribed decomposition (2.2) there exists a corresponding representation (2.1). If (2.2) is the JORDAN decomposition [30, p. 11] of  $\mu$ , then (2.1) is called canonical. We may assume without loss of generality that (2.1) is canonical and valid throughout  $\Omega$  (cf. M. G. ARSOVE [2, p. 331]).

Let  $\gamma$  be a JORDAN curve in  $\Omega$  which encloses  $\Delta$ . We introduce the *flux* of  $u$  through  $\gamma$  in accordance with the theory of F. RIESZ [27]. If  $u$  is of class  $C^2$  and if  $\gamma$  is analytic, then we simply define

$$\Phi(u, \gamma; \Gamma) = \frac{1}{2\pi} \int_{\gamma} \frac{\partial u}{\partial n} |dz| , \quad (2.3)$$

$n$  denoting the outer normal. In the case of general  $u$  and  $\gamma$  we introduce a sequence  $\{\delta_k\}$  of JORDAN curves such that the annular regions  $(\Delta, \delta_k)$ , bounded by  $\Delta$  and  $\delta_k$ , tend increasingly to  $(\Delta, \gamma)$  as  $k \rightarrow \infty$ . Let

$$h_k(z) = h_{1k}(z) - h_{2k}(z) ,$$

where  $h_{1k}$  and  $h_{2k}$  are the best harmonic majorants of  $u_1$  and  $u_2$ , respectively, in  $(\delta_k, \gamma)$ .  $h_k$  is independent of the choice of the decomposition (2.1). Let now  $\delta'_k$  denote an analytic JORDAN curve in  $(\delta_k, \gamma)$  which encloses  $\delta_k$ . Then  $\Phi(h_k, \delta'_k; \gamma)$  is well defined and its value is the same for every such  $\delta'_k$ , since  $h_k$  is harmonic. We define

$$\Phi(u, \gamma; \Gamma) = \lim_{k \rightarrow \infty} \Phi(h_k, \delta'_k; \gamma) . \quad (2.4)$$

F. RIESZ [27] proved that if  $u$  is subharmonic then this limit always exists, being finite and independent of the choice of  $\{\delta_k\}$ . It is easy to conclude from this that the same is true in our case.

Now, let  $\{\gamma_l\}$ ,  $l = 1, 2, 3, \dots$ , be an arbitrary sequence of JORDAN curves, enclosing  $\Delta$ , whose interior regions tend increasingly to  $\Omega_0$ . In Theorems 1, 2 and 3 (below) we make the hypothesis

(A) *For any such sequence the limit*

$$\Phi(\Gamma) = \lim_{l \rightarrow \infty} \Phi(u, \gamma_l; \Gamma) \quad (2.5)$$

*exists, admitting the values  $+\infty$  and  $-\infty$ .* Of course,  $\Phi(\Gamma)$  is necessarily independent of the sequence  $\{\gamma_l\}$ .

The theory of F. RIESZ implies that

$$\Phi(u_1, \gamma_{l+1}; \Gamma) - \Phi(u_1, \gamma_l; \Gamma) = \mu_1[\gamma_l \cup (\gamma_l, \gamma_{l+1})]$$

for all  $l$ . Hence the sequence  $\{\Phi(u_1, \gamma_l; \Gamma)\}$  is non-decreasing. The same is true for  $\{\Phi(u_2, \gamma_l; \Gamma)\}$ . Consequently, the limits

$$\Phi_1(\Gamma) = \lim_{l \rightarrow \infty} \Phi(u_1, \gamma_l; \Gamma)$$

and

$$\Phi_2(\Gamma) = \lim_{l \rightarrow \infty} \Phi(u_2, \gamma_l; \Gamma)$$

always exist, being finite or  $+\infty$ .

*Hypothesis (A) is equivalent to the assumption:*

(B)  $\Phi_1(\Gamma)$  and  $\Phi_2(\Gamma)$  are not both infinite. Furthermore

$$\Phi(\Gamma) = \Phi_1(\Gamma) - \Phi_2(\Gamma) . \quad (2.6)$$

Let us briefly indicate a proof of this statement. If (B) is fulfilled, then (A) and (2.6) follow immediately from the relation

$$\Phi(u, \gamma_l; \Gamma) = \Phi(u_1, \gamma_l; \Gamma) - \Phi(u_2, \gamma_l; \Gamma) \quad (l = 1, 2, 3, \dots) ,$$

which is an obvious consequence of the definition of  $\Phi$ . The second half of the equivalence proof has to be based on the fact that  $\mu = \mu_1 - \mu_2$  is a JORDAN decomposition. (We have supposed that the representation (2.1) is canonical.) Without entering into details we mention that the assumption  $\Phi_1(\Gamma) = \Phi_2(\Gamma) = +\infty$  makes it possible to construct two sequences  $\{\gamma_l\}$  and  $\{\gamma'_l\}$  of the above mentioned type for which  $\Phi(\Gamma) = +\infty$  and  $\Phi(\Gamma) = -\infty$ , respectively. Clearly, this yields a contradiction to (A).

A path  $\sigma$  in  $\Omega$  will be said to *tend to*  $\Gamma$  if the following conditions are fulfilled:

(1)  $\sigma$  is the topological image  $z = z(t)$  of the half-open interval  $0 \leq t < 1$ ;

(2) given an arbitrary subcompact  $K$  of  $\Omega_0$  there exists a number  $t'(K) < 1$ , such that  $z(t)$  lies outside  $K$  for  $t > t'$ .

**Theorem 1.** *If  $\Phi(\Gamma) < -1$ , then there exists a locally rectifiable path  $\sigma$  in  $\Omega$ , tending to  $\Gamma$ , such that*

$$\int_{\sigma} e^{u(z)} |dz| < +\infty. \quad (2.7)$$

**Remarks.** There is no value of  $\Phi(\Gamma)$  outside the above mentioned interval for which the theorem is also true. This can be seen from the following counterexamples:

- (I)  $\Omega_0 =$  finite  $z$ -plane,  $u_1 = |z|^2$ ,  $u_2 \equiv 0$ . Then  $\Phi(\Gamma) = +\infty$ .
- (II)  $\Omega_0 =$  finite  $z$ -plane,  $u_1 = \alpha \log |z|$ ,  $u_2 \equiv 0$ . Then  $\Phi(\Gamma) = \alpha$  ( $0 \leq \alpha < +\infty$ ).
- (III)  $\Omega_0 =$  finite  $z$ -plane,  $u_1 \equiv 0$ ,  $u_2 = -\alpha \log |z|$ . Then  $\Phi(\Gamma) = \alpha$  ( $-1 \leq \alpha \leq 0$ ).
- (IV) Let  $\Gamma$  contain at least two points. Then there exists a conformal mapping  $w = \varphi(z)$  of  $\Omega_0$  onto  $|w| < 1$ . We define

$$u_1 = \frac{1}{1 - |\varphi(z)|} + \log |\varphi'(z)|, \quad u_2 \equiv 0.$$

Then  $\Phi(\Gamma) = +\infty$ .

The reader will easily verify that in each of these examples there exists no path  $\sigma$  having the properties postulated in Theorem 1. (It should be observed that the choice of  $\Delta$  is irrelevant.)

It is natural to ask whether the hypothesis  $\Phi(\Gamma) < -1$  can be weakened if more restrictive conditions are imposed on  $\Omega$ . Examples (I), (II) and (III) show that the condition  $\Phi(\Gamma) < -1$  cannot be replaced by a weaker inequality for those domains  $\Omega$  whose boundary component  $\Gamma$  consists of only the point at infinity. We shall now demonstrate that for all other regions the hypothesis  $\Phi(\Gamma) < -1$  in Theorem 1 can be replaced by  $\Phi(\Gamma) < +\infty$ . Since, on the other hand, for the case  $\Phi(\Gamma) = +\infty$  we have given counterexamples for any  $\Omega$ , this settles the question completely.

**Theorem 2.** *If  $\Gamma$  contains more than one point, and if  $\Phi(\Gamma) < +\infty$ , then there exists a locally rectifiable path  $\sigma$ , tending to  $\Gamma$ , such that (2.7) is fulfilled.*

**Proof of Theorem 2.** From (2.6) and the hypothesis  $\Phi(\Gamma) < +\infty$  we infer that  $\Phi_1(\Gamma) < +\infty$ . We choose an arbitrary positive number  $\alpha > \Phi_1(\Gamma) + 1$  and consider the function  $u^* = u_1^* - u_2^*$ , where  $u_1^*(z) \equiv u_1(z)$  and  $u_2^*(z) = u_2(z) - \alpha g_0(z, z_0)$ . Here  $g_0$  denotes GREEN's function of  $\Phi_0$  and  $z_0$  is an ar-

bitrary but fixed point in this region. Throughout  $\Omega$ ,  $u < u^*$ . The functions  $u_1^*$  and  $u_2^*$  are obviously subharmonic. Because of the choice of  $\alpha$ ,  $\Phi^*(\Gamma) < -1$ . Therefore, Theorem 1 may be applied to  $u^*(z)$ .<sup>3)</sup> There exists a locally rectifiable path  $\sigma$ , tending to  $\Gamma$ , such that

$$\int_{\sigma} e^u |dz| < \int_{\sigma} e^{u^*} |dz| < +\infty.$$

This completes the proof of Theorem 2.

It is also possible to weaken the hypothesis  $\Phi(\Gamma) < -1$  in Theorem 1 by making further assumptions about the function  $u$ . We shall now discuss the effect of a condition which is natural from the point of view of both the theory of functions and the applications to differential geometry.

A sequence of curves  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , will be said to *come arbitrarily near to  $\Gamma$* , if the point set  $\bigcup_n \gamma_n$  is not contained in any subcompact of  $\Omega_0$ .

**Theorem 3.** *Suppose there exists a sequence  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , of rectifiable JORDAN curves, enclosing  $\Delta$ , in  $\Omega$  and a number  $M$  such that*

(a)  $\{\gamma_n\}$  comes arbitrarily near to  $\Gamma$ ,

(b)  $\int_{\gamma_n} e^u |dz| < M$  for all  $n$ .

Then, if  $\Phi(\Gamma) \neq -1$ , there exists a locally rectifiable path  $\sigma$ , tending to  $\Gamma$ , such that (2.7) is fulfilled.

**Remarks.** The hypothesis  $\Phi(\Gamma) \neq -1$  cannot be dropped. This follows from the example

$$\begin{aligned} \text{(V)} \quad \Omega_0 &= \text{finite } z\text{-plane, } u_1 = 0, \quad u_2 = \log |z|, \quad \gamma_n = [|z| = n], \\ \Phi(\Gamma) &= -1. \end{aligned}$$

By Theorem 2 no such counterexamples exist if  $\Omega_0$  is of hyperbolic type. In this case Theorem 3 actually gives new information only for  $\Phi(\Gamma) = +\infty$ .

If  $\Phi(\Gamma) < -1$ , then Theorem 3 is, of course, superseded by Theorem 1 for any  $\Omega$ .

We are left to prove Theorems 1 and 3.

**Preliminary considerations.** A point  $z_0$  in  $\Omega$  will be called a *singular point of the measure  $\mu$*  if  $\mu(z_0) \geq 1$ . (The symbol  $z_0$  is used here to denote the set consisting of the point  $z_0$ .) In every subcompact of  $\Omega$  there are at most a finite number of such points.

**Lemma 1.** *Let  $\alpha$  be an analytic arc which contains no singular point of  $\mu$ . Then*  

$$\int_{\alpha} e^u |dz| < +\infty.$$

---

<sup>3)</sup> The proof of Theorem 1 will later be given.

**Remark.** It is easy to construct examples which show that in this lemma the word “analytic” cannot be replaced by “rectifiable”.

**Proof.** Let  $\tilde{\alpha}$  denote the segment  $0 \leq \xi \leq 1$  on the real axis of a complex  $\zeta$ -plane ( $\zeta = \xi + i\eta$ ). There exists a conformal mapping  $z = \varphi(\zeta)$  of a neighborhood  $\tilde{V}$  of  $\tilde{\alpha}$  onto a neighborhood  $V$  of  $\alpha$  such that  $\tilde{\alpha}$  corresponds to  $\alpha$ . We now consider in  $\tilde{V}$  the subharmonic functions

$$\begin{aligned}\tilde{u}_1(\zeta) &\equiv u_1(\varphi(\zeta)) + \log|\varphi'(\zeta)| \\ \tilde{u}_2(\zeta) &\equiv u_2(\varphi(\zeta))\end{aligned}\tag{2.8}$$

and define  $\tilde{u} = \tilde{u}_1 - \tilde{u}_2$ , so that  $e^u|dz| = e^{\tilde{u}}|d\zeta|$ . One proves, without difficulty, that  $\mu_1(e) = \tilde{\mu}_1(\tilde{e})$  and  $\mu_2(e) = \tilde{\mu}_2(\tilde{e})$  for corresponding sets  $e$  and  $\tilde{e}$ ,  $\tilde{\mu}_1$  and  $\tilde{\mu}_2$  denoting the measures associated with  $\tilde{u}_1$  and  $\tilde{u}_2$ , respectively. Because of the existence of such a transformation we may, without loss of generality, assume  $\alpha$  to be the segment  $0 \leq x \leq 1$  on the real axis.

Given an arbitrary point  $x_0$  on  $\alpha$ , there always exists a radius  $r_0(x_0) > 0$  such that  $\mu_2(|z - x_0| < 2r_0) = p < 1$ . By a well known theorem of F. RIESZ [27, II, p. 350] we have the representations

$$u_1(z) = h_1(z) + \int_{|\zeta - x_0| < 2r_0} \log|z - \zeta| d\mu_1(e_\zeta) \tag{2.9}$$

and

$$u_2(z) = h_2(z) + \int_{|\zeta - x_0| < 2r_0} \log|z - \zeta| d\mu_2(e_\zeta) \tag{2.10}$$

in  $|z - x_0| < 2r_0$ , the functions  $h_1$  and  $h_2$  being harmonic. An obvious covering argument yields the existence of  $\int_0^1 e^u dx$  if we can show that

$$\int_{x_0 - r_0}^{x_0 + r_0} e^u dx < +\infty.$$

But, by (2.9) and (2.10), this will be achieved if we can prove that

$$I = \int_{x_0 - r_0}^{x_0 + r_0} \exp \left\{ - \int_{|\zeta - x_0| < 2r_0} \log|x - \zeta| d\mu_2(e_\zeta) \right\} dx < +\infty. \tag{2.11}$$

We first consider the special case where the mass distribution  $\mu_2$  is concentrated in one point  $\zeta_0$ . Then

$$I = \int_{x_0 - r_0}^{x_0 + r_0} |x - \zeta_0|^{-p} dx \leq \int_{x_0 - r_0}^{x_0 + r_0} |x - x_0|^{-p} dx = \frac{2r_0^{1-p}}{1-p}. \tag{2.12}$$

Let us now proceed to a measure  $\mu_2$  which consists of a finite number of concentrated masses,  $\alpha_1 p$  in  $\zeta_1, \dots, \alpha_n p$  in  $\zeta_n$ ,  $\sum \alpha_i = 1$ ,  $\alpha_i > 0$  ( $i = 1, 2, \dots, n$ ). We introduce the notation  $I_N$  for the integral  $I$  in which  $\log|z - \zeta|$

has been replaced by  $\log^N |z - \zeta| = \max [\log |z - \zeta|, -N]$ ,  $N$  denoting an arbitrarily large constant. By an application of HÖLDER's inequality [15, p. 140] and of (2.12) we obtain

$$\begin{aligned} I_N &= \int_{x_0-r_0}^{x_0+r_0} \exp \left\{ - \sum_{i=1}^n \alpha_i p \log^N |x - \zeta_i| \right\} dx \\ &\leq \prod_{i=1}^n \left[ \int_{x_0-r_0}^{x_0+r_0} \exp \{ -p \log^N |x - \zeta_i| \} dx \right]^{\alpha_i} \\ &\leq \prod_{i=1}^n \left[ \int_{x_0-r_0}^{x_0+r_0} |x - \zeta_i|^{-p} dx \right]^{\alpha_i} \leq \frac{2r_0^{1-p}}{1-p} . \end{aligned}$$

Let us now drop every special assumption about  $\mu_2$ . In the general case  $I_N$  can always be approximated arbitrarily close for fixed  $N$  by substituting for  $\mu_2$  a suitable measure of the special type considered above. Hence

$$I_N \leq 2r_0^{1-p}/(1-p)$$

holds without restriction. Letting  $N \rightarrow +\infty$  we obtain (2.11) in the limit. Q. E. D.

**Lemma 2.** *Suppose there exists a sequence  $\{\sigma_n\}$ ,  $n = 1, 2, 3, \dots$ , of rectifiable curves in  $\Omega$ , a subcompact  $K$  of  $\Omega_0$  and a number  $C$  such that the following is true:*

- (a) *each  $\sigma_n$  has a non-empty intersection with  $K$ ,*
- (b)  *$\{\sigma_n\}$  comes arbitrarily near to  $\Gamma$ ,*
- (c)  *$\int_{\sigma_n} e^u |dz| < C$  for all  $n$ .*

*Then there exists a locally rectifiable path  $\sigma$  in  $\Omega$ , tending to  $\Gamma$ , such that (2.7) is satisfied.*

**Proof.** We may assume that  $\Omega$  is a circular ring

$$R_1 < |z| < R_2 \quad (0 \leq R_1 < R_2 \leq +\infty) .$$

For, if this lemma has been proved for a particular region  $\Omega$ , then it is immediately seen to be valid for the whole class of conformally equivalent domains. This is easily proved by transplanting the metric  $e^{u(z)} |dz|$  under conformal representation.

We introduce a sequence of circles  $\gamma_m = [|z| = r_m]$ ,  $m = 1, 2, 3, \dots$ , in  $\Omega$ , supposing that  $\gamma_1$  encloses  $K$  and that  $r_m \nearrow R_2$  for  $m \nearrow +\infty$ . We further assume that no singular point of  $\mu$  lies on these circles. If one of the curves  $\sigma_n$  tends to  $\Gamma$ , then we have nothing to prove. If this is not the case, then there exists a subsequence  $\{\sigma'_n\}$  of  $\{\sigma_n\}$  such that  $\sigma'_n$  intersects  $\gamma_n$  for all  $n$ . From

this and hypothesis (a) we conclude that  $\sigma'_n$  contains an arc which leads from  $\gamma_1$  to  $\gamma_n$ . We subdivide it into  $\sigma'_n(1, 2)$  (leading from  $\gamma_1$  to  $\gamma_2$ ),  $\dots$ ,  $\sigma'_n(n-1, n)$  (leading from  $\gamma_{n-1}$  to  $\gamma_n$ ). By making use of CANTOR's diagonal process we select a subsequence  $\{\sigma''_n\}$  of  $\{\sigma'_n\}$  such that the common endpoints of  $\sigma''_n(m-1, m)$  and  $\sigma''_n(m, m+1)$  on  $\gamma_m$  converge to a limit point  $z_m$  for all  $m$ . Let now  $m$  be fixed. By Lemma 1, there exists an open arc  $\alpha_m$  of  $\gamma_m$ , containing  $z_m$ , such that  $\int_{\alpha_m} e^u |dz| < 2^{-m}$ . Furthermore, there exists an index  $N(m)$  such that the following conditions are satisfied:

(a) The inner and outer endpoints of  $\sigma''_N(m, m+1)$  lie on  $\alpha_m$  and  $\alpha_{m+1}$ , respectively,

$$(b) \quad \int_{\sigma''_N(m, m+1)} e^u |dz| < \inf_{n > N} \int_{\sigma''_n(m, m+1)} e^u |dz| + 2^{-m-1}.$$

We define  $\sigma$  to consist of the curves  $\sigma''_{N(1)}(1, 2)$ ,  $\sigma''_{N(2)}(2, 3)$ ,  $\dots$ , joined by subarcs of  $\alpha_2, \alpha_3, \dots$ . The reader will easily convince himself that

$$\int_{\sigma} e^u |dz| < C + 1.$$

This proves Lemma 2.

**Proof of Theorem 1.** There is some expository advantage in assuming that  $\Omega$  is a circular ring  $R_1 < |z| < R_2$  ( $0 \leq R_1 < R_2 \leq +\infty$ ). This can be done without loss of generality. Indeed, an arbitrary  $\Omega$  can always be mapped conformally onto a circular annulus  $R_1 < |\zeta| < R_2$  in such a way that  $\Gamma$  corresponds to the outer circle  $|\zeta| = R_2$ . Under such a representation both length element  $ds$  and flux  $\Phi$  are invariant if  $u$  is transformed according to (2.8).

It follows from the hypotheses that there exists a number  $\eta$  ( $0 < \eta < 1$ ) and a radius  $r_1$  ( $R_1 < r_1 < R_2$ ) such that

$$\Phi(u_2, |z| = r_1; \Gamma) > 1 + 2\eta, \quad (2.13)$$

$$\Phi_1(\Gamma) - \Phi(u_1, |z| = r_1; \Gamma) < \eta. \quad (2.14)$$

**Lemma 3.** *Let the radius  $\varrho_1$  ( $r_1 < \varrho_1 < R_2$ ) be chosen arbitrarily. Then there is a number  $C$  with the following property: Given any  $\varrho_2$  ( $\varrho_1 < \varrho_2 < R_2$ ), there exists a rectifiable curve  $\alpha$ , leading from  $|z| = \varrho_1$  to  $|z| = \varrho_2$ , such that*

$$\int_{\alpha} e^u |dz| < C. \quad (2.15)$$

**Proof.** We introduce three radii  $s_1, s_2$  and  $r_2$  satisfying the inequalities

$$R_1 < r_1 < s_1 < \varrho_1 < \varrho_2 < s_2 < r_2 < R_2. \quad (2.16)$$

In addition we require that there should be no singular point of  $\mu$  on  $|z| = s_1$ .

In the following all radii with index 1 have to be considered fixed. On the other hand,  $\varrho_2$ ,  $s_2$  and  $r_2$  are variable within the above limits and our estimation ultimately does not depend on their choice. By a theorem of F. RIESZ [27, II, p. 357] the representations

$$u_1(z) = h_1(z) - \int_{\omega} g(z, \zeta) d\mu_1(e_{\zeta}) \quad (2.17)$$

and

$$u_2(z) = h_2(z) - \int_{\omega} g(z, \zeta) d\mu_2(e_{\zeta}) \quad (2.18)$$

hold in  $\omega = [r_1 < |z| < r_2]$ , where  $h_1$  and  $h_2$  are the best harmonic majorants of  $u_1$  and  $u_2$ , respectively, in  $\omega$  and  $g$  denotes GREEN's function for this domain. We define

$$v(z) = h_1(z) - u_2(z) = u(z) + \int_{\omega} g(z, \zeta) d\mu_1(e_{\zeta}) . \quad (2.19)$$

$v$  is superharmonic in  $\omega$  and admits the representation (F. RIESZ [27, II, p. 350])

$$v(z) = H(z) - \int_{\omega} \log|z - \zeta| d\mu_2(e_{\zeta}) , \quad (2.20)$$

where  $H$  is harmonic in  $\omega$ . It follows from (2.13) and (2.14) that

$$\frac{1}{2\pi} \int_{\gamma} \frac{\partial H}{\partial n} |dz| < -1 - \eta , \quad (2.21)$$

$\gamma$  being an arbitrary smooth JORDAN curve in  $\omega$  which encloses  $|z| = r_1$ ,  $n$  denoting the outer normal.

Let  $\vartheta = (\varrho_1 - s_1)/2$ . We define

$$\omega_0 = [r_1 < |z| \leq s_1 + \vartheta] , \quad \omega_1 = [s_1 + \vartheta < |z| < r_2] ,$$

$\mu_{20}(e) = \mu_2(e \cap \omega_0)$ ,  $\mu_{21}(e) = \mu_2(e \cap \omega_1)$ ,  $m_0 = \mu_2(\omega_0)$ ,  $m_1 = \mu_2(\omega_1)$  and

$$v_1(z) = H(z) - \int_{\omega_1} \log|z - \zeta| d\mu_{21}(e_{\zeta}) . \quad (2.22)$$

Further, let

$$V(z) = u(z) + \int_{|z| > r_1} G(z, \zeta) d\mu_1(e_{\zeta}) , \quad (2.23)$$

where  $G$  denotes GREEN's function for  $|z| > r_1$ . For  $|z| = s_1$ ,  $v_1(z) \leq v(z) + m_0 \log(2\varrho_1)$ . Furthermore,  $V$  is superharmonic in  $\omega$ , the associated measure being  $\mu_2(e)$ , and  $v \leq V$ . Since the circle  $|z| = s_1$  contains no singular point of  $\mu$ , we have, by Lemma 1

$$\int_{|z|=s_1} e^{v_1} |dz| \leq (2\varrho_1)^{m_0} \int_{|z|=s_1} e^v |dz| \leq (2\varrho_1)^{m_0} \int_{|z|=s_1} e^V |dz| = C_1 < +\infty , \quad (2.24)$$

where  $C_1$  is a constant not depending on the choice of  $\varrho_2$ ,  $s_2$  and  $r_2$ .

For  $\varrho_1 \leq |z| \leq \varrho_2$ ,  $u \leq v \leq v_1 - m_0 \log \vartheta$ . Hence, for any curve  $\alpha$  contained in this annulus

$$\int_{\alpha} e^u |dz| \leq \vartheta^{-m_0} \int_{\alpha} e^{v_1} |dz| . \quad (2.25)$$

We are now going to demonstrate that there exists a rectifiable curve  $\alpha$ , leading from  $|z| = \varrho_1$  to  $|z| = \varrho_2$ , and such that

$$\int_{\alpha} e^{v_1} |dz| \leq C_2 \int_{|z|=s_1} e^{v_1} |dz| + C_3 , \quad (2.26)$$

where  $C_2$  and  $C_3$  are constants not depending on the choice of  $\varrho_2$ ,  $r_2$  and  $s_2$ . Lemma 3 will be an immediate consequence of (2.24), (2.25) and (2.26).

In order to establish (2.26) we first approximate the measure  $\mu_{21}(e)$  by a finite number of concentrated masses. This is done by the following construction: Choose an integer  $N \geq 2$  such that

$$m_1 \log \frac{N+1+2\sqrt{2}}{N+1-2\sqrt{2}} < \log 2 . \quad (2.27)$$

Let  $\zeta_1, \zeta_2, \dots, \zeta_m$  designate those points (necessarily finite in number) which support a concentrated mass of weight  $\geq \frac{1}{4(2N+1)^2}$  in the measure  $\mu_{21}$ . Denoting the corresponding masses by  $p_1, p_2, \dots, p_m$ , we define

$$w(z) = H(z) - \sum_{k=1}^m p_k \log |z - \zeta_k| . \quad (2.28)$$

We introduce the compact set  $K$  which is obtained by subtracting from the annulus  $s_1 \leq |z| \leq s_2$  the open disks  $|z - \zeta_k| < \delta_k$  ( $k = 1, 2, \dots, m$ ), where the radii  $\delta_k$  are chosen small enough so that the following conditions are satisfied:

(a) the sets  $|z| = s_1$ ,  $|z - \zeta_1| \leq 2\delta_1, \dots, |z - \zeta_m| \leq 2\delta_m$  are disjoint,

(b) whenever  $\zeta_k$  is a singular point of  $\mu$  (i. e.  $p_k \geq 1$ ), then we choose  $\delta_k$  so small that

$$\int_{\kappa} e^w |dz| > \frac{2^{m_1+2} r_2^{m_1}}{1 - \cos(\pi\eta)} \int_{|z|=s_1} e^{v_1} |dz| \quad (2.29)$$

for any rectifiable curve  $\kappa$  leading from  $|z| = s_1$  to  $|z - \zeta_k| = 2\delta_k$ ,<sup>4)</sup>

(c) we require the remaining  $\delta_k$ 's to be so small that

$$\sum_{p_k < 1} \int_{|z - \zeta_k| = 2\delta_k} e^{v_1} |dz| < 1 . \quad (2.30)$$

<sup>4)</sup> It is easy to prove that such a choice is always possible.

This condition can always be satisfied since

$$\lim_{\delta \rightarrow 0} \int_{|z - \zeta_k| = \delta} e^{v_1} |dz| = 0 \quad (2.31)$$

at any non-singular point  $\zeta_k$ . (2.31) can be proved by a reasoning which is quite similar to the one used in the proof of Lemma 1 and which we do not reproduce here.

Let  $\nu(e)$  denote the measure which is obtained from  $\mu_{21}(e)$  after subtracting the concentrated masses  $p_1$  in  $\zeta_1, \dots, p_m$  in  $\zeta_m$ . We have

$$v_1(z) = w(z) - \int_{\omega_1} \log |z - \zeta| d\nu(e_\zeta) . \quad (2.32)$$

There exists a number  $d > 0$  such that  $\nu(e) \leq \frac{1}{4(2N+1)^2}$  for all BOREL sets  $e$  of diameter  $\leq d$ . Let such a  $d$  be chosen, requiring in addition that  $d < 1$ .

The function  $w(z)$  is uniformly continuous on  $K$ .

We now choose a number  $L > 0$ , so small that the conditions (2.33) to (2.37) are satisfied:

$$|z_1 - z_2| \leq 3\sqrt{2}L \quad \text{implies} \quad (2.33)$$

$$|w(z_1) - w(z_2)| < \log 2 \quad \text{for all } z_1, z_2 \in K ,$$

$$m_1 \log \frac{\vartheta}{\vartheta - L} < \log 2 , \quad (2.34)$$

$$\sqrt{2}L < d , \quad \sqrt{2}(2N+1)L < 1 , \quad (2.35)$$

$$2\sqrt{2}L < \delta_k \quad (k = 1, 2, \dots, m) , \quad (2.36)$$

$$m_1 d^{-m_1} e^M (N+1)^2 N^{-\frac{1}{4}} \lambda^{\frac{1}{2}} |\log \lambda| < 1 \quad (2.37)$$

for  $0 < \lambda \leq \sqrt{2}L$ ,  $M$  denoting the maximum of  $w(z)$  on  $K$ .

Now we cover the  $(x, y)$ -plane by a net of squares  $\Sigma$  with sides of length  $L$  ( $x = iL, y = jL; i, j = 0, \pm 1, \pm 2, \dots$ ) and replace the measure  $\nu(e)$  by a finite number of concentrated masses, assigning to the points  $((i + \frac{1}{2})L, (j + \frac{1}{2})L)$  the weights  $\nu([iL \leq x < (i+1)L] \cap [jL \leq y < (j+1)L])$ . Let these be the masses  $p_{m+1}$  in  $\zeta_{m+1}, \dots, p_n$  in  $\zeta_n$ . We define

$$w_1(z) = w(z) - \sum_{k=m+1}^n p_k \log |z - \zeta_k| = H(z) - \sum_{k=1}^n p_k \log |z - \zeta_k| . \quad (2.38)$$

We now state: There exists a polygon  $\beta$ , leading from  $|z| = s_1$  to  $|z| = s_2$ , such that

$$\int_{\beta} e^{w_1} |dz| < \frac{2}{1 - \cos(\pi\eta)} \int_{|z|=s_1} e^{w_1} |dz| . \quad (2.39)$$

We should like to point out that the proof of this statement constitutes the kernel of our demonstration of Lemma 3. In fact, if  $u$  were assumed to be superharmonic and of the type  $w_1$  (i. e. harmonic with isolated logarithmic singularities) throughout  $\Omega$ , then the following reasoning would represent the complete proof of this lemma.

Let  $z_0$  be an arbitrary point of the annulus  $A = [s_1 \leq |z| \leq s_2]$ . We define  $A(z_0) = \inf_{\gamma} \int_{\gamma} e^{w_1} |dz|$ , where  $\gamma$  varies on the set  $P(z_0)$  consisting of all rectifiable, closed curves on  $A$  which are not nullhomotopic and pass through  $z_0$ . If  $z_0$  is not a singular point of  $\mu$ , then  $A(z_0)$  is finite and there exists at least one *minimal curve*  $\bar{\gamma}(z_0) \in P(z_0)$  such that

$$A(z_0) = \int_{\bar{\gamma}} e^{w_1} |dz| . \quad (2.40)$$

Indeed,  $\bar{\gamma}$  can be constructed in the usual way as the limit of a suitably chosen minimal sequence.  $A(z)$  is continuous on  $A - \bigcup_{k \geq 1} \zeta_k$ .

We decompose the set  $D = [s_1 \leq |z| \leq s_2]$  into three disjoint subsets  $D = \bigcup_{i=1}^3 D_i$ , where

$$(a) \quad D_1 = D \cap \bigcup_{k \geq 1} [|z - \zeta_k| < 2\delta_k] ,$$

(b)  $D_2$  consists of those points of  $D - D_1$  which possess at least one minimal curve contained in  $D$ ,

$$(c) \quad D_3 = D - (D_1 \cup D_2) .$$

Some of these sets may be empty.  $D_2$  is closed.

We now discuss some properties of minimal curves which contain no double point. Assume first, for simplicity, that  $\bar{\gamma}(z_0)$  is completely in the interior of  $A$  and that none of the  $\zeta_k$ 's lie on it ( $k = 1, 2, \dots, n$ ). Then  $w_1$  is harmonic in some (doubly connected) neighborhood  $V$  of  $\bar{\gamma}$ . Hence the function

$$\zeta = \Psi(z) = \int_{z_0}^z e^{w_1 + i w_1^*} dz , \quad (2.41)$$

where  $w_1^*$  is conjugate harmonic to  $w_1$ , yields a conformal mapping of the universal covering surface  $\tilde{V}$  of  $V$  onto some simply connected RIEMANN surface extending over the  $\zeta$ -plane. Since  $e^{w_1} |dz| = |d\zeta|$ , the image of any subarc of  $\bar{\gamma}$  not containing  $z_0$  in its interior (and considered on any sheet of  $\tilde{V}$ ) is a straight line segment in the  $\zeta$ -plane. Hence

$$\arg d\zeta = w_1^* + \arg dz = \text{const.} \quad (2.42)$$

along such an arc. Furthermore, (2.21) implies

$$\int_{\bar{\gamma}} \frac{\partial w_1^*}{\partial s} ds = \int_{\bar{\gamma}} \frac{\partial w_1}{\partial n} ds = \int_{\bar{\gamma}} \frac{\partial H}{\partial n} ds - 2\pi \sum_{\zeta_k \text{ inside } \bar{\gamma}} p_k < -2\pi(1 + \eta) , \quad (2.43)$$

where  $s$  denotes the arc length,  $n$  the exterior normal, and the integration is performed in the positive sense. From (2.42) and (2.43) we conclude that, in the neighborhood of  $z_0$ ,  $\bar{\gamma}$  consists of two analytic arcs which intersect at an exterior angle  $\Theta(\bar{\gamma}, z_0)$  satisfying the inequality

$$\Theta(\bar{\gamma}, z_0) < \pi(1 - 2\eta) . \quad (2.44)$$

If we now allow  $\bar{\gamma}$  to have points in common with  $|z| = s_1$ , then (2.42) will no more be true in general. However, it follows easily from a consideration of the mapping (2.41) that  $w_1^* + \arg dz$  increases monotonically if  $\bar{\gamma}$  is followed in the positive sense. From this and (2.42) the inequality (2.44) is again obtained.

Finally, we also admit that  $\bar{\gamma}$  may pass through some of the  $\zeta_k$ 's. In such points  $\bar{\gamma}$  will not possess corners because these would make shortcuts (in the metric  $e^{w_1}|dz|$ ) possible, contrary to the definition of minimal curves. Near these points we replace  $\bar{\gamma}$  temporarily by small circular arcs lying in the interior region. By making use of the mapping (2.41) (integrating in a neighborhood of the modified curve) and by our knowledge of the behavior of  $w_1^*$  along the small circles, we verify again the monotonicity of  $w_1^* + \arg dz$  and the ensuing relation (2.44). No essential difficulty arises if  $z_0$  itself coincides with one of the  $\zeta_k$ 's.

If  $\bar{\gamma}$  has points in common with  $|z| = s_2$ , then (2.44) will not be fulfilled in general.

We shall now investigate the behavior of the function  $A(z)$  in the neighborhood  $U$  of a point  $z_0$  on  $D_2$ . To this end we map  $U$ , by (2.41), conformally onto a domain  $\tilde{U}$  in the  $\zeta$ -plane. (Assume first, for simplicity, that  $w_1$  is harmonic at  $z_0$ .) There exists a minimal curve  $\bar{\gamma}(z_0)$  which is contained in  $D$ . For the present we make the additional assumption that  $\bar{\gamma}$  is a simple closed curve. Then the image of  $\bar{\gamma}$  in  $\tilde{U}$  consists of two straight line segments,  $l_1$  and  $l_2$ , intersecting at  $\zeta_0 = \Psi(z_0)$  under the angle  $\Theta(\bar{\gamma}, z_0)$ . Let  $\zeta'_0$  be a point on the bisector  $b$  of  $\Theta$ , and let  $\zeta_1$  and  $\zeta_2$  denote the points of intersection of the normal  $n$  to  $b$  through  $\zeta'_0$  with  $l_1$  and  $l_2$ , respectively. (Choose  $|\zeta_0 - \zeta'_0|$  small enough so that the triangle  $\zeta_0\zeta_1\zeta_2$  lies in  $\tilde{U}$ .) We have

$$|\zeta_0 - \zeta'_0| = \frac{\cos(\Theta/2)}{2(1 - \sin(\Theta/2))} [(|\zeta_0 - \zeta_1| + |\zeta_0 - \zeta_2|) - (|\zeta'_0 - \zeta_1| + |\zeta'_0 - \zeta_2|)] .$$

The (Euclidean) bisector of  $\Theta$  in the  $z$ -plane is transformed into an analytic curve  $a$  tangential to  $b$  at  $z_0$ . Let  $\zeta_0''$  denote the point of intersection of  $a$  with  $n$ .

Then

$$\int_{\zeta_0}^{\zeta_0''} |d\zeta| = |\zeta_0 - \zeta_0'| + o(|\zeta_0 - \zeta_0'|) .$$

Hence, for sufficiently small  $|\zeta_0 - \zeta_0'|$

$$\int_{\zeta_0}^{\zeta_0''} |d\zeta| < \frac{\cos(\Theta/2)}{1 - \sin(\Theta/2)} [ (|\zeta_0 - \zeta_1| + |\zeta_0 - \zeta_2|) - (|\zeta_0'' - \zeta_1| + |\zeta_0'' - \zeta_2|) ] . \quad (2.45)$$

Let  $z_0'' = \Psi^{-1}(\zeta_0'')$ . From (2.44), (2.45) and the definition of  $A$  we conclude that

$$\int_{z_0}^{z_0''} e^{w_1} |dz| < \frac{1}{1 - \cos(\pi\eta)} [A(z_0) - A(z_0'')] , \quad (2.46)$$

where  $z_0 z_0''$  denotes a (sufficiently small) segment on the (Euclidean) bisector of  $\Theta(\bar{\gamma}, z_0)$ .

Only slight modifications are needed for the case where  $z_0$  coincides with one of the  $\zeta_k$ 's. Again we make use of the mapping (2.41). In order to obtain uniqueness we slit the domain  $U$  along a line which leads from  $z_0$  into the interior of  $\bar{\gamma}$ . The mapping  $\Psi$  is no more conformal at  $z_0$ . However, the reader will easily convince himself that  $\Theta$  is decreased. So our estimations hold a fortiori.

Finally, if  $\bar{\gamma}$  contains double points, then it can be proved without difficulty that (2.46) is satisfied if  $z_0 z_0''$  is defined to be a (sufficiently small) segment on the tangent to either of the two branches of  $\bar{\gamma}$  issuing from  $z_0$ .

On  $D_2$  we now define a complex-valued function  $T(z)$ . Let  $z \in D_2$  and  $\varphi$  ( $0 \leq \varphi < 2\pi$ ) be fixed and let  $t(z, \varphi)$  denote the largest number with the following property: For all  $\tau$  in the interval  $0 < \tau < t$  the point  $z = \tau e^{i\varphi}$  is contained in  $A$  and the inequality

$$\int_0^\tau \exp \{w_1(z + re^{i\varphi})\} dr < \frac{1}{1 - \cos(\pi\eta)} [A(z) - A(z + \tau e^{i\varphi})]$$

holds. Suppose  $\{\varphi_i(z)\}$  is a sequence of arguments such that  $\{t(z, \varphi_i)\}$  tends to the least upper bound  $t_m(z)$  of  $t(z, \varphi)$ . By (2.46),  $t_m(z)$  is always positive. We choose an arbitrary limit point  $\varphi_m$  of  $\{\varphi_i\}$  and define  $T(z) = t_m e^{i\varphi_m}$ . It follows from the definition of  $T(z)$  and the continuity of  $A(z)$  that, for all  $z \in D_2$

$$\int_0^{t_m} \exp \{w_1(z + re^{i\varphi_m})\} dr \leq \frac{1}{1 - \cos(\pi\eta)} [A(z) - A(z + T(z))] . \quad (2.47)$$

Furthermore, by making use of the definitions it can be verified in a straightforward way that  $t_m(z)$  is lower semicontinuous. Let  $t_{min}$  denote the (positive) minimum of  $t_m(z)$  on the (compact) set  $D_2$ .

Let us now construct  $\beta$ . We choose an arbitrary point  $z_0$  on  $|z| = s_1$ . Obviously

$$A(z_0) \leq \int_{|z|=s_1} e^{w_1} |dz| . \quad (2.48)$$

If  $z_0 \in D_2$ , then we define  $z_1 = z_0 + T(z_0)$ ,  $z_2 = z_1 + T(z_1)$ , etc., until we arrive at a point  $z_n$  not contained in  $D_2$ . This will always happen after a finite number of steps. This follows from  $A(z) > 0$  and the inequalities (implied by (2.47))

$$A(z_k) - A(z_{k+1}) \geq t_{\min} e^{w_{1\min}} [1 - \cos(\pi\eta)]$$

( $k = 0, 1, 2, \dots$ ),  $w_{1\min}$  denoting the (finite) minimum of  $w_1$  on  $A$ .

Let  $\beta_1$  denote the polygon  $(z_0, z_1, \dots, z_n)$ . From (2.47) (formulated for  $z = z_0, z_1, \dots, z_{n-1}$ ) and (2.48) we conclude that

$$\int_{\beta_1} e^{w_1} |dz| \leq \frac{1}{1 - \cos(\pi\eta)} \int_{|z|=s_1} e^{w_1} |dz| . \quad (2.49)$$

Assume that  $z_n \in D_3$ . Then every minimal curve  $\bar{\gamma}(z_n)$  intersects  $|z| = \varrho_2$ . Hence there exists a rectifiable curve, joining  $z_n$  with some point on  $|z| = \varrho_2$ , and of length  $\leq A(z_n)/2 \leq A(z_0)/2$  in the metric  $e^{w_1} |dz|$ . Combined with (2.48) this implies the existence of a polygon  $\beta_2$  with the same endpoints and such that

$$\int_{\beta_2} e^{w_1} |dz| < \int_{|z|=s_1} e^{w_1} |dz| . \quad (2.50)$$

We define  $\beta = \beta_1 + \beta_2$ . (2.39) follows from (2.49) and (2.50).

If  $|z_n| > \varrho_2$ , then  $\beta$  reduces to a portion of  $\beta_1$ . If  $z_0 \in D_3$ , then  $\beta$  consists of  $\beta_2$  only. In these cases (2.39) is true a fortiori.

We are left with the possibility  $z_n \in D_1$ . This never occurs. Indeed, by (2.38), we have

$$w \leq w_1 + m_1 \log(2r_2) . \quad (2.51)$$

Let  $S_k$  be an arbitrary closed square of the net  $\Sigma$  which intersects with  $\omega_1$ . Then, for any  $z \in [|z| = s_1]$  and arbitrary  $\zeta \in S_k$

$$\log \left| \frac{z - \zeta}{z - \zeta_k} \right| < \log \frac{\vartheta}{\vartheta - L} ,$$

$\zeta_k$  denoting the center of  $S_k$ . Hence (2.32), (2.34) and (2.38) imply

$$w_1 \leq v_1 + \log 2 \quad \text{on} \quad |z| = s_1 .$$

Therefore

$$\int_{|z|=s_1} e^{w_1} |dz| \leq 2 \int_{|z|=s_1} e^{v_1} |dz| . \quad (2.52)$$

It follows from (2.51), (2.49) and (2.52) that

$$\int_{\beta_1} e^w |dz| \leq \frac{2^{m_1+1} r_2^{m_1}}{1 - \cos(\pi\eta)} \int_{|z|=s_1} e^{v_1} |dz| .$$

Now, if  $z_n \in D_1$ , this would contradict (2.29).

We thus have proved the existence of  $\beta$ . By applying (2.39) instead of (2.49) in the above reasoning one finds that  $\beta$  does not intersect  $D_1$ .

With the polygon  $\beta$  we now associate a curve  $\alpha$  which leads from  $|z| = \varrho_1$  to  $|z| = \varrho_2$ . This is done by the following construction: Let  $E(z)$  denote the set consisting of those (1, 2 or 4) closed squares of the net  $\Sigma$  which contain the point  $z$ . We connect the endpoint  $z_0$  of  $\beta$  on  $|z| = s_1$  with the last point of intersection  $z'_1$  of  $\beta$  with  $E(z_0)$ ,  $z'_1$  with the last point of intersection  $z'_2$  of  $\beta$  with  $E(z'_1)$ , and so forth, until we arrive at the endpoint of  $\beta$  on  $|z| = \varrho_2$ . We obtain a polygon  $\beta'$ . Now, if  $\beta'$  should penetrate into some of the disks

$$|z - \zeta_k| < 2\delta_k \quad (p_k < 1; k = 1, 2, \dots, m) ,$$

then we replace in each case the subpolygon between the first entry and the last exit by an arc on  $|z - \zeta_k| = 2\delta_k$ . If the endpoint of  $\beta$  on  $|z| = \varrho_2$  should lie in  $|z - \zeta_k| < 2\delta_k$  ( $p_k < 1; k = 1, 2, \dots, m$ ), then we follow the circle  $|z - \zeta_k| = 2\delta_k$  from the first entry until we reach  $|z| = \varrho_2$ . The resulting curve contains a portion  $\alpha$  which leads from  $|z| = \varrho_1$  to  $|z| = \varrho_2$  and is contained in  $\varrho_1 \leq |z| \leq \varrho_2$ . We are going to prove that

$$\int_{\alpha} e^{v_1} |dz| \leq \frac{32}{3} \int_{\beta} e^{v_1} |dz| + \frac{259}{3} . \quad (2.53)$$

The construction clearly implies that  $\alpha$  does not enter any of the disks  $|z - \zeta_k| < \delta_k$  for which  $p_k < 1$  ( $k = 1, 2, \dots, m$ ). From (2.36) and the fact that  $\beta$  does not intersect  $D_1$  it follows that the same is true for  $p_k \geq 1$ . So  $\alpha$  lies in  $K$ .

We decompose  $\alpha = \alpha_P + \alpha_C$ ,  $\alpha_P$  consisting of a finite number of polygons,  $\alpha_C$  of a finite number of circular arcs. We have, by (2.30)

$$\int_{\alpha_C} e^{v_1} |dz| < 1 . \quad (2.54)$$

$\alpha_P$  is composed of straight line segments  $\alpha_{P_1}, \alpha_{P_2}, \dots, \alpha_{P_r}$  which are contained in some closed squares  $S_1, S_2, \dots, S_r$  of the net  $\Sigma$ . It follows from the construction of  $\alpha$  that these squares are different from each other. We complete the finite sequence  $S_1, S_2, \dots, S_r$  to an infinite sequence  $\{S_i\}$  which enumerates all squares of  $\Sigma$ . Let  $Z_i$  denote the centers of  $S_i$  ( $i = 1, 2, 3, \dots$ ). Some of the  $Z_i$ 's are identical with  $\zeta_{m+1}, \zeta_{m+2}, \dots, \zeta_n$  (introduced for the definition of  $w_1$ ), say  $Z_{i_1} = \zeta_{m+1}, Z_{i_2} = \zeta_{m+2}, \dots, Z_{i_{n-m}} = \zeta_n$ . For the corresponding masses

we introduce the change of notation  $P_{i_1} = p_{m+1}, P_{i_2} = p_{m+2}, \dots, P_{i_{n-m}} = p_n$ , defining  $P_i = 0$  for all other indices.

Let  $S'_i$  and  $S''_i$  denote the closed squares of center  $Z_i$  with sides of lengths  $3L$  and  $(2N+1)L$ , respectively. We define

$$A_i = \max_{z \in (S'_i \cap K)} w(z) \quad \text{and} \quad a_i = \min_{z \in (S'_i \cap K)} w(z)$$

(2.33) implies

$$A_i - a_i < \log 2. \quad (2.55)$$

Further, let  $b_{ik}$  and  $B_{ik}$  denote the maximum and minimum, respectively, of the function  $\log|z - \zeta|$ ,  $z$  varying on  $S'_i$ ,  $\zeta$  on  $S_k$ . If  $S_k$  is not contained in  $S''_i$ , then we have, by (2.27)

$$m_1(b_{ik} - B_{ik}) < \log 2. \quad (2.56)$$

Let  $Q_i = \nu(S''_i)$ . It follows from (2.35) and the definition of  $d$  that

$$Q_i \leq \frac{1}{4} \quad (i = 1, 2, 3, \dots). \quad (2.57)$$

An arbitrary point  $z$  of the plane is contained in at most  $(2N+2)^2$  of the sets  $S''_i$ . From this we infer that

$$\sum_{i=1}^{\infty} Q_i \leq 4(N+1)^2 m_1. \quad (2.58)$$

We have

$$\begin{aligned} \int_{\alpha_{P_i}} e^{v_1} |dz| &\leq \int_{\alpha_{P_i}} \exp \left\{ A_i - \sum_{k=1}^{\infty} P_k B_{ik} - \int_{S''_i} \log|z - \zeta| d\nu(e_{\zeta}) \right\} |dz| \\ &\leq 2 \exp \left\{ A_i - \sum_{k=1}^{\infty} P_k B_{ik} \right\} \int_0^{\lambda_i/2} x^{-Q_i} dx \leq \frac{32}{3} \exp \left\{ a_i - \sum_{k=1}^{\infty} P_k b_{ik} \right\} \lambda_i^{1-Q_i}. \end{aligned} \quad (2.59)$$

Here, as indicated by the double prime, we do not extend the summation over those indices  $k$  for which  $S_k$  is contained in  $S''_i$ . This estimation is based on the representation (2.32). In the first step we replace  $w(z)$  as well as the potential of the masses  $\nu$  outside  $S''_i$  by a constant. Then we observe that the integral attains its maximum when the total mass  $Q_i$  is concentrated in the center of  $\alpha_{P_i}$ . This follows from an application of HÖLDER'S inequality. (The reader is referred to the proof of Lemma 1 where exactly the same reasoning has been reproduced in full detail.) Finally, in the third step we evaluate the integral and make use of (2.55), (2.56), (2.57) and the inequality  $\sum'' P_k \leq m_1$ .

The segment  $\alpha_{P_i}$  has been introduced as a shortcut of some subpolygon  $\beta_i$  of  $\beta$  which, at least for a portion of length  $\geq \lambda_i$ , is contained in  $S'_i$ . By (2.36) and the construction of  $\alpha$ ,  $S'_i \subset K$  ( $i = 1, 2, \dots, r$ ). Hence

$$\int_{\beta_i} e^{w_1} |dz| \geq \lambda_i \exp \left\{ a_i - \sum_{k=1}^{\infty} P_k b_{ik} \right\}. \quad (2.60)$$

Here  $w(z)$  as well as the potential of the masses  $\nu$  outside  $S_i''$  have been replaced by a constant. The potential of the masses on  $S_i''$  has been neglected. (We are allowed to do this because of the second relation of (2.35).)

From (2.59) and (2.60) we conclude that

$$\begin{aligned} \int_{\alpha_{P_i}} e^{v_1} |dz| &\leq \frac{32}{3} \int_{\beta_i} e^{w_1} |dz| + \frac{32}{3} \exp \{a_i - \sum_{k=1}^{\infty} P_k b_{ik}\} Q_i \frac{\lambda_i^{1-Q_i} - \lambda_i}{Q_i} \\ &\leq \frac{32}{3} \int_{\beta_i} e^{w_1} |dz| + \frac{32}{3} \exp \{a_i - \sum_{k=1}^{\infty} P_k b_{ik}\} Q_i \lambda_i^{3/4} |\log \lambda_i| \quad (i = 1, 2, \dots, r). \end{aligned} \quad (2.61)$$

The first step is obvious. In the second inequality we make use of the mean value theorem of differential calculus, (2.57) and the inequality  $\lambda_i < 1$  which is implied by the second relation of (2.35). We observe that the final estimation in (2.61) is also valid if  $Q_i = 0$ , although in this case the intermediate step has no meaning.

Obviously

$$a_i \leq M \quad (i = 1, 2, \dots, r), \quad (2.62)$$

$M$  denoting the maximum of  $w(z)$  on  $K$ .

It is easy to verify that

$$\sum_{k=1}^{\infty} P_k b_{ik} \geq \frac{1}{4} \log (NL) + m_1 \log d. \quad (2.63)$$

We have, by (2.62), (2.63), (2.37) and (2.58)

$$\begin{aligned} \sum_{i=1}^r \exp \{a_i - \sum_{k=1}^{\infty} P_k b_{ik}\} Q_i \lambda_i^{3/4} |\log \lambda_i| &\leq \sum_{i=1}^r e^M (NL)^{-1/4} d^{-m_1} Q_i \lambda_i^{3/4} |\log \lambda_i| \\ &\leq 2 \sum_{i=1}^r Q_i e^M N^{-1/4} d^{-m_1} \lambda_i^{1/2} |\log \lambda_i| \leq \frac{2}{m_1 (N+1)^2} \sum_{i=1}^r Q_i \leq 8. \end{aligned} \quad (2.64)$$

We add the inequalities (2.61) and obtain, by (2.64)

$$\int_{\alpha_P} e^{v_1} |dz| \leq \frac{32}{3} \int_{\beta} e^{w_1} |dz| + \frac{256}{3}. \quad (2.65)$$

(2.53) follows from (2.54) and (2.65). Since (2.26) is implied by (2.53), (2.39) and (2.52), this completes the proof of Lemma 3. Theorem 1 is a consequence of Lemmas 2 and 3.

**Proof of Theorem 3.** If there exists a subcompact of  $\Omega_0$  which intersects infinitely many  $\gamma_n$ , then Theorem 3 follows immediately from an application of Lemma 2. Otherwise we may suppose that (eventually after extraction of a subsequence) the interior regions of  $\{\gamma_n\}$  tend monotonically to  $\Omega_0$ .

We distinguish between two cases depending on whether  $\Omega_0$  is parabolic (i. e. identical with the entire plane) or hyperbolic.

(I) *Let  $\Omega_0$  be parabolic.* If  $\Phi(\Gamma) < -1$ , then Theorem 3 is superseded by Theorem 1. We complete the proof of Theorem 3 for parabolic  $\Omega_0$  by showing that in the remaining case the hypotheses are incompatible:

**Lemma 4.** *Suppose that  $\Omega_0$  is the entire plane and that  $\Phi(\Gamma) > -1$ . Let  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , be a sequence of rectifiable JORDAN curves whose interior regions tend increasingly to  $\Omega_0$ . Then*

$$\lim_{n \rightarrow \infty} \int_{\gamma_n} e^{u(z)} |dz| = +\infty. \quad (2.66)$$

**Proof.** There exists an index  $N$  and a number  $\eta > 0$  such that

$$\Phi(u_1, \gamma_n; \Gamma) - \Phi_2(\Gamma) < -1 + \eta \quad (2.67)$$

for all  $n \geq N$ . We choose a circle  $\gamma = [|z| = R]$  which encloses  $\gamma_N$ . Let  $\omega_n$  denote the annular region bounded by  $\gamma$  and  $\gamma_n$ ,  $n$  being arbitrary but sufficiently large so that  $|z| \leq R$  lies within  $\gamma_n$ . The theory of F. RIESZ [27] implies

$$\frac{1}{2\pi} \int_{\delta} \frac{\partial h_1}{\partial n} |dz| \geq \Phi(u_1, \gamma; \Gamma) \quad (2.68)$$

and

$$\frac{1}{2\pi} \int_{\delta} \frac{\partial h_2}{\partial n} |dz| \leq \Phi(u_2, \gamma_n; \Gamma) \quad (2.69)$$

for every smooth JORDAN curve  $\delta$  in  $\omega_n$  which encloses  $\gamma$ . Here  $h_1$  and  $h_2$  denote the best harmonic majorants of  $u_1$  and  $u_2$ , respectively, in  $\omega_n$ , and  $n$  designates the outer normal. We define

$$h(z) = h_1(z) - h_2(z) \quad (2.70)$$

and conclude from relations (2.67) to (2.70) that

$$\frac{1}{2\pi} \int_{\delta} \frac{\partial h}{\partial n} |dz| > -1 + \eta. \quad (2.71)$$

There exists a (uniquely determined) conformal representation  $z = \varphi_n(\zeta)$  of a (suitable) circular ring  $R < |\zeta| < R_n$  onto  $\omega_n$  such that  $\varphi_n(+R) = +R$ . We introduce the function

$$\tilde{h}_n(\zeta) = h(\varphi_n(\zeta)) + \log |\varphi_n'(\zeta)|.$$

$\tilde{h}_n$  is harmonic in  $R < |\zeta| < R_n$  and we have

$$\int_{\gamma} \frac{\partial \tilde{h}_n}{\partial n} |d\zeta| = \int_{\delta} \frac{\partial h}{\partial n} |dz|, \quad (2.72)$$

$\gamma$  and  $\delta$  denoting arbitrary smooth JORDAN curves, not null-homotopic and lying in  $R < |\zeta| < R_n$  and  $\omega_n$ , respectively. We state that

$$\begin{aligned} \int_{-\pi}^{+\pi} \tilde{h}_n(\varrho_2 e^{i\varphi}) d\varphi &= \int_{-\pi}^{+\pi} \tilde{h}_n(\varrho_1 e^{i\varphi}) d\varphi + 2\pi \int_{\varrho_1}^{\varrho_2} \left[ \frac{1}{2\pi} \int_{-\pi}^{+\pi} \frac{\partial \tilde{h}_n(\varrho e^{i\varphi})}{\partial \varrho} \varrho d\varphi \right] \frac{d\varrho}{\varrho} \\ &\geq \int_{-\pi}^{+\pi} \tilde{h}_n(\varrho_1 e^{i\varphi}) d\varphi + 2\pi(-1 + \eta)(\log \varrho_2 - \log \varrho_1) \\ &> A + 2\pi(-1 + \eta) \log \varrho_2 \quad (R < \varrho_1 < \varrho_2 < R_n), \end{aligned} \quad (2.73)$$

where  $A$  is a constant not depending on  $n$ ,  $\varrho_1$  and  $\varrho_2$ . In this inequality we first apply (2.71) and (2.72). Then we let  $\varrho_1 \rightarrow R$ . We complete the proof of (2.73) by showing that the limits

$$L_n = \lim_{\varrho_1 \rightarrow R} \int_{-\pi}^{+\pi} \tilde{h}_n(\varrho_1 e^{i\varphi}) d\varphi \quad (n = 1, 2, 3, \dots)$$

exist and are uniformly bounded. To this end we consider the function

$$U_1(z) = \begin{cases} h_1(z) & \text{in } \omega_n \\ u_1(z) & \text{on } \gamma \cup (\Delta, \gamma), \end{cases}$$

which is defined and subharmonic throughout the annulus  $(\Delta, \gamma_n)$ . The conformally transplanted functions,  $\tilde{U}_{1n}(\zeta) = U_1(\varphi_n(\zeta))$ ,  $n = 1, 2, 3, \dots$ , are subharmonic in circular rings  $R'_n < |\zeta| < R_n$  ( $R'_n < R$ ). It is well known (see e. g. [25, p. 8]) that the arithmetic mean of a subharmonic function on concentric circles is a continuous function of the radius. Consequently

$$\begin{aligned} \lim_{\varrho_1 \rightarrow R} \frac{1}{2\pi} \int_{-\pi}^{+\pi} h_1(\varphi_n(\varrho_1 e^{i\varphi})) d\varphi &= \lim_{\varrho_1 \rightarrow R} \frac{1}{2\pi} \int_{-\pi}^{+\pi} \tilde{U}_{1n}(\varrho_1 e^{i\varphi}) d\varphi \\ &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \tilde{U}_{1n}(R e^{i\varphi}) d\varphi = \frac{1}{2\pi} \int_{-\pi}^{+\pi} u_1(\varphi_n(R e^{i\varphi})) d\varphi. \end{aligned}$$

An analogous relation holds with index 2. By subtraction we obtain

$$\lim_{\varrho_1 \rightarrow R} \int_{-\pi}^{+\pi} h(\varphi_n(\varrho_1 e^{i\varphi})) d\varphi = \int_{-\pi}^{+\pi} u(\varphi_n(R e^{i\varphi})) d\varphi. \quad (2.74)$$

Furthermore

$$\lim_{\varrho_1 \rightarrow R} \int_{-\pi}^{+\pi} \log |\varphi'_n(\varrho_1 e^{i\varphi})| d\varphi = \int_{-\pi}^{+\pi} \log |\varphi'_n(R e^{i\varphi})| d\varphi, \quad (2.75)$$

since  $\varphi_n$  is analytic on  $|\zeta| = R$ . From (2.74), (2.75) and the definition of  $\tilde{h}_n$  we infer that

$$L_n = \int_{-\pi}^{+\pi} u(\varphi_n(R e^{i\varphi})) d\varphi + \int_{-\pi}^{+\pi} \log |\varphi'_n(R e^{i\varphi})| d\varphi \quad (n = 1, 2, 3, \dots). \quad (2.76)$$

We have yet to verify that the sequence  $\{L_n\}$  is bounded. Actually we shall prove more, namely

$$\lim_{n \rightarrow \infty} L_n = \int_{-\pi}^{+\pi} u(Re^{i\varphi}) d\varphi. \quad (2.77)$$

we first observe that (2.77) will be an immediate consequence of (2.76) after it has been shown that

$$\lim_{n \rightarrow \infty} \varphi_n(\zeta) = \zeta \quad (2.78)$$

(and, consequently,  $\lim_{n \rightarrow \infty} \varphi'_n(\zeta) \equiv 1$ ), uniformly on  $|\zeta| = R$ . Indeed, then

$$\lim_{n \rightarrow \infty} \int_{-\pi}^{+\pi} u(\varphi_n(Re^{i\varphi})) d\varphi = \lim_{n \rightarrow \infty} \int_{-\pi}^{+\pi} u(Re^{i\psi}) \left| \frac{d\varphi_n^{-1}(Re^{i\psi})}{dz} \right| d\psi = \int_{-\pi}^{+\pi} u(Re^{i\psi}) d\psi,$$

whereas the second term on the right-hand side of (2.76) tends to 0 as  $n \rightarrow \infty$ .

In order to prove (2.78) we first extend the definition of  $\varphi_n(\zeta)$  by reflection at  $|\zeta| = R$ . The resulting function is schlicht in  $R^2/R_n < |\zeta| < R_n$ , and it maps  $|\zeta| = R$  onto  $|z| = R$ . We mention that  $R_n \rightarrow \infty$  as  $n \rightarrow \infty$ , due to the fact that  $\omega_n$  tends to  $R < |z| < +\infty$ .

By making use of CANTOR's diagonal process we select a subsequence  $\{\varphi_n^*\}$  of  $\{\varphi_n\}$  which converges at an enumerable set of points possessing limit points in both  $0 < |\zeta| < R$  and  $R < |\zeta| < +\infty$ .  $\{\varphi_n^*\}$  is normal [7, pp. 176, 179] in these regions. Hence, by a theorem of VITALI [7, p. 186], it converges there, uniformly on every subcompact. From this we conclude that  $\{\varphi_n^*\}$  converges even in  $0 < |\zeta| < +\infty$ , uniformly on every subcompact. The limit function  $\varphi$  is either schlicht meromorphic or a constant [7, p. 193]. The latter possibility can immediately be excluded, since convergence to a constant, uniformly on  $|\zeta| = R$ , is incompatible with the fact that  $\varphi_n$  maps  $|\zeta| = R$  onto  $|z| = R$  for all  $n$ .

The image region has to be of the same conformal type as  $0 < |\zeta| < +\infty$ . Furthermore, it follows from results of A. HURWITZ [7, pp. 191–192] that  $\varphi$  does not assume the values 0 and  $\infty$ . Hence  $0 < |\zeta| < +\infty$  is mapped onto  $0 < |z| < +\infty$  and  $\varphi$  is necessarily a linear transformation. The points 0,  $\infty$  and  $+R$  are fixed. Consequently,  $\varphi(\zeta) \equiv \zeta$ .

Finally, if  $\{\varphi_n(\zeta)\}$  does not converge to  $\zeta$ , then  $\{\varphi_n^*\}$  can be chosen such that, for some  $\zeta_0$  ( $0 < |\zeta_0| < +\infty$ ),  $\lim_{n \rightarrow \infty} \varphi_n^*(\zeta_0) = \zeta_1 \neq \zeta_0$ . We thus obtain a contradiction, since the above reasoning yields  $\lim_{n \rightarrow \infty} \varphi_n^*(\zeta_0) = \zeta_0$ . This completes the proof of (2.78) and, with it, of (2.73).

By the theorem of the arithmetic and geometric means [15, p. 137]

$$\int_{-\pi}^{+\pi} \exp \{ \tilde{h}_n(\varrho_2 e^{i\varphi}) \} \varrho_2 d\varphi \geq 2\pi \varrho_2 \exp \left\{ \frac{1}{2\pi} \int_{-\pi}^{+\pi} \tilde{h}_n(\varrho_2 e^{i\varphi}) d\varphi \right\}. \quad (2.79)$$

Furthermore

$$\limsup_{\varrho_2 \rightarrow R_n - \pi} \int_{-\pi}^{+\pi} \exp \{ \tilde{h}_n(\varrho_2 e^{i\varphi}) \} \varrho_2 d\varphi \leq \int_{\gamma_n} e^u |dz| . \quad (2.80)$$

This relation will be verified in the proof of Lemma 5 (see formula (2.130)). By (2.73), (2.79) and (2.80)

$$\int_{\gamma_n} e^u |dz| \geq 2\pi e^{A/2\pi} R_n^\eta . \quad (2.81)$$

(2.66) is implied by (2.81), since  $R_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

(II) *Let  $\Omega_0$  be hyperbolic.* Again we may suppose that the interior regions of  $\{\gamma_n\}$  tend increasingly to  $\Omega_0$ . Furthermore, we assume that  $\Phi(\Gamma) = +\infty$ , since otherwise Theorem 3 is superseded by Theorem 2. Then there exists an index  $N$  such that

$$\Phi(u_1, \gamma_N; \Gamma) - \Phi_2(\Gamma) > 0 \quad (2.82)$$

and

$$\Phi_2(\Gamma) - \Phi(u_2, \gamma_N; \Gamma) < \frac{1}{16} . \quad (2.83)$$

Let  $n > N$  be arbitrary. Theorem 3 (for the case remaining to be treated) is an obvious consequence of Lemma 2 and the following result:

**Lemma 5.** *There exists a rectifiable curve  $\sigma_n$ , leading from  $\gamma_N$  to  $\gamma_n$ , such that*

$$\int_{\sigma_n} e^u |dz| \leq \frac{M}{\sin(\pi/8) \cos(\pi/32)} . \quad (2.84)$$

**Proof.** By F. RIESZ [27, II, p. 357] the representations

$$u_1(z) = h_1(z) - \int_{\omega} g(z, \zeta) d\mu_1(e_\zeta) \quad (2.85)$$

and

$$u_2(z) = h_2(z) - \int_{\omega} g(z, \zeta) d\mu_2(e_\zeta) \quad (2.86)$$

hold in the annulus  $\omega = (\gamma_N, \gamma_n)$ . Here  $h_1$  and  $h_2$  denote the best harmonic majorants of  $u_1$  and  $u_2$ , respectively, in  $\omega$  and  $g$  is GREEN's function for this region. The theory of F. RIESZ implies further that

$$\Phi(u_1, \gamma_N; \Gamma) \leq \frac{1}{2\pi} \int_{\gamma} \frac{\partial h_1}{\partial n} |dz| \leq \Phi(u_1, \gamma_n; \Gamma) \quad (2.87)$$

and

$$\Phi(u_2, \gamma_N; \Gamma) \leq \frac{1}{2\pi} \int_{\gamma} \frac{\partial h_2}{\partial n} |dz| \leq \Phi(u_2, \gamma_n; \Gamma) \quad (2.88)$$

for every smooth JORDAN curve  $\gamma$  in  $\omega$  which encloses  $\gamma_N$ .  $n$  denotes the outer

normal. We define

$$h(z) = h_1(z) - h_2(z) \quad (2.89)$$

and introduce the abbreviation

$$\alpha = \frac{1}{2\pi} \int_{\gamma} \frac{\partial h}{\partial n} |dz| . \quad (2.90)$$

$\alpha$  does not depend on the choice of  $\gamma$ . (2.85), (2.86) and (2.89) imply

$$u(z) \leq h(z) + \int_{\omega} g(z, \zeta) d\mu_2(e_{\zeta}) . \quad (2.91)$$

From (2.83) follows

$$\mu_2(\omega) < \frac{1}{16} . \quad (2.92)$$

We begin with the special case where  $\gamma_N$  and  $\gamma_n$  are circles  $|z| = R_N$  and  $|z| = R_n$  ( $0 < R_N < R_n < +\infty$ ). We state that

$$\int_{R_N}^{R_n} e^{u(x)} dx \leq \frac{M}{\sin(\pi/8) \cos(\pi/32)} . \quad (2.93)$$

More generally, (2.93) holds if the integration is extended over any radial segment  $(R_N e^{i\Theta}, R_n e^{i\Theta})$ ,  $0 \leq \Theta < 2\pi$ . First we prove (2.93) under the additional assumption that the measure  $\mu_2(e_{\zeta})$  is concentrated in one point  $\zeta_0$ . We have, for positive real  $x$

$$g(x, \zeta_0) \leq g(x, |\zeta_0|) . \quad (2.94)$$

This estimation can be verified by considering, for fixed  $x$ , the FOURIER expansion of  $g(x, \zeta_0)$  on the circle  $|\zeta_0| = \text{const}$ . We conclude from (2.91), (2.92) and (2.94) that, for positive real  $x$

$$u(x) \leq h(x) + \frac{1}{16} g(x, |\zeta_0|) . \quad (2.95)$$

If we slit  $\omega$  along the negative real axis we obtain a simply connected region  $\omega'$ . We define

$$f(z) = \exp \{ (h(z) + i h^*(z)) / 2 \} , \quad (2.96)$$

where  $h^*$  is conjugate harmonic to  $h$  and, for example,  $h^*((R_N + R_n)/2) = 0$ .  $f$  is regular analytic in  $\omega'$ . It can be continued into  $\omega$  as a multiple-valued function with (multiplicative) period  $e^{\pi\alpha i}$ . We consider

$$\varphi(z) = z^{-\alpha/2} f(z) , \quad (2.97)$$

defining  $z^{-\alpha/2}$  to be positive real on the interval  $(R_N, R_n)$ .  $\varphi$  is analytic in  $\omega'$

and can be continued into  $\omega$  in a unique way. There exists a decomposition of the form

$$\varphi(z) = \varphi_1(z) + i\varphi_2(z) , \quad (2.98)$$

where  $\varphi_1$  and  $\varphi_2$  are analytic and one-valued in  $\omega$ , real for real  $z$ . Indeed, just put  $\varphi_1(z) = \Sigma a_k z^k$  and  $\varphi_2(z) = \Sigma b_k z^k$ , where  $\Sigma(a_k + ib_k)z^k$  is the LAURENT expansion of  $\varphi$  in  $\omega$  ( $a_k, b_k$  real;  $k = 0, \pm 1, \pm 2, \dots$ ). From (2.97) and (2.98) we conclude that  $f$  admits the decomposition

$$f(z) = f_1(z) + if_2(z) , \quad (2.99)$$

where  $f_1$  and  $f_2$  are analytic in  $\omega'$  and

$$\arg f_1 \equiv \arg f_2 \equiv 0 \pmod{\pi} \quad (2.100)$$

on  $(R_N, R_n)$ . Furthermore, on  $(-R_N, -R_n)$  we have

$$\arg f_1 \equiv \arg f_2 \equiv \pi\alpha/2 \pmod{\pi} \quad (2.101)$$

or

$$\arg f_1 \equiv \arg f_2 \equiv -\pi\alpha/2 \pmod{\pi} \quad (2.102)$$

depending on whether the real axis is approached from above or below. For real  $z$

$$|f(z)|^2 = |f_1(z)|^2 + |f_2(z)|^2 . \quad (2.103)$$

We further have, for  $-\pi < t < +\pi$  and  $R_N < r < R_n$

$$|f(re^{it})|^2 = |f_1(re^{it})|^2 + |f_2(re^{it})|^2 + i[f_1(re^{-it})f_2(re^{it}) - f_1(re^{it})f_2(re^{-it})] .$$

Since the bracket contains an odd function of  $t$ , it follows that

$$\int_{-\pi}^{+\pi} |f(re^{it})|^2 dt = \int_{-\pi}^{+\pi} (|f_1(re^{it})|^2 + |f_2(re^{it})|^2) dt \quad (2.104)$$

for  $R_N < r < R_n$ . Let  $\omega'' = \omega \cap [\operatorname{Im} z > 0]$ . Let  $g^*(z, |\zeta_0|)$  be conjugate harmonic to  $g(z, |\zeta_0|)$  in  $\omega''$ , satisfying

$$g^*(x, |\zeta_0|) = 0 \quad \text{for} \quad R_N < x < |\zeta_0| . \quad (2.105)$$

We then have

$$g^*(x, |\zeta_0|) = \pi \quad \text{for} \quad |\zeta_0| < x < R_n \quad (2.106)$$

and

$$0 < g^*(x, |\zeta_0|) < \pi \quad (2.107)$$

throughout  $\omega''$ . We define

$$F_1(z) = f_1^2(z) \exp \left\{ \frac{1}{16} [g(z, |\zeta_0|) + ig^*(z, |\zeta_0|)] \right\} . \quad (2.108)$$

We state that

$$\begin{aligned} & \sin(\pi/8) \cos(\pi/32) \int_{r_N}^{r_n} |f_1(x)|^2 \exp \left\{ \frac{1}{16} g(x, |\zeta_0|) \right\} dx \\ & \leq r_N \exp \left\{ \frac{1}{16} g(r_N, |\zeta_0|) \right\} \int_0^\pi |f_1(r_N e^{it})|^2 dt + r_n \exp \left\{ \frac{1}{16} g(r_n, |\zeta_0|) \right\} \int_0^\pi |f_1(r_n e^{it})|^2 dt \end{aligned} \quad (2.109)$$

for any two radii  $r_N$  and  $r_n$  ( $R_N < r_N < r_n < R_n$ ). For the proof of this relation we distinguish between two cases:

(A)  $|\alpha - q| \geq 1/4$ , where  $q$  denotes the or one of the odd integers nearest to  $\alpha$ .

(B) There exists an odd integer  $q$  such that  $|\alpha - q| < 1/4$ .

*Case (A):* We apply CAUCHY's integral theorem to  $F_1$ . We integrate along the boundary of  $\omega''$  (described in the positive sense), replacing the circular boundary by the approximating circles  $|z| = r_N$  and  $|z| = r_n$ . At first we bypass  $\zeta_0$  on a small semicircle of radius  $\varepsilon$ . But we observe that the integral along this semicircle tends to 0 with  $\varepsilon$ . So we have

$$I_1 + I_2 + I_3 + I_4 = 0, \quad (2.110)$$

where  $I_1, I_2, I_3$  and  $I_4$  denote the respective integrals along  $(r_N, |\zeta_0|)$ ,  $(|\zeta_0|, r_n)$ ,  $(-r_n, -r_N)$  and the two semicircles. (2.100), (2.105) and (2.108) imply

$$\arg I_1 = 0. \quad (2.111)$$

By (2.100), (2.106) and (2.108) we have

$$\arg I_2 = \pi/16. \quad (2.112)$$

From (2.111) and (2.112) we infer

$$|I_1 + I_2| \geq \cos(\pi/32)(|I_1| + |I_2|). \quad (2.113)$$

We further have

$$0 \leq \arg(I_1 + I_2) \leq \pi/16. \quad (2.114)$$

From the assumption that  $|\alpha - q| \geq 1/4$  ( $q$  denoting the or one of the odd integers nearest to  $\alpha$ ), (2.101), (2.107) and (2.108) we conclude that  $\arg I_3$  distinguishes itself from the nearest odd multiple of  $\pi$  by at least  $3\pi/16$ . Consider the triangle with sides  $I_1 + I_2$ ,  $I_3$  and  $I_4$  in the complex number plane. Since the angle between  $I_1 + I_2$  and  $I_3$  is  $\geq \pi/8$  we obtain the estimation

$$|I_4| \geq \sin(\pi/8)|I_1 + I_2|. \quad (2.115)$$

From (2.113), (2.115) and the property (2.94) of Green's function it follows indeed that

$$\begin{aligned}
& \sin(\pi/8) \cos(\pi/32) \int_{r_N}^{r_n} |f_1(x)|^2 \exp \left\{ \frac{1}{16} g(x, |\zeta_0|) \right\} dx \\
&= \sin(\pi/8) \cos(\pi/32) (|I_1| + |I_2|) \leq |I_4| \\
&\leq r_N \exp \left\{ \frac{1}{16} g(r_N, |\zeta_0|) \right\} \int_0^\pi |f_1(r_N e^{it})|^2 dt \\
&+ r_n \exp \left\{ \frac{1}{16} g(r_n, |\zeta_0|) \right\} \int_0^\pi |f_1(r_n e^{it})|^2 dt .
\end{aligned}$$

*Case (B):* From (2.82), (2.87), (2.88), (2.90) and the inequality  $|\alpha - q| < 1/4$  we conclude that  $q > 0$ .

We first make the additional hypothesis that  $f_1 \neq 0$  on

$$D = [r_N \leq |z| \leq r_n] \cap [\operatorname{Im} z \geq 0] .$$

Our method is again based on an application of CAUCHY's integral theorem to the function  $F_1$ , but this time we do not integrate along the boundary of  $D$ . (The previous argument breaks down because it does not yield any more an inequality of the type (2.115)).

We state that there exists an *analytic curve*  $\tau$  with the following properties:

(a)  $\tau$  is contained in  $D$  and leads from  $|z| = r_n$  to  $|z| = r_N$ , (b)  $\arg[f_1^2(z)dz] \equiv 0$  along  $\tau$ .

In order to verify the existence of  $\tau$  we decompose the function  $\log f_1^2$  into its real and imaginary parts,  $\log f_1^2 = H_1(z) + iH_1^*(z)$ . With every point  $z$  on  $D$  we associate the unit vector  $\exp\{-iH_1^*(z)\}$ . The thus defined vector field has no singular points, since  $f_1 \neq 0$  on  $D$ . Through every point  $z$  on  $D$  there passes exactly one streamline (i. e. solution of the differential equation

$$dz/dt = \exp\{-iH_1^*(z)\} ,$$

which begins and ends on the boundary of  $D$ . Obviously,  $\arg[f_1^2(z)dz] \equiv 0$  along these lines. They are analytic since the conformal mapping  $w = \int_{z_0}^z f_1^2(z) dz$  transforms them into straight lines. So the existence of  $\tau$  will be asserted if we can verify that at least one of these streamlines leads from  $|z| = r_n$  to  $|z| = r_N$ .

On the intervals  $(r_N, r_n)$  and  $(-r_n, -r_N)$ ,  $H_1^* = 0$  and  $H_1^* = \pi\alpha$ , respectively. Since  $q \geq 1$ ,  $\alpha > \frac{3}{4}$ . The continuity of  $H_1^*$  and the relations  $H_1^*(r_N) = 0$ ,  $H_1^*(-r_N) = \pi\alpha > \frac{3}{4}\pi$  imply: There exists at least one open sub-arc  $S = [\Theta_1 > \Theta = \arg z > \Theta_2]$  of the semicircle

$$C_N = [|z| = r_N] \cap [\operatorname{Im} z \geq 0]$$

with the following properties:

- (a)  $H_1^*(r_N e^{i\theta_1}) \equiv -\Theta_1 + \frac{3\pi}{2} \pmod{2\pi}$ ,
- (b)  $H_1^*(r_N e^{i\theta_2}) \equiv -\Theta_2 + \frac{\pi}{2} \pmod{2\pi}$ ,
- (c)  $-\Theta + \frac{3\pi}{2} > H_1^*(r_N e^{i\theta}) > -\Theta + \frac{\pi}{2} \pmod{2\pi}$

for all  $\Theta$  in  $S$ .

Since  $H_1^*(r_N e^{i\theta})$  is of bounded variation in  $\Theta$ , there exists but a finite number of such arcs. We denote them by  $S_1, S_2, \dots, S_m$  and suppose that they have been arranged in such a way that  $\Theta_{k-1,2} > \Theta_{k1}$  ( $k = 2, 3, \dots, m$ ), where  $S_k = [\Theta_{k1} > \Theta > \Theta_{k2}]$ . In each  $S_k$  there is (at least) one point  $\Theta_{k0}$  at which  $H_1^*(r_N e^{i\theta_{k0}}) = -\Theta_{k0} + \pi \pmod{2\pi}$ .

The geometrical meaning of these conditions is the following: At  $r_N e^{i\theta_{k1}}$  and at  $r_N e^{i\theta_{k2}}$  the field vector is tangential to  $C_N$ , directed away from  $S_k$ . It points into  $|z| < r_N$  at all points in  $S_k$  and is, in particular, normal to  $C_N$  at  $r_N e^{i\theta_{k0}}$ .

We are now going to prove that at least one of the  $m$  streamlines ending at  $r_N e^{i\theta_{10}}, r_N e^{i\theta_{20}}, \dots, r_N e^{i\theta_{m0}}$ , must begin on  $|z| = r_n$ . This will complete the existence proof for  $\tau$ .

We state that  $H_1^*(r_N e^{i\theta_{10}}) \geq \pi q - \Theta_{10}$ . Indeed, otherwise

$$H_1^*(r_N e^{i\theta_{10}}) \leq \pi(q - 2) - \Theta_{10}.$$

From this would follow

$$H_1^*(r_N e^{i\theta_{11}}) \leq \pi\left(q - \frac{3}{2}\right) - \Theta_{11},$$

which, in turn, would imply the existence of at least one  $S_k$  between  $-r_N$  and  $r_N e^{i\theta_{11}}$ , contrary to hypothesis.

Let  $z_0$  be the point at which the streamline  $\gamma$  ending at  $r_N e^{i\theta_{10}}$  begins. We consider all possibilities:

- (a)  $|z_0| = r_n$ . Then there is nothing left to prove.
- (b)  $z_0$  positive real. This never occurs because the positive real axis is itself a streamline.
- (c)  $z_0 = r_N e^{i\theta_0}$  ( $\Theta_{10} > \Theta_0 > 0$ ). It is easy to verify that  $\arg dz$  increases by at least  $\frac{\pi}{2} + \Theta_{10} - \Theta_0$  if we follow  $\gamma$  from  $z_0$  to  $r_N e^{i\theta_{10}}$ . From this, the inequality  $H_1^*(r_N e^{i\theta_{10}}) \geq \pi q - \Theta_{10}$  and the fact that  $\arg[f_1^2(z)dz] \equiv 0$  along  $\gamma$  we conclude that  $H_1^*(z_0) \geq \pi(q + \frac{1}{2}) - \Theta_0$ . Hence, there exists at

least one more  $S_k$  between  $z_0$  and  $r_N$ . Furthermore, we have again

$$H_1^*(r_N e^{i\Theta_{k_0}}) \geq \pi q - \Theta_{k_0} .$$

(d)  $z_0 = r_N e^{i\Theta_0}$  ( $\pi > \Theta_0 > \Theta_{10}$ ). It is easy to see that  $\arg dz$  decreases by at least  $\frac{\pi}{2} + \Theta_0 - \Theta_{10}$  if we follow  $\gamma$  from  $z_0$  to  $r_N e^{i\Theta_{10}}$ . Hence

$$H_1^*(r_N e^{i\Theta_{10}}) - H_1^*(z_0) \geq \frac{\pi}{2} + \Theta_0 - \Theta_{10} .$$

Furthermore,  $H_1^*(z_0) \geq \pi(q + \frac{1}{2}) - \Theta_0$ . (Otherwise, since the vector  $\exp\{-iH_1^*(z_0)\}$  points into  $D$ , we would have  $H_1^*(z_0) \leq \pi(q - \frac{1}{2}) - \Theta_0$ . But this would imply the existence of an  $S_k$  between  $-r_N$  and  $z_0$ , contrary to hypothesis). We conclude that  $H_1^*(r_N e^{i\Theta_{10}}) \geq \pi(q + 1) - \Theta_{10}$ . But, knowing that  $H_1^*(r_N e^{i\Theta_{10}}) + \Theta_{10}$  is an odd multiple of  $\pi$ , we infer that even  $H_1^*(r_N e^{i\Theta_{10}}) \geq \pi(q + 2) - \Theta_{10}$  is satisfied. Consequently,

$$H_1^*(r_N e^{i\Theta_{12}}) \geq \pi\left(q + \frac{3}{2}\right) - \Theta_{12} .$$

So there must exist a second  $\text{arc} S_2$  and we have  $H_1^*(r_N e^{i\Theta_{20}}) \geq \pi q - \Theta_{20}$ .

(e)  $z_0$  negative real. The previous argument applies also to this case and yields the same conclusion.

As a result of this discussion we have now the following alternative: Either (a) occurs, and then the proof is completed, or the above constructions lead to another  $S_k$  such that  $H_1^*(r_N e^{i\Theta_{k_0}}) \geq \pi q - \Theta_{k_0}$ . In the second case we repeat the above reasoning. (The reader will convince himself that this can be done without difficulty. The following trivial observation is useful in this connection: Streamlines do not intersect. Hence, for example, the one ending at  $r_N e^{i\Theta_{k_0}}$  does not begin on  $|z| = r_N$  between the points  $z_0$  and  $r_N e^{i\Theta_{10}}$ .) We arrive at the same alternative again. But there are only a finite number of  $S_k$ 's. Hence, if we iterate this argument we must meet with case (a) after a finite number of steps. So  $\tau$  exists.

We now apply CAUCHY's integral theorem to  $F_1$ , integrating in the positive sense along the closed curve consisting of  $\tau$ ,  $(r_N, r_n)$  and subarcs of  $|z| = r_N$  and  $|z| = r_n$ . We have

$$I_1 + I_2 + I_3' + I_4' = 0 , \quad (2.116)$$

where  $I_1, I_2, I_3'$  and  $I_4'$  denote the respective integrals along  $(r_N, |\zeta_0|)$ ,  $(|\zeta_0|, r_n)$ ,  $\tau$  and the two connecting circular arcs. From (2.107) and the fact that  $\arg [f_1^2(z) dz] \equiv 0$  along  $\tau$  we infer

$$0 \leq \arg I_3' \leq \pi/16 . \quad (2.117)$$

Consider the triangle with sides  $I_1 + I_2$ ,  $I'_3$  and  $I'_4$  in the complex number plane. By (2.114) and (2.117), the angle between  $I_1 + I_2$  and  $I'_3$  is  $\geq 15\pi/16$ . Hence

$$|I'_4| \geq |I_1 + I_2| . \quad (2.118)$$

It follows from (2.113), (2.118) and the property (2.94) of GREEN's function that

$$\begin{aligned} \cos(\pi/32) \int_{r_N}^{r_n} |f_1(x)|^2 \exp\left\{\frac{1}{16}g(x, |\zeta_0|)\right\} dx &= \cos(\pi/32)(|I_1| + |I_2|) \leq |I'_4| \\ &\leq r_N \exp\left\{\frac{1}{16}g(r_N, |\zeta_0|)\right\} \int_0^\pi |f_1(r_N e^{it})|^2 dt + r_n \exp\left\{\frac{1}{16}g(r_n, |\zeta_0|)\right\} \int_0^\pi |f_1(r_n e^{it})|^2 dt . \end{aligned}$$

This inequality implies (2.109). It has been deduced under the assumption that  $f_1 \neq 0$  on  $D$ . We now admit zeros, but still assume that  $f_1 \neq 0$  on the circles  $|z| = r_N$  and  $|z| = r_n$ . Let  $a_1, a_2, \dots, a_s$  denote the zeros of  $f_1$  on  $D$ . We introduce the function

$$f_{10}(z) = f_1(z) \prod_{i=1}^s \exp\{[g(z, a_i) + g(z, \bar{a}_i)] + i[g^*(z, a_i) + g^*(z, \bar{a}_i)]\} . \quad (2.119)$$

Here  $g$  denotes GREEN's function for the annulus  $r_N < |z| < r_n$  and  $g^*$  is conjugate harmonic to  $g$ , defined to vanish at  $z = r_N$ .  $f_{10}$  is analytic and  $\neq 0$  on  $D$ . Obviously

$$|f_{10}(z)| = |f_1(z)| \quad (2.120)$$

on  $|z| = r_N$  and on  $|z| = r_n$ . Furthermore

$$|f_{10}(z)| \geq |f_1(z)| , \quad (2.121)$$

everywhere on  $D$ . We conclude from (2.100) and (2.119) that

$$\arg f_{10} \equiv 0 \pmod{\pi} \quad (2.122)$$

(2.101) and (2.119) imply

$$\arg f_{10} \equiv \pi\alpha_0/2 \pmod{\pi} , \quad (2.123)$$

where  $\alpha_0 > \alpha$ . We can write  $\alpha_0 = q_0 + \vartheta_0$ , where  $q_0$  is a positive odd integer and  $|\vartheta_0| \leq 1$ . We distinguish between two cases:

(B<sub>1</sub>)  $|\vartheta_0| < 1/4$ . The above reasoning can be applied to  $f_{10}(z)$  since this function has no zeros on  $D$ . We obtain relation (2.109), but with  $f_{10}$  taking the place of  $f_1$ . It follows from (2.120) and (2.121) that the unmodified inequality (2.109) is true a fortiori.

(B<sub>2</sub>)  $|\vartheta_0| \geq 1/4$ . By the method used in case (A) we prove (2.109),  $f_1$  being replaced by  $f_{10}$ . The unmodified inequality (2.109) follows as mentioned.

We can easily free ourselves from the assumption that  $f_1 \neq 0$  on the circles  $|z| = r_N$  and  $|z| = r_n$ . For, if this hypothesis should not be fulfilled, then we first prove (2.109) for neighboring circles and afterwards pass to the limit.

(2.109) is also satisfied by  $f_2$ . Analogous estimations hold for the lower half-annulus. By adding these four inequalities and making use of (2.95), (2.96), (2.103) and (2.104) we obtain

$$\begin{aligned} 2 \sin(\pi/8) \cos(\pi/32) \int_{r_N}^{r_n} e^{u(x)} dx &\leq r_N \exp \left\{ \frac{1}{16} g(r_N, |\zeta_0|) \right\} \int_{-\pi}^{+\pi} |f(r_N e^{it})|^2 dt \\ &+ r_n \exp \left\{ \frac{1}{16} g(r_n, |\zeta_0|) \right\} \int_{-\pi}^{+\pi} |f(r_n e^{it})|^2 dt . \end{aligned} \quad (2.124)$$

The best harmonic majorants  $h_1$  and  $h_2$  are limits of decreasing sequences,  $\{h_{1k}\}$  and  $\{h_{2l}\}$  ( $k, l = 1, 2, 3, \dots$ ), consisting of functions which are harmonic in the region  $\omega$ , continuous on its closure. Furthermore, on the boundary  $h_{1k} \searrow u_1$  and  $h_{2l} \searrow u_2$  for  $k, l \nearrow \infty$ .

Let  $k, l$  be fixed. The function  $|z| \exp \{h_{1k}(z) - h_{2l}(z)\}$  is subharmonic in  $\omega$ . Hence the integral

$$L(r \exp \{h_{1k} - h_{2l}\}; r) = \int_{-\pi}^{+\pi} r \exp \{h_{1k}(re^{it}) - h_{2l}(re^{it})\} dt$$

is a convex function of  $\log r$ . Consequently, for  $R_N < r < R_n$

$$\begin{aligned} L(r \exp \{h_{1k} - h_{2l}\}; r) &\leq \max [L(r \exp \{h_{1k} - h_{2l}\}; R_N), \\ &L(r \exp \{h_{1k} - h_{2l}\}; R_n)] . \end{aligned}$$

Letting first  $k \rightarrow \infty$ , then  $l \rightarrow \infty$ , we obtain, by hypothesis (b) of Theorem 3

$$L(r|f|^2; r) = L(re^h; r) \leq \max [L(re^u; R_N), L(re^u; R_n)] < M .$$

Since, furthermore,  $g(r_N, |\zeta_0|) \rightarrow 0$  for  $r_N \rightarrow R_N$  and  $g(r_n, |\zeta_0|) \rightarrow 0$  for  $r_n \rightarrow R_n$ , (2.93) is a consequence of (2.124).

Let us now proceed to general measures  $\mu_2$ . We define

$$S_K = \int_{R_N}^{R_n} \exp \{h^K(x) + \int_{\omega} g^K(x, \zeta) d\mu_2(e_{\zeta})\} dx , \quad (2.125)$$

where  $g^K(x, \zeta) = \min [K, g(x, \zeta)]$  and  $h^K(x) = \min [K, h(x)]$ ,  $K$  denoting an arbitrary positive constant. We first examine the case where  $\mu_2(e_{\zeta})$  consists of a finite number of concentrated masses:  $\alpha_1 p$  in  $\zeta_1$ ,  $\alpha_2 p$  in  $\zeta_2$ ,  $\dots$ ,  $\alpha_m p$  in  $\zeta_m$ ,  $\sum_1^m \alpha_i = 1$ ,  $\alpha_i > 0$  ( $i = 1, 2, \dots, m$ ) and  $p < 1/16$ . An application of HÖLDER's inequality [15, p. 140] and the above result yields

$$\begin{aligned}
& \int_{R_N}^{R_n} \exp \{h^K(x) + p \sum_{i=1}^m \alpha_i g^K(x, \zeta_i)\} dx = \int_{R_N}^{R_n} \prod_{i=1}^m [\exp \{h^K(x) + p g^K(x, \zeta_i)\}]^{\alpha_i} dx \\
& \leq \prod_{i=1}^m \int_{R_N}^{R_n} \exp \{h^K(x) + p g^K(x, \zeta_i)\} dx^{\alpha_i} \leq \prod_{i=1}^m \int_{R_N}^{R_n} \exp \{h(x) + p g(x, \zeta_i)\} dx^{\alpha_i} \\
& \leq \frac{M}{\sin(\pi/8) \cos(\pi/32)} .
\end{aligned} \tag{2.126}$$

One proves without difficulty that it is always possible to approximate  $S_K$  arbitrarily close by substituting for  $\mu_2(e_\zeta)$  a finite number of concentrated masses of total weight  $p = \mu_2(\omega) < 1/16$ . Therefore, we infer from (2.126) that

$$S_K \leq \frac{M}{\sin(\pi/8) \cos(\pi/32)} . \tag{2.127}$$

Letting  $K \rightarrow \infty$  we obtain (2.93) as a consequence of (2.91), (2.125) and (2.127).

We now admit arbitrary rectifiable JORDAN curves  $\gamma_N$  and  $\gamma_n$ . Then there exists a conformal representation  $z = \varphi(\zeta)$  of some suitable circular ring  $R_N < |\zeta| < R_n$  ( $0 < R_N < R_n < +\infty$ ) onto the annular region  $\omega$ , bounded by  $\gamma_N$  and  $\gamma_n$ , such that the boundary components  $\gamma_N$  and  $\gamma_n$  correspond to  $|\zeta| = R_N$  and  $|\zeta| = R_n$ , respectively. Then the flux  $\Phi(u, \gamma; \Gamma)$  is invariant if  $u$  is transformed according to (2.8). We are going to prove that (2.84) is satisfied if  $\sigma_n$  is the image of the interval  $(R_N, R_n)$  on the  $\xi$ -axis ( $\zeta = \xi + i\eta$ ). Let  $\tilde{u}(\zeta)$  be defined by (2.8). Obviously, (2.84) is equivalent to

$$\int_{R_N}^{R_n} e^{\tilde{u}(\xi)} d\xi \leq \frac{M}{\sin(\pi/8) \cos(\pi/32)} . \tag{2.128}$$

All relevant quantities ( $e^u |dz|$ , mass, flux) are invariant under the transformation (2.8). For this reason the proof of (2.128) is essentially an application of the already treated special case to the function  $\tilde{u}$ . The only difficulty which arises stems from the boundary behavior of  $\varphi$ .

First we assume that  $\mu_2$  is concentrated in one point. Let  $h_1(z)$  and  $h_2(z)$  denote the best harmonic majorants in  $\omega$  of  $u_1(z)$  and  $u_2(z)$ , respectively. We define  $h(z) = h_1(z) - h_2(z)$ ,  $f(z) = \exp \{(h(z) + i h^*(z))/2\}$  ( $h^*$  being conjugate harmonic to  $h$ ) and introduce the transplanted functions  $\tilde{h}(\zeta) = h(\varphi(\zeta))$ ,  $\tilde{f}(\zeta) = f(\varphi(\zeta))$ . Assume for a moment that the inequalities

$$\limsup_{r \rightarrow R_N} \int_{-\pi}^{+\pi} |\tilde{f}^2(re^{it}) \varphi'(re^{it})| r dt \leq \int_{\gamma_N} e^{u(z)} |dz| \tag{2.129}$$

and

$$\limsup_{r \rightarrow R_n} \int_{-\pi}^{+\pi} |\tilde{f}^2(re^{it}) \varphi'(re^{it})| r dt \leq \int_{\gamma_n} e^{u(z)} |dz| \tag{2.130}$$

have been demonstrated. Then we would obtain (2.128) as an immediate consequence of (2.129), (2.130), Hypothesis (b) of Theorem 3 and (2.124) ( $u, f^2$  being replaced by  $\tilde{u}, \tilde{f}^2 \varphi'$ ). Since an application of HÖLDER's inequality would enable us again to get rid of the special hypothesis about  $\mu_2$ , the proof of Lemma 5 would thus be complete.

We are left to verify (2.129) and (2.130). Let  $\{h_{1k}(z)\}$  and  $\{h_{2l}(z)\}$  be defined as above, i. e. sequences of functions which are harmonic in  $\omega$ , continuous on the closure and which tend decreasingly to  $h_1(z)$  and  $h_2(z)$ , respectively. Let  $h_{kl} = h_{1k} - h_{2l}$ . We introduce the functions

$$f_{kl}(z) = \exp \{ (h_{kl}(z) + i h_{kl}^*(z)) / 2 \} \quad (k, l = 1, 2, 3, \dots), \quad (2.131)$$

where  $h_{kl}^*$  is conjugate harmonic to  $h_{kl}$ , and define

$$\tilde{f}_{kl}(\zeta) = f_{kl}(\varphi(\zeta)) \quad (k, l = 1, 2, 3, \dots). \quad (2.132)$$

We state that

$$\limsup_{r \rightarrow R_N} \int_{-\pi}^{+\pi} |\tilde{f}_{kl}^2(re^{it}) \varphi'(re^{it})| r dt \leq \int_{\gamma_N} |f_{kl}(z)|^2 |dz| \quad (2.133)$$

and

$$\limsup_{r \rightarrow R_n} \int_{-\pi}^{+\pi} |\tilde{f}_{kl}^2(re^{it}) \varphi'(re^{it})| r dt \leq \int_{\gamma_n} |f_{kl}(z)|^2 |dz|. \quad (2.134)$$

We briefly indicate a proof of these two inequalities<sup>5)</sup>. (For more details the reader is referred to the quoted articles.) Certain statements concerning the boundary behavior of  $\varphi$  will have to be verified. The analogous properties of the conformal mapping of simply connected domains are well known and we shall make extensive use of them.

Let us first verify that  $\varphi$  is continuous and of bounded variation on the boundary. We consider the outer boundary,  $|\zeta| = R_n$ . Let  $z = \varphi_1(w)$  denote a conformal mapping of a suitable circular disk  $|w| < R$  onto the interior of  $\gamma_n$ . We define  $w = \varphi_2(\zeta) = \varphi_1^{-1}(\varphi(\zeta))$ . This function represents the domain  $R_N < |\zeta| < R_n$  conformally onto an annulus  $\Omega$  with outer boundary  $|w| = R$ .  $\varphi$  has thus been decomposed into two steps,  $\varphi = \varphi_1(\varphi_2(\zeta))$ .  $\varphi_1$  is known to be continuous on  $|w| = R$  (for references see C. GATTEGNO and A. OSTROWSKI [14, p. 27]). Since  $\gamma_n$  is rectifiable,  $\varphi_1$  is also of bounded variation. Furthermore,  $w = \varphi_2(\zeta)$  is analytic on  $|\zeta| = R_n$  because this part of the boundary is mapped onto a circle. Consequently,  $z = \varphi(\zeta)$  is continuous and of bounded variation on  $|\zeta| = R_n$ . The same is true for the inner circle  $|\zeta| = R_N$ .

Next we prove that  $\varphi$  is absolutely continuous on the boundary. From the LAURENT expansion we conclude that  $\varphi$  admits the representation  $\varphi(\zeta) = \varphi_I(\zeta) + \varphi_{II}(\zeta)$ ,  $\varphi_I$  and  $\varphi_{II}$  being analytic in  $|\zeta| < R_n$  and in  $|\zeta| > R_N$ , respec-

<sup>5)</sup> Professor M. RIESZ kindly suggested to us the following demonstration.

tively. Since  $\varphi$  is continuous and of bounded variation on  $|\zeta| = R_n$ , the same is true for  $\varphi_{II}$ . Hence, by a well known theorem of F. and M. RIESZ [28],  $\varphi_{II}$  is absolutely continuous on  $|\zeta| = R_n$ . Consequently,  $\varphi$  is absolutely continuous on  $|\zeta| = R_n$ . An analogous reasoning proves the same for  $|\zeta| = R_N$ .

$\varphi$  can be written as the POISSON integral of its boundary values

$$\varphi(\zeta) = \int K(\zeta, Z) \varphi(Z) |dZ| ,$$

where  $K$  denotes the POISSON kernel,  $\zeta$  an interior and  $Z$  a boundary point of  $R_N < |\zeta| < R_n$ . Because of the circular symmetry of the domain the kernel  $K$  depends only on  $|\zeta|$ ,  $|Z|$  and  $\arg(\zeta - Z)$ . Consequently, if we put  $\zeta = \rho e^{i\theta}$  and  $Z = R_N e^{it}$  (or  $Z = R_n e^{it}$ ), we have  $\partial K / \partial \theta = -\partial K / \partial t$ . This relation allows us to convert the differentiation of  $\int K(\zeta, Z) \varphi(Z) |dZ|$  with respect to  $\zeta$  into a differentiation along the boundary (see M. RIESZ [29, p. 55]). A partial integration then permits us to conclude that  $\varphi'(\zeta)$  is given by the POISSON integral of its (almost everywhere existing) values

$$\varphi'(R_N e^{it}) = \frac{1}{i R_N e^{it}} \frac{\partial \varphi(R_N e^{it})}{\partial t} \quad \text{and} \quad \varphi'(R_n e^{it}) = \frac{1}{i R_n e^{it}} \frac{\partial \varphi(R_n e^{it})}{\partial t}$$

on the bounding circles. It follows that

$$|\varphi'(\zeta)| \leq A_1(\zeta) + A_2(\zeta) , \quad (2.135)$$

where

$$A_1(\zeta) = \int_{-\pi}^{+\pi} K(\zeta, R_N e^{it}) |\varphi'(R_N e^{it})| R_N dt$$

and

$$A_2(\zeta) = \int_{-\pi}^{+\pi} K(\zeta, R_n e^{it}) |\varphi'(R_n e^{it})| R_n dt .$$

For  $|Z| = R_N$ ,  $|\zeta| \rightarrow R_n$  implies  $K(\zeta, Z) \rightarrow 0$ , the convergence being uniform in  $Z$  and  $\arg \zeta$ . Consequently

$$\lim_{r \rightarrow R_n} \int_{-\pi}^{+\pi} A_1(r e^{it}) r dt = 0 . \quad (2.136)$$

Let  $K_0$  denote the POISSON kernel for the domain  $|\zeta| < R_n$ . For  $|Z| = R_n$ ,  $K(\zeta, Z) \leq K_0(\zeta, Z)$ . Hence

$$\begin{aligned} \int_{-\pi}^{+\pi} A_2(r e^{it}) r dt &\leq \int_{|\zeta|=r} \int_{|Z|=R_n} K_0(\zeta, Z) |\varphi'(Z)| |d\zeta| |dZ| \\ &\leq \int_{-\pi}^{+\pi} |\varphi'(R_n e^{it})| R_n dt = \int_{\gamma_n} |dz| . \end{aligned} \quad (2.137)$$

From (2.135), (2.136) and (2.137) we infer

$$\limsup_{r \rightarrow R_n} \int_{-\pi}^{+\pi} |\varphi'(r e^{it})| r dt \leq \int_{-\pi}^{+\pi} |\varphi'(R_n e^{it})| R_n dt = \int_{\gamma_n} |dz| . \quad (2.138)$$

We state that, more generally

$$\limsup_{r \rightarrow R_n} \int_{t_1}^{t_2} |\varphi'(re^{it})| r dt \leq \int_{t_1}^{t_2} |\varphi'(R_n e^{it})| R_n dt = \int_{\gamma'_n} |dz| \quad (2.139)$$

holds for any pair of values  $t_1$  and  $t_2$  ( $-\pi \leq t_1 < t_2 \leq +\pi$ ),  $\gamma'_n$  denoting the portion of  $\gamma_n$  which corresponds to the arc  $(R_n e^{it_1}, R_n e^{it_2})$ . Indeed, suppose that (2.139) is not fulfilled for some such arc. Then there exists a sequence of radii  $\{r_k\} \rightarrow R_n$  such that

$$\lim_{k \rightarrow \infty} \int_{t_1}^{t_2} |\varphi'(r_k e^{it})| r_k dt > \int_{t_1}^{t_2} |\varphi'(R_n e^{it})| R_n dt$$

and

$$\lim_{k \rightarrow \infty} \int_{-\pi}^{+\pi} |\varphi'(r_k e^{it})| r_k dt \leq \int_{-\pi}^{+\pi} |\varphi'(R_n e^{it})| R_n dt .$$

These two inequalities imply

$$\lim_{k \rightarrow \infty} \int_{t_2}^{t_1+2\pi} |\varphi'(r_k e^{it})| r_k dt < \int_{t_2}^{t_1+2\pi} |\varphi'(R_n e^{it})| R_n dt .$$

But since  $|\varphi'(r_k e^{it})| \rightarrow |\varphi'(R_n e^{it})|$  for almost all  $t$ , we have thus obtained a contradiction to FATOU's lemma [30, p. 29]. This proves (2.139). (2.134) is an immediate consequence of (2.139). (2.133) can be proved by the same method.

Now, let  $k \rightarrow \infty$ . We conclude from (2.133) and (2.134) that

$$\begin{aligned} & \limsup_{r \rightarrow R_N} \int_{-\pi}^{+\pi} \exp \{h_1(\varphi(re^{it})) - h_{2l}(\varphi(re^{it}))\} |\varphi'(re^{it})| r dt \\ & \leq \int_{\gamma_N} \exp \{u_1(z) - h_{2l}(z)\} |dz| \end{aligned} \quad (2.140)$$

and

$$\begin{aligned} & \limsup_{r \rightarrow R_n} \int_{-\pi}^{+\pi} \exp \{h_1(\varphi(re^{it})) - h_{2l}(\varphi(re^{it}))\} |\varphi'(re^{it})| r dt \\ & \leq \int_{\gamma_n} \exp \{u_1(z) - h_{2l}(z)\} |dz| \quad (l = 1, 2, 3, \dots) . \end{aligned} \quad (2.141)$$

Indeed, the symbol „ $>$ ” in either of these inequalities leads to a contradiction: An easy argument then yields the conclusion that, for any fixed  $l$  and sufficiently large  $k(l)$ , the integral on the left-hand sides of (2.133) and (2.134) is *not* a convex function of  $\log r$ . But, on the other hand, it should possess this convexity property, since it represents the mean value of a subharmonic function.

Finally we let  $l \rightarrow \infty$ . By making again use of the convexity we obtain (2.129) and (2.130) as consequences of (2.140) and (2.141), respectively. This completes the proof of Lemma 5.

Theorem 1, 2 and 3 are concerned with the behavior of  $u$  near  $\Gamma$ . Corresponding results can be obtained for the neighborhood of  $\Delta$ . We state them without giving detailed proofs. Let  $\Phi(\Delta)$  be defined analogously to  $\Phi(\Gamma)$ . (Interchange  $\Gamma$  with  $\Delta$  and assume that  $n$  designates the inner normal in all definitions.) Suppose that  $\Phi(\Delta)$  exists.

**Theorem 4.** *If  $\Phi(\Delta) < +1$ , then there exists a locally rectifiable path  $\sigma$  in  $\Omega$ , tending to  $\Delta$ , such that (2.7) is fulfilled.*

**Theorem 5.** *If  $\Delta$  contains more than one point, and if  $\Phi(\Delta) < +\infty$ , then there exists a locally rectifiable path  $\sigma$ , tending to  $\Delta$ , such that (2.7) is satisfied.*

**Theorem 6.** *Suppose there exists a sequence  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , of rectifiable JORDAN curves, enclosing  $\Delta$ , in  $\Omega$  and a number  $M$  such that*

- (a)  $\{\gamma_n\}$  comes arbitrarily near to  $\Delta$ ,
- (b)  $\int_{\gamma_n} e^u |dz| < M$  for all  $n$ .

*Then, if  $\Phi(\Delta) \neq +1$ , there exists a locally rectifiable path  $\sigma$ , tending to  $\Delta$ , such that (2.7) is fulfilled.*

These results follow from the previous ones by an inversion. We mention that Theorems 4 and 6 can also be obtained as corollaries of Theorems 8 and 9 (section 4), respectively.

We conclude this section with a remark concerning a special case. Let  $w = f(z)$  be a (not necessarily single-valued) complex analytic function which is defined and  $\neq 0$  throughout  $\Omega$  except possibly at a (finite or infinite) number of isolated points  $a_1, a_2, a_3, \dots$ . Suppose that in the neighborhood of these  $f$  admits the representation

$$f(z) = (z - a_k)^{p_k} g(z) \quad (k = 1, 2, 3, \dots),$$

where  $g$  is regular and  $\neq 0$  at  $a_k$ , and  $p_k$  denotes an arbitrary real number. (In particular,  $f$  may be of the form  $[F(z)]^\lambda$ , where  $F$  denotes a meromorphic function and  $\lambda$  is a real number.) Then the (single-valued) function

$$u(z) = \log |f(z)|$$

is harmonic throughout  $\Omega$  except at  $a_1, a_2, a_3, \dots$ , where it possesses isolated logarithmic singularities. Hence  $u$  admits the representation (2.1). Let  $\gamma$  denote a JORDAN curve in  $\Omega$ , enclosing  $\Delta$ , which does not pass through any of the  $a_k$ 's. Then

$$\Phi(u, \gamma; \Gamma) = \frac{1}{2\pi} \int_{\gamma} d[\arg f(z)],$$

$\gamma$  being described in the positive sense. Since  $e^{u(z)} |dz| = |f(z)| |dz|$ , Theo-

lems 1 to 6 thus imply certain results on integrals of moduli of analytic functions in the case where something is known about the variation of the argument along closed curves.

If the region of definition of  $f$  can be extended to  $\Omega_0$ , then

$$\Phi(u, \gamma; \Gamma) = \sum_{a_k \in \omega} p_k ,$$

where  $\omega$  designates the interior region of  $\gamma$ .

### 3. A characteristic property of polynomials

**Theorem 7.** *An entire analytic function  $w = f(z)$  is a polynomial if and only if there exists a positive number  $\lambda$  such that*

$$\int_{\sigma} |f(z)|^{-\lambda} |dz| = +\infty \quad (3.1)$$

for every locally rectifiable path  $\sigma$  tending to infinity.

**Remarks.** It is natural to ask whether this theorem remains valid if the class of admissible curves  $\sigma$  is more restricted. We have no results in this direction. For example, the following question is still open: *Let  $w = f(z)$  be an entire analytic function. Suppose there exists a positive number  $\lambda$  such that*

$$\int_1^{+\infty} |f(\varrho e^{i\Theta})|^{-\lambda} d\varrho = +\infty$$

for all  $\Theta$  ( $0 \leq \Theta < 2\pi$ ). Does this imply that  $f$  is a polynomial?

One might also consider other regions of definition of  $f(z)$  instead of the entire plane. In this respect Theorem 2 immediately yields the following statement: *Suppose the function  $w = f(z) \not\equiv 0$  is defined and analytic in a simply-connected, proper subregion  $\Omega_0$  of the  $z$ -plane. Then, given any  $\lambda > 0$ , there exists a locally rectifiable path  $\sigma$ , tending to the boundary  $\Gamma$  of  $\Omega_0$ , such that*

$$\int_{\sigma} |f(z)|^{-\lambda} |dz| < +\infty .$$

**Proof of Theorem 7.** Suppose the function  $w = f(z)$  satisfies condition (3.1) for some  $\lambda > 0$ . Let  $N$  denote the number of zeros of  $f$  ( $N \leq +\infty$ ). We exclude the trivial function  $f \equiv 0$  and define  $u(z) = u_1(z) - u_2(z)$ , where  $u_1(z) \equiv 0$  and  $u_2(z) = \lambda \log |f(z)|$ . Let  $\Omega_0$  denote the entire  $z$ -plane. Using the notations of section 2 we then have  $\Phi(\Gamma) = -\lambda N$ . Since  $u$  does not satisfy the statement of Theorem 1,  $\Phi(\Gamma) \geq -1$ . Hence  $N \leq 1/\lambda < +\infty$ . So  $f$  admits the representation

$$f(z) = e^{\Psi(z)} \prod_{k=1}^N (z - a_k) , \quad (3.2)$$

where  $\Psi$  is an entire function and  $a_1, a_2, \dots, a_N$  are the zeros of  $f$ . The function

$$\zeta = \varphi(z) = \int_0^z e^{-\lambda \psi(z)} dz \quad (3.3)$$

yields a conformal mapping of the finite  $z$ -plane onto a RIEMANN surface  $R$  without branch points, extending over the  $\zeta$ -plane. We have to distinguish between two cases

(a)  $R$  coincides with the entire finite plane. Then  $\zeta = \varphi(z)$  is necessarily an entire linear function. So  $\psi(z)$  is a constant and, therefore,  $f(z)$  is a polynomial.

(b) There exists a finite point  $\zeta_0$  with the following property: On some sheet of  $R$  the half-open segment  $\zeta(t) = \zeta_0 t$  ( $0 \leq t < 1$ ) belongs to  $R$ , whereas  $\zeta_0$  lies on the boundary. This segment is the image of an analytic curve  $\sigma_0$  in the  $z$ -plane which tends to infinity. By (3.3) we have

$$\int_{\sigma_0} |e^{\psi(z)}|^{-\lambda} |dz| = \int_{\sigma_0} |\varphi'(z)| |dz| = |\zeta_0| < +\infty. \quad (3.4)$$

From (3.2) and (3.4) we conclude that

$$\int_{\sigma_0} |f(z)|^{-\lambda} |dz| < +\infty, \quad (3.5)$$

contrary to hypothesis. (It is understood that  $\sigma_0$  has to be slightly modified if zeros of  $f$  should lie on it.)

So  $f$  must be a polynomial. The converse is obvious.

#### 4. On complete conformal metrics defined on finitely connected, open RIEMANN surfaces

We first give a conformally invariant formulation of Theorems 1 and 3. Let  $\Omega$  be a doubly connected, open RIEMANN surface on which a conformal metric (1.7) is defined. We assume that  $u$  admits the local representation (1.9). Let  $\Gamma$  and  $\Delta$  denote the ideal boundary components of  $\Omega$  and let  $\gamma$  be a JORDAN curve<sup>6)</sup> in  $\Omega$  which is not nullhomotopic.

Assume for a moment that  $u$  is of class  $C^2$  and that  $\gamma$  is analytic. Then we define

$$I(e^u |dz|, \gamma; \Gamma) = \int_{\gamma} \left( k_e + \frac{\partial u}{\partial n} \right) |dz|, \quad (4.1)$$

---

<sup>6)</sup> The definitions of the concepts JORDAN curve, homotopic, locally rectifiable, analytic are to be understood with respect to the underlying RIEMANN surface.

where  $k_e$  and  $n$  are to be determined as in (1.5), the orientation of  $\gamma$  being chosen such that  $\Gamma$  lies on the right. The integral (4.1) (in fact, even the differential  $\left(k_e + \frac{\partial u}{\partial n}\right)|dz|$ ) is a conformal invariant. This is immediately clear from the geometrical interpretation (1.6), but it can also be verified by direct calculation.

In order to define  $I(e^u|dz|, \gamma; \Gamma)$  for general  $u$  and  $\gamma$  we consider a sequence  $\{\delta_k\}$ ,  $k = 1, 2, 3, \dots$ , of JORDAN curves in  $\Omega$ , chosen such that the annuli  $(\Delta, \delta_k)$  tend increasingly to  $(\Delta, \gamma)$  as  $k \rightarrow \infty$ . In  $(\delta_k, \gamma)$  we introduce a conformal metric  $e^{h_k(z)}|dz|$ , defining

$$h_k(z) = u(z) + \int_{(\delta_k, \gamma)} g_k(z, \zeta) d\mu(e_\zeta) . \quad (4.2)$$

Here  $g_k$  denotes GREEN's function for  $(\delta_k, \gamma)$  and  $\mu = \mu_1 - \mu_2$ , where  $\mu_1$  and  $\mu_2$  are the measures associated with  $u_1$  and  $u_2$ , respectively. (It should be noticed that  $\mu$  does not depend on the choice of decomposition or uniformizer. Hence the integral on the right-hand side of (4.2) is a scalar. Consequently,  $e^{h_k(z)}|dz|$  is indeed a conformal invariant.) Let  $\delta'_k$  be an arbitrary analytic JORDAN curve in  $(\delta_k, \gamma)$  which is not nullhomotopic. Then  $I(e^{h_k}|dz|, \delta'_k; \gamma)$  is well defined. If  $\Omega$  is a schlicht region in the finite  $z$ -plane, then  $h_k(z)$ , defined by (4.2), is identical with the function designated in the same way in (2.4). This is implied by the decomposition theorem of F. RIESZ [27, II, p. 357]). If, furthermore,  $\Gamma$  denotes the outer boundary of  $\Omega$ , then, obviously

$$I(e^{h_k}|dz|, \delta'_k; \gamma) = 2\pi[\Phi(h_k, \delta'_k; \gamma) + 1] . \quad (4.3)$$

We observe that  $I(e^{h_k}|dz|, \delta'_k; \gamma)$  does not depend on the choice of  $\delta'_k$ , this being true for the right-hand side of (4.3). We further conclude from (4.3) and the existence of the limit (2.4) that

$$I(e^u|dz|, \gamma; \Gamma) = \lim_{k \rightarrow \infty} I(e^{h_k}|dz|, \delta'_k; \gamma) \quad (4.4)$$

always exists, being finite and independent of the choice of  $\{\delta_k\}$ . Clearly

$$I(e^u|dz|, \gamma; \Gamma) = 2\pi[\Phi(u, \gamma; \Gamma) + 1] . \quad (4.5)$$

These relations hold under the above mentioned special assumptions. But  $\Omega$  can always be mapped conformally onto a schlicht annulus such that  $\Gamma$  corresponds to the outer boundary. Furthermore, since  $I(e^{h_k}|dz|, \delta'_k; \gamma)$  is a conformal invariant, the existence of  $I(e^u|dz|, \gamma; \Gamma)$  is thus assured in any case.

Now, let  $\{\gamma_l\}$ ,  $l = 1, 2, 3, \dots$ , be an arbitrary sequence of JORDAN curves which are not nullhomotopic and such that the regions  $(\Delta, \gamma_l)$  tend increas-

ingly to  $\Omega$  as  $l \rightarrow \infty$ . Assume that the limit

$$I(\Gamma) = \lim_{l \rightarrow \infty} I(e^u |dz|, \gamma_l; \Gamma) \quad (4.6)$$

exists for any such sequence, admitting the values  $+\infty$  and  $-\infty$ . Of course,  $I(\Gamma)$  is necessarily independent of the choice of  $\{\gamma_l\}$ .

From (4.5) we infer that

$$I(\Gamma) = 2\pi[\Phi(\Gamma) + 1] \quad (4.7)$$

in the case where  $\Omega$  is a schlicht region and  $\Gamma$  denotes its outer boundary. Since  $I(\Gamma)$  is a conformal invariant, this relation gives rise immediately to the following extensions of Theorems 1 and 3<sup>7)</sup>:

**Theorem 8.** *If  $I(\Gamma) < 0$ , then there exists a locally rectifiable path  $\sigma$ , tending to  $\Gamma$ , such that  $\int_{\sigma} e^{u(z)} |dz| < +\infty$ .*

**Theorem 9.** *Suppose there exists a sequence  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , of locally rectifiable JORDAN curves which are not nullhomotopic and a number  $M$  such that*

(a)  $\{\gamma_n\}$  comes arbitrarily near to  $\Gamma$ ,

(b)  $\int_{\gamma_n} e^{u(z)} |dz| < M$  for all  $n$ .

*Then, if  $I(\Gamma) \neq 0$ , there exists a locally rectifiable path  $\sigma$ , tending to  $\Gamma$ , such that  $\int_{\sigma} e^{u(z)} |dz| < +\infty$ .*

After this preparation we take up the concepts developed in the introduction. Consider an open RIEMANN surface  $S$  on which a conformal metric (1.7) is defined. Assume that  $u$  admits the local representation (1.9). We define: The metric  $e^{u(z)} |dz|$  is said to be *complete* if  $\int_{\sigma} e^{u(z)} |dz| = +\infty$  for every locally rectifiable path  $\sigma$  which tends to the ideal boundary of  $S$ .

**Theorem 10.** *Let  $S$  be a finitely connected, open RIEMANN surface on which a complete conformal metric  $e^{u(z)} |dz|$  is defined. Suppose that the curvatura integra  $C$  exists. Then  $C \leq 2\pi\chi$ , where  $\chi$  denotes the EULER-POINCARÉ characteristic of  $S$ .*

**Remark.** This is a result of S. COHN-VOSSEN (Satz 6, p. 79 in [10]) in extended form. (For further comments see introduction.)

---

<sup>7)</sup> The meaning of " $\sigma$  tends to  $\Gamma$ " and of " $\{\gamma_n\}$  comes arbitrarily near to  $\Gamma$ " has been defined in section 2 for schlicht annular regions. It is clear how these definitions have to be reformulated for arbitrary RIEMANN surfaces.

**Proof.**  $S$  is homeomorphic to a closed surface from which a finite number (say  $N$ ) of points have been removed<sup>8)</sup>. There exists a subcompact  $K$  of  $S$ , bounded by  $N$  JORDAN curves,  $\Delta_1, \Delta_2, \dots, \Delta_N$ , such that the open set  $S - K$  consists of  $N$  doubly connected components,  $\Omega_1, \Omega_2, \dots, \Omega_N$ . Thereby each  $\Omega_r$  is bounded by  $\Delta_r$  and a second (ideal) boundary component  $\Gamma_r$ . Let  $\gamma_r$  be an arbitrary JORDAN curve in  $\Omega_r$  which is not nullhomotopic with respect to  $\Omega_r$  ( $r = 1, 2, \dots, N$ ). Let  $\Sigma$  denote the subregion of  $S$  whose boundary consists of  $\gamma_1, \dots, \gamma_{N-1}$  and  $\gamma_N$ . We are going to prove the GAUSS-BONNET formula<sup>9)</sup>

$$2\pi\mu(\Sigma) + 2\pi\chi = \sum_{r=1}^N I(e^u|dz|, \gamma_r; \Gamma_r) . \quad (4.8)$$

Assume first that  $u$  is of class  $C_2$  and that  $\gamma_1, \dots, \gamma_{N-1}$  and  $\gamma_N$  are analytic. Consider a triangulation of the closure of  $\Sigma$  consisting of analytic arcs. Let  $T_j$  denote the interiors,  $B_j$  the boundaries, and  $\alpha_{jl}$  ( $l = 1, 2, 3$ ) the exterior angles of the triangles ( $j = 1, 2, \dots, M$ ). We may suppose that one and the same local uniformizer can be used in a neighborhood of  $T_j \cup B_j$ . By GAUSS's theorem and the definition of  $k_e$

$$\iint_{T_j} \Delta u dx_j dy_j + 2\pi - \sum_{l=1}^3 \alpha_{jl} = \int_{B_j} \left( k_e + \frac{\partial u}{\partial n} \right) |dz_j| \quad (j = 1, 2, \dots, M) ,$$

if we integrate along  $B_j$  in the positive sense. We add all these relations. Because of the conformal invariance of  $\left( k_e + \frac{\partial u}{\partial n} \right) |dz|$ , and the coherence of the orientation most integrals on the right-hand side drop out. Furthermore, the EULER-POINCARÉ characteristic  $\chi$  of  $S$  appears in a well known way. We obtain

$$\iint_{\Sigma} \Delta u dx dy + 2\pi\chi = \sum_{r=1}^N \int_{\gamma_r} \left( k_e + \frac{\partial u}{\partial n} \right) |dz| ,$$

i. e. relation (4.8) for the considered special case.

We now proceed to the general case but still assume that the  $\gamma_r$ 's are analytic and free of mass. Without losing generality we then may suppose that the entire „skeleton” of the triangulation,  $L = \bigcup_{j=1}^M B_j$ , is free of mass. Let  $\{D_k\}$ ,  $k = 1, 2, 3, \dots$ , be a sequence of regions, tending decreasingly to  $L$  as  $k \rightarrow \infty$ , each of which is bounded by  $M + N$  JORDAN curves lying, respectively, in  $T_1, T_2, \dots, T_M, (\gamma_1, \Gamma_1), (\gamma_2, \Gamma_2), \dots, (\gamma_N, \Gamma_N)$ . Consider the conformal metric

<sup>8)</sup> cf. B. v. KERÉKJÁRTÓ [19, chapter 5].

<sup>9)</sup> For the functiontheoretical aspects of the GAUSS-BONNET formula see also R. NEVANLINNA [21].

$e^{H_k(z)}|dz|$  in  $D_k$ , defining

$$H_k(z) = u(z) + \int_{D_k} G_k(z, \zeta) d\mu(e_\zeta) , \quad (4.9)$$

where  $G_k$  designates GREEN's function for  $D_k$  ( $k = 1, 2, 3, \dots$ ). Then

$$\int_{B_j} \frac{\partial H_k}{\partial n} |dz_j| + 2\pi - \sum_{l=1}^3 \alpha_{jl} = \int_{B_j} \left( k_e + \frac{\partial H_k}{\partial n} \right) |dz_j| \quad (j = 1, 2, \dots, M) ,$$

and, by addition

$$\sum_{j=1}^M \int_{B_j} \frac{\partial H_k}{\partial n} |dz_j| + 2\pi\chi = \sum_{r=1}^N \int_{\gamma_r} \left( k_e + \frac{\partial H_k}{\partial n} \right) |dz| . \quad (4.10)$$

We state that

$$\lim_{k \rightarrow \infty} \int_{B_j} \frac{\partial H_k}{\partial n} |dz_j| = 2\pi\mu(T_j) \quad (j = 1, 2, \dots, M) \quad (4.11)$$

and

$$\lim_{k \rightarrow \infty} \int_{\gamma_r} \left( k_e + \frac{\partial H_k}{\partial n} \right) |dz| = I(e^u |dz|, \gamma_r; \Gamma_r) \quad (r = 1, 2, \dots, N) . \quad (4.12)$$

In order to verify (4.11) we introduce (for a fixed  $j$ ) a sequence of doubly connected regions  $\{E_k\}$  which are bounded by analytic JORDAN curves and tend decreasingly to  $B_j$ . Consider the conformal metric  $e^{h_k(z)}|dz|$  in  $E_k$ , defining

$$h_k(z) = u(z) + \int_{E_k} g_k(z, \zeta) d\mu(e_\zeta) , \quad (4.13)$$

where  $g_k$  denotes GREEN's function for  $E_k$  ( $k = 1, 2, 3, \dots$ ). From the results of F. RIESZ [27] one concludes without difficulty that

$$\lim_{k \rightarrow \infty} \int_{B_j} \frac{\partial h_k}{\partial n} |dz_j| = 2\pi\mu(T_j) . \quad (4.14)$$

(4.9) and (4.13) imply

$$\int_{B_j} \left( \frac{\partial H_k}{\partial n} - \frac{\partial h_k}{\partial n} \right) |dz_j| = \int_{D_k} \left[ \int_{B_j} \frac{dG_k(z, \zeta)}{\partial n_z} |dz| \right] d\mu(e_\zeta) - \int_{E_k} \left[ \int_{B_j} \frac{\partial g_k(z, \zeta)}{\partial n_z} |dz| \right] d\mu(e_\zeta) . \quad (4.15)$$

Obviously we have

$$\left| \frac{1}{2\pi} \int_{B_j} \frac{\partial G_k(z, \zeta)}{\partial n_z} |dz| \right| \leq 1 \quad (4.16)$$

and

$$\left| \frac{1}{2\pi} \int_{B_j} \frac{\partial g_k(z, \zeta)}{\partial n_z} |dz| \right| \leq 1 , \quad (4.17)$$

if we interpret these integrals as fluxes. From (4.15), (4.16) and (4.17) we infer that

$$\left| \int_{B_j} \left( \frac{\partial H_k}{\partial n} - \frac{\partial h_k}{\partial n} \right) |dz_j| \right| \leq 2\pi [\mu(E_k) + \mu(D_k)] . \quad (4.18)$$

The right-hand side of this inequality tends to 0 as  $k \rightarrow \infty$ , because  $\mu(L) = \mu(B_j) = 0$ . Consequently, (4.11) follows from (4.14) and (4.18). (4.12) can be demonstrated in a similar way.

Since  $\sum_{j=1}^M \mu(T_j) = \mu(\Sigma)$ , (4.11) and (4.12) allow us to conclude that (4.8) is the limit of (4.10) as  $k \rightarrow \infty$ .

In order to get rid of the hypotheses that the  $\gamma_r$ 's are analytic and free of mass we exhaust an arbitrary  $\Sigma$  by an increasing sequence of regions whose bounding curves satisfy these conditions. Relation (4.8), formulated for  $\Sigma$ , is immediately obtained as the limit of the corresponding equalities already verified for the subregions.

From (4.8) one can conclude that the limits  $I(\Gamma_r)$ ,  $r = 1, 2, \dots, N$ , exist. (This is a consequence of the existence of  $C = -2\pi\mu(S)$  and proved by letting an arbitrary one of the  $\gamma_r$ 's move to the boundary while all others are kept fixed.) Now, if  $\Sigma$  tends increasingly to  $S$ , then (4.8) yields in the limit

$$C = 2\pi\chi - \sum_{r=1}^N I(\Gamma_r) . \quad (4.19)$$

From Theorem 8 and the completeness of the metric we infer that  $I(\Gamma_r) \geq 0$  for  $r = 1, 2, \dots, N$ . Hence, by (4.19),  $C \leq 2\pi\chi$ . Q. E. D.

**Theorem 11.** *Let  $S$  be a finitely connected, open RIEMANN surface on which a complete conformal metric  $e^{u(z)}|dz|$  is defined. Suppose there exists a sequence  $\{\gamma_n\}$ ,  $n = 1, 2, 3, \dots$ , of locally rectifiable JORDAN curves with the following properties:*

- (1) *they are not nullhomotopic,*
- (2) *their lengths  $\int_{\gamma_n} e^{u(z)}|dz|$  are uniformly bounded,*
- (3)  *$\{\gamma_n\}$  comes arbitrarily near to every boundary component of  $S$  <sup>10)</sup>.*

*Assume further that the curvatura integra  $C$  exists. Then  $C = 2\pi\chi$ , where  $\chi$  denotes the EULER-POINCARÉ characteristic of  $S$ .*

**Remarks.** This result implies a theorem of S. COHN-VOSSEN (Satz 7, p. 79 in [10]) which states that  $C = 2\pi\chi$  for every finitely connected, open, two-

<sup>10)</sup> i. e. the  $\gamma_n$ 's penetrate into each  $\Omega_r$ ,  $r = 1, 2, \dots, N$ , for any choice of  $K$ .

dimensional RIEMANNIAN manifold whose curvatura integra exists and which does not possess a so-called "eigentlicher Kelch". By going back to the definition of this concept and using the notations introduced for the proof of Theorem 10 one arrives to the following formulation of COHN-VOSSEN's hypothesis: Let

$$m(\Omega_r) = \inf_{\gamma} \left[ \int_{\gamma} e^{u(z)} |dz| \right],$$

admitting to competition all locally rectifiable JORDAN curves in  $\Omega_r$  which are not nullhomotopic. A sequence  $\{\gamma_n\}$  of such curves is called a *minimal sequence* of  $\Omega_r$  if

$$\lim_{n \rightarrow \infty} \int_{\gamma_n} e^{u(z)} |dz| = m(\Omega_r).$$

COHN-VOSSEN postulated that, given an arbitrary subcompact  $K_0$  of  $S$ , there should always exist a connected subcompact  $K$ , containing  $K_0$ , such that all components  $\Omega_r$ ,  $r = 1, 2, \dots, N$ , of  $S - K$  have the following property: Each minimal sequence  $\{\gamma_n\}$  of  $\Omega_r$  comes arbitrarily near to  $\Gamma_r$ .

It is clear that COHN-VOSSEN's hypothesis is stronger than ours. The ordinary circular cylinder imbedded in 3-space is a trivial example of a manifold to which our result applies while COHN-VOSSEN's theorem does not.

The following statement is also a corollary of Theorem 11: *Let  $M$  be a finitely connected, complete, open, two-dimensional RIEMANNIAN manifold whose curvatura integra  $C$  exists. Suppose there exists a sequence of subcompacts, tending increasingly to  $M$ , the boundaries of which are of uniformly bounded length. Then  $C = 2\pi\chi$ .* Professor H. HOPF points out to us that if we make the additional assumptions that  $M$  is analytic and has everywhere non-negative curvature (and is therefore necessarily simply connected), then this result follows from two theorems of F. FIALA (Theorems A and D, pp. 299–300 in [12]) and the previously mentioned result of COHN-VOSSEN (Satz 6, p. 79 in [10]).

**Proof.** If infinitely many of the  $\gamma_n$ 's would intersect  $K$ , then a reasoning quite similar to the one used in the proof of Lemma 2 would yield a contradiction to the hypothesis of completeness.

Hence only a finite number of  $\gamma_n$ 's intersect  $K$ . From this we conclude that each  $\Omega_r$  contains a subsequence  $\{\gamma_n^{(r)}\}$  of  $\{\gamma_n\}$  which comes arbitrarily near to  $\Gamma_r$ . Therefore, by Theorem 9 and the completeness of the metric,  $I(\Gamma_r) = 0$  for  $r = 1, 2, \dots, N$ . Consequently, by (4.19),  $C = 2\pi\chi$ . Q. E. D.

**Theorem 12.** *Let  $S$  be a finitely connected, open RIEMANN surface on which a complete conformal metric  $e^{u(z)} |dz|$  of finite total area  $A = \iint_S e^{2u} dx dy$  is defined. Suppose that the curvatura integra  $C$  exists. Then  $C = 2\pi\chi$ , where  $\chi$  denotes the EULER-POINCARÉ characteristic of  $S$ .*

**Proof.** Again we make use of the concepts introduced for the proof of Theorem 10. Considering an arbitrary one of the regions  $\Omega_r$ , we shall prove that  $I(\Gamma_r) = 0$ . This, combined with relation (4.9), will demonstrate the theorem.

The region  $\Omega_r$  is conformally equivalent to a schlicht circular ring in the  $z$ -plane,  $R_1 < |z| < R_2$  ( $0 < R_1 < R_2 \leq +\infty$ ). We distinguish between two cases depending on whether  $R_2$  is finite or infinite. Let us begin by showing that the first possibility cannot occur.

(I)  $R_2 < +\infty$ . Let  $R$  ( $R_1 < R < R_2$ ) be chosen arbitrarily. By SCHWARZ'S inequality

$$\begin{aligned} \int_{R_1}^R \int_0^{2\pi} e^{2u(\varrho e^{i\varphi})} \varrho d\varrho d\varphi &\geq R_1 \int_0^{2\pi} \left[ \int_{R_1}^R e^{2u(\varrho e^{i\varphi})} d\varrho \right] d\varphi \geq \frac{R_1}{R - R_1} \int_0^{2\pi} \left[ \int_{R_1}^R e^{u(\varrho e^{i\varphi})} d\varrho \right]^2 d\varphi \\ &\geq \frac{2\pi R_1}{R - R_1} \left[ \inf_{0 \leq \varphi < 2\pi} \int_{R_1}^R e^{u(\varrho e^{i\varphi})} d\varrho \right]^2. \end{aligned} \quad (4.20)$$

Furthermore

$$\lim_{R \rightarrow R_2} \left[ \inf_{0 \leq \varphi < 2\pi} \int_{R_1}^R e^{u(\varrho e^{i\varphi})} d\varrho \right] = +\infty, \quad (4.21)$$

since otherwise an application of Lemma 2 would yield a contradiction to the completeness of the metric. It follows from (4.20) and (4.21) that

$$\iint_{\Omega_r} e^{2u} dx dy = +\infty,$$

contrary to hypothesis. Hence  $R_2 = +\infty$ .

(II)  $R_2 = +\infty$ . Since

$$\iint_{\Omega_r} e^{2u(\varrho e^{i\varphi})} \varrho d\varrho d\varphi < +\infty,$$

there must exist a sequence of radii  $\{\varrho_n\} \rightarrow \infty$  for which

$$\int_0^{2\pi} e^{2u(\varrho_n e^{i\varphi})} \varrho_n d\varphi < \frac{1}{\varrho_n}.$$

Hence, by SCHWARZ'S inequality

$$\left[ \int_0^{2\pi} e^{u(\varrho_n e^{i\varphi})} \varrho_n d\varphi \right]^2 \leq 2\pi \varrho_n \int_0^{2\pi} e^{2u(\varrho_n e^{i\varphi})} \varrho_n d\varphi < 2\pi.$$

Now we apply Theorem 9, letting  $\gamma_n = [|z| = \varrho_n]$  and  $M = \sqrt{2\pi}$ . It follows that  $I(\Gamma_r) = 0$ . This completes the proof of Theorem 12.

We mention that there exist finitely connected, complete, open, two-dimensional RIEMANNIAN manifolds which belong to any prescribed topological type.

In order to construct such examples we take a parabolic RIEMANN surface  $S^{11)}$  which possesses the required topological structure. (Such a RIEMANN surface can always be obtained by removing a finite number of points from a suitable closed surface.) Then the  $\Omega_r$ 's are all conformally equivalent to schlicht circular rings of the type  $R_1 < |z| < +\infty$  ( $R_1 > 0$ ). In these we define the conformal metric

$$ds = \frac{|dz|}{|z| \log(|z| + 1)} .$$

On the remaining portion of  $S$  the metric is "filled out" arbitrarily, but such that it is everywhere positive definite and of class  $C^2$ . By making use of the fact that

$$\int_1^{+\infty} \frac{d\varrho}{\varrho \log(\varrho + 1)} = +\infty, \quad \text{but} \quad \int_1^{+\infty} \frac{d\varrho}{\varrho \log^2(\varrho + 1)} < +\infty ,$$

one verifies easily that this metric is complete, but of finite total area.

## 5. On complete conformal metrics defined on infinitely connected RIEMANN surfaces

**Theorem 13.** *Suppose that the conformal metric  $e^{u(z)}|dz|$ , defined on an infinitely connected RIEMANN surface  $S$ , is complete. Then  $C^- = +\infty$ .*

**Remark.** This result complements Theorem 10. It was suggested to us as a conjecture by Professor H. HOPF.

**Proof.** Assume that  $C^- < +\infty$ . We are going to show that  $e^{u(z)}|dz|$  cannot be complete under this hypothesis.

We exhaust  $S$  by an increasing sequence  $\{\Sigma_r\}$  of subcompacts, each being bounded by a finite number of analytic JORDAN curves  $(\beta_{r1}, \beta_{r2}, \dots, \beta_{rm_r})$  which we suppose to be free of mass. We may further request that each  $\beta_{rs}$  constitutes the boundary of exactly one component  $\Omega_{rs}$  of  $S - \Sigma_r$  ( $s = 1, 2, \dots, m_r$ ). By the GAUSS-BONNET formula (4.8)

$$2\pi\mu(\Sigma_r) + 2\pi\chi_r = \sum_{s=1}^{m_r} I(e^u|dz|, \beta_{rs}; B) , \quad (5.1)$$

where  $\chi_r$  denotes the EULER-POINCARÉ characteristic of  $\Sigma_r$  and  $B$  designates the (ideal) boundary of  $S$ . The left-hand side of (5.1) tends to  $-\infty$  as  $r \rightarrow \infty$ , since  $2\pi\mu(S) = -C < +\infty$ . Hence, for sufficiently large  $r$

$$\sum_{s=1}^{m_r} I(e^u|dz|, \beta_{rs}; B) < -4\pi\mu^+(S) .$$

<sup>11)</sup> i. e. a RIEMANN surface with nullboundary (cf. R. NEVANLINNA [24, p. 319]).

Then, for at least one index  $s$

$$I(e^u|dz|, \beta_{rs}; B) = -4\pi[\mu^+(\Omega_{rs}) + \eta] , \quad (5.2)$$

where  $\eta > 0$ . Let such  $r$  and  $s$  be chosen. We change the notation by writing  $\beta_1, \Omega, \mu_1$  and  $\mu_2$  instead of  $\beta_{rs}, \Omega_{rs}, \mu^+$  and  $\mu^-$ , respectively, and introduce an analytic JORDAN curve  $\delta_1$  in  $\Omega$  which is homotopic to  $\beta_1$ , free of mass and so close to  $\beta_1$  that  $\mu_2(\beta_1, \delta_1) < \eta$ .

**Lemma 6.** *There is a number  $C$  with the following property: Given an arbitrary index  $r$ , there exists a rectifiable curve  $\alpha$ , leading from  $\delta_1$  to the boundary of  $\Sigma_r$ , such that*

$$\int_{\alpha} e^u|dz| < C . \quad (5.3)$$

**Remark.** Theorem 13 follows immediately from this result by means of an obvious generalization of Lemma 2.

**Proof.** Our demonstration is similar to the one of Lemma 3. Let  $\gamma_1$  and  $\gamma_0$  be analytic JORDAN curves in  $(\beta_1, \delta_1)$ , both homotopic to  $\beta_1$  and free of mass,  $\gamma_0$  lying in  $(\gamma_1, \delta_1)$ . We introduce the notations  $\delta_2, \gamma_2$  and  $\beta_2$  for the respective intersections (non-empty for sufficiently large  $r$ ) of the boundaries of  $\Sigma_r, \Sigma_{r+1}$  and  $\Sigma_{r+2}$  with  $\Omega$ . Further, let  $\omega = (\beta_1, \beta_2)$ ,  $\omega_0 = (\beta_1, \gamma_0)$ ,  $\omega_1 = (\gamma_0, \beta_2)$ ,  $\mu_{20}(e) = \mu_2(e \cap \omega_0)$ ,  $\mu_{21}(e) = \mu_2(e \cap \omega_1)$ ,  $m_0 = \mu_2(\omega_0)$  and  $m_1 = \mu_2(\omega_1)$ . We have

$$m_0 \leq \mu_2(\beta_1, \delta_1) < \eta . \quad (5.4)$$

We define the metrics  $e^{h(z)}|dz|$ ,  $e^{v(z)}|dz|$ ,  $e^{v_1(z)}|dz|$  and  $e^{V(z)}|dz|$  by putting

$$h(z) = u(z) + \int_{\omega} g(z, \zeta) d\mu(e_{\zeta}) , \quad (5.5)$$

$$v(z) = u(z) + \int_{\omega} g(z, \zeta) d\mu_1(e_{\zeta}) \quad (5.6)$$

$$= h(z) + \int_{\omega} g(z, \zeta) d\mu_2(e_{\zeta}) ,$$

$$v_1(z) = h(z) + \int_{\omega} g(z, \zeta) d\mu_{21}(e_{\zeta}) \quad (5.7)$$

and

$$V(z) = u(z) + \int_{\omega} G(z, \zeta) d\mu_1(e_{\zeta}) , \quad (5.8)$$

where  $G$  and  $g$  denote the GREEN's functions of  $\Omega$  and  $\omega$ , respectively<sup>12)</sup>. Since  $0 \leq g \leq G$  throughout  $\omega$ ,  $v_1(z) \leq v(z) \leq V(z)$  and, consequently

$$\int_{\gamma_1} e^{v_1}|dz| \leq \int_{\gamma_1} e^{V(z)}|dz| = C_1 < +\infty .^{13)} \quad (5.9)$$

<sup>12)</sup> Cf. R. NEVANLINNA [24].

<sup>13)</sup>  $C_1, \dots, C_5$  are constants not depending on the choice of  $\beta_2, \gamma_2$  and  $\delta_2$ .

Furthermore, throughout  $(\delta_1, \delta_2)$ ,  $u(z) \leq v(z) \leq v_1(z) + C_2$ , where  $C_2$  denotes the (finite) upper bound of

$$\int_{\omega_0} G(z, \zeta) d\mu_{20}(e_\zeta) \quad \text{for } z \text{ varying on } \Omega - (\beta_1, \delta_1) .$$

(In the case where  $\mu_{20}$  is concentrated in one point the existence of such a bound is an immediate consequence of the properties of  $G$ . One proceeds to general measures  $\mu_{20}$  by an application of HÖLDER's inequality.) Hence, for every curve  $\alpha$  in  $(\delta_1, \delta_2)$

$$\left| \int_{\alpha} e^u dz \right| \leq C_3 \left| \int_{\alpha} e^{v_1} dz \right| . \quad (5.10)$$

We are now going to prove that there exists a rectifiable curve  $\alpha$ , leading from  $\delta_1$  to  $\delta_2$ , such that

$$\left| \int_{\alpha} e^{v_1} dz \right| \leq C_4 \left| \int_{\gamma_1} e^{v_1} dz \right| + C_5 . \quad (5.11)$$

Lemma 6 is an immediate consequence of (5.9), (5.10) and (5.11).

In order to establish (5.11) we first approximate the measure  $\mu_{21}$  by a finite number of concentrated masses. This is the purpose of the following construction.

Consider a triangulation  $T_0$  of the closure  $\bar{\omega}$  of  $\omega$ , consisting of the triangles  $\Delta_k^{014}$  ( $k = 1, 2, \dots, M$ ) whose boundaries we suppose to be analytic and free of mass.

We subdivide  $T_0$  in the following way: There exists a conformal representation  $t_k = \varphi_k(z)$  of the interior of  $\Delta_k^0$  onto the equilateral triangle  $E$  ( $1/2, -1/2, i\sqrt{3}/2$ ) in a  $t_k$ -plane such that the vertices of  $\Delta_k^0$  correspond to those of  $E$  ( $k = 1, 2, \dots, M$ ). (In the following the letters  $z$  and  $\zeta$  are used to designate points on  $S$  whereas  $t_k$  and  $\tau_k$  denote the corresponding values of the just introduced uniformizers.) We join the mid-points of the sides of  $E$ , thus breaking it up into four smaller triangles. To this subdivision of  $E$  there corresponds a triangulation  $T_1$  of  $\bar{\omega}$ , consisting of the triangles

$$\Delta_{kj}^1 \quad (k = 1, 2, \dots, M; j = 1, 2, 3, 4) .$$

By iterating the subdivision of  $E$  we obtain a sequence  $\{T_n\}$  of triangulations, each  $T_n$  being composed of  $4^n M$  triangles  $\Delta_{kj}^n$  ( $k = 1, 2, \dots, M; j = 1, 2, \dots, 4^n$ ). We are going to select an index  $n$  large enough for our purpose.

We define a subcompact  $K_1$  of  $\bar{\omega}$  by subtracting neighborhoods of all vertices of  $T_0$ . We require that these neighborhoods be bounded by analytic

---

<sup>14</sup>) By  $\Delta_k^0$  and  $\Delta_{kj}^n$  (below) we understand the *closures* of the respective triangles.

curves (let  $\Gamma_1$  denote their totality) such that

$$\int_{\Gamma_1 \cap \bar{\omega}} e^{v_1} |dz| < 1. \quad (5.12)$$

Further, let  $K_2$  be a second compact of the same type, containing  $K_1$  and satisfying the condition

$$\mu_{21}(\omega - K_2)g(z, \zeta) < \log 2 \quad (5.13)$$

for all  $z \in K_1$  and  $\zeta \in (\omega - K_2)$ . It is always possible to fulfill the conditions (5.12) and (5.13), since, by hypothesis, the vertices of  $T_0$  support no mass.

The region of definition of the conformal representation  $t_k = \varphi_k(z)$  can be extended to include an open set  $O_k$  containing  $\Delta_k^0 \cap K_2$ . Let  $G_k$  denote a region which also contains  $\Delta_k^0 \cap K_2$  and whose closure lies in  $O_k$  ( $k = 1, 2, \dots, M$ ).  $G_k$ 's belonging to adjacent triangles intersect. It can be inferred from the construction that there exists a number  $A$  ( $1 \leq A < +\infty$ ) such that, uniformly in  $k$  and  $l$

$$\frac{1}{A} \leq \left| \frac{dt_k}{dt_l} \right| \leq A \quad (5.14)$$

throughout  $G_k \cap G_l$ .

We define the notion of  $N$ -neighborhood ( $N = 0, 1, 2, \dots$ ) of a triangle  $\Delta_{kj}^n$  by recursion as follows: The 0-neighborhood is identical with  $\Delta_{kj}^n$ . The  $N$ -neighborhood of  $\Delta_{kj}^n$  consists of those (closed) triangles  $\Delta_{k'j'}^n$  which intersect the  $(N-1)$ -neighborhood of  $\Delta_{kj}^n$  ( $N = 1, 2, 3, \dots$ ).

The following property is obvious:

(I) If  $\Delta_{k'j'}^n$  lies in the  $N$ -neighborhood of  $\Delta_{kj}^n$ , then  $\Delta_{kj}^n$  belongs to the  $N$ -neighborhood of  $\Delta_{k'j'}^n$ .

Given an arbitrary positive integer  $N$ , there always exists an index  $n_0(N)$  such that the following condition is fulfilled for all  $k, j$  and any  $n > n_0(N)$ : If the  $N$ -neighborhood of  $\Delta_{kj}^n$  intersects  $K_2$ , then it lies in  $G_k$  and overlaps from  $\Delta_k^0$  into at most one  $\Delta_l^0$  ( $l \neq k$ ). In this case every such  $N$ -neighborhood (considered in the  $t_k$ -plane) is contained in a circle of radius  $(N+1)A/n$ . Since, on the other hand, the area of every composing triangle is  $\geq \sqrt{3}/4 A^2 n^2$ , we conclude:

(II) Each  $N$ -neighborhood intersecting  $K_2$  contains less than  $8(N+1)^2 A^4$  triangles  $\Delta_{kj}^n$ .

The statements (I) and (II) imply:

(III) Each  $\Delta_{kj}^n$  intersecting  $K_2$  is contained in less than  $8(N+1)^2 A^4$  different  $N$ -neighborhoods.

We choose an integer  $N_1 > A + 1$  and introduce the abbreviation  $U_{kj}^n$  for the  $N_1$ -neighborhood of  $\Delta_{kj}^n$ . It is easy to verify that the following is true for all  $n > n_0(N_1)$ :

(IV) Let  $\varrho$  be a rectifiable curve on  $S$ , leading from  $z$  to  $\zeta$ , both points lying in  $\Delta_{kj}^n$ . Suppose that  $\Delta_{kj}^n \cap K_2 \neq 0$ . Then  $\varrho \cap U_{kj}^n$  has at least the length<sup>15)</sup>  $|t_k - \tau_k|$ .

We state that there exists an integer  $N_2 > N_1 + 1$  with the following property:

(V) Let  $V_{kj}^n$  denote the  $N_2$ -neighborhood of  $\Delta_{kj}^n$ . Suppose that  $\Delta_{kj}^n \cap K_2 \neq 0$ . Assume further that  $n > n_0(N_2)$ . Then the inequality

$$m_1 |g(z, \zeta) - g(z', \zeta')| < \log 2 \quad (5.15)$$

holds for arbitrary  $z, z' \in \Delta_{kj}^n$  and  $\zeta, \zeta' \in \Delta_{k'j'}^n$ , admitting any  $\Delta_{k'j'}^n$  which is not contained in  $V_{kj}^n$ .

It is sufficient to prove the existence of  $N_2$  for a fixed index  $k$ . Throughout  $G_k$  we have the representation

$$g(z, \zeta) = \log \frac{1}{|t_k - \tau_k|} + r_k(t_k, \tau_k), \quad (5.16)$$

where  $r_k$  is a regular function. We first limit ourselves to those triangles  $\Delta_{k'j'}^n$  which lie in  $G_k$ . If  $n$  is large enough, then, by the continuity of  $r_k$

$$m_1 |r_k(t_k, \tau_k) - r_k(t'_k, \tau'_k)| < \frac{\log 2}{2} \quad (5.17)$$

for  $z, z' \in \Delta_{kj}^n$  and  $\zeta, \zeta' \in \Delta_{k'j'}^n \subset G_k$ . Since  $n_0(N_2) \rightarrow \infty$  as  $N_2 \rightarrow \infty$ , (5.17) is fulfilled for all  $n > n_0(N_2)$  if only  $N_2$  is chosen large enough. Furthermore, the diameter<sup>15)</sup> of any  $\Delta_{k'j'}^n$  in  $G_k$  is at most  $A/n$ , whereas the distance<sup>15)</sup> between  $U_{kj}^n$  and the boundary of  $V_{kj}^n$  is at least  $(N_2 - N_1 - 1)/An$ . Hence, if  $N_2$  is sufficiently large, then

$$m_1 \log \left| \frac{t_k - \tau_k}{t'_k - \tau'_k} \right| \leq m_1 \log \frac{\frac{N_2 - N_1 - 1}{An} + \frac{2A}{n}}{\frac{N_2 - N_1 - 1}{An} - \frac{2A}{n}} \leq \frac{\log 2}{2} \quad (n > n_0(N_2)) \quad (5.18)$$

for  $z, z' \in U_{kj}^n$  and  $\zeta, \zeta' \in \Delta_{k'j'}^n$ , where  $\Delta_{k'j'}^n$  is supposed to lie in  $G_k$  but not in  $V_{kj}^n$ . (5.15) follows from (5.16), (5.17) and (5.18). We have yet to treat the case where  $\Delta_{k'j'}^n$  intersects  $\omega - G_k$ . But then (5.15) (for sufficiently large  $n$ ) is an obvious consequence of the continuity of  $g$ .

From now on  $N_1$  and  $N_2$  are to be considered fixed.  $n$  is still variable. Let  $\zeta_1, \zeta_2, \dots, \zeta_m$  designate those points which support a concentrated mass of weight  $\geq \frac{1}{32(N_2 + 1)^2 A^4}$  in the measure  $\mu_{21}(e \cap K_2)$ . We denote the corresponding masses by  $p_1, p_2, \dots, p_m$  and define the metric  $e^{w(z)}|dz|$  by putting

<sup>15)</sup> with respect to the Euclidean metric in the  $t_k$ -plane.

$$w(z) = h(z) + \sum_{l=1}^m p_l g(z, \zeta_l) . \quad (5.19)$$

We enclose each  $\zeta_l$  by two analytic JORDAN curves  $\varepsilon_l$  and  $\varepsilon'_l$  which satisfy the following conditions:

(a)  $\varepsilon'_l$  is contained in the interior of  $\varepsilon_l$  ( $l = 1, 2, \dots, m$ );

(b)  $\gamma_1, \varepsilon_1, \varepsilon_2, \dots, \varepsilon_m$  do not intersect;

(c) whenever  $\zeta_l$  is a singular point of  $\mu$  (i. e.  $p_l \geq 1$ ), then  $\varepsilon_l$  shall be so close to  $\zeta_l$  that

$$\int_{\kappa} e^w |dz| > \frac{4}{1 - \cos(\pi\eta)} \int_{\gamma_1} e^{v_1} |dz| \quad (5.20)$$

for every rectifiable curve  $\kappa$  leading from  $\gamma_1$  to  $\varepsilon_l$ ;

$$(d) \quad \sum_{p_l < 1} \int_{\varepsilon_l} e^{v_1} |dz| < 1 . \quad (5.21)$$

Let  $\nu(e)$  denote the measure which originates from  $\mu_{21}(e \cap K_2)$  after removal of the concentrated masses  $p_1$  in  $\zeta_1, \dots, p_m$  in  $\zeta_m$ . For all  $z \in K_1$  we have, by (5.13) and (5.19)

$$v_1(z) = v_2(z) + \int_{\omega - K_2} g(z, \zeta) d\mu_{21}(e_\zeta) \leq v_2(z) + \log 2 , \quad (5.22)$$

where

$$v_2(z) = w(z) + \int_{\omega} g(z, \zeta) d\nu(e_\zeta) . \quad (5.23)$$

Let  $C$  designate the compact which is obtained by subtracting the interiors of  $\varepsilon'_1, \varepsilon'_2, \dots, \varepsilon'_l$  from the closure of  $(\gamma_1, \gamma_2)$ .

Now we choose  $n$ , large enough so that conditions (5.24) to (5.30) are satisfied for all  $k, j$  and  $l$ :

$$n > n_0(N_2) ; \quad (5.24)$$

$$\Delta_{kj}^n \cap \varepsilon_l \neq 0 \quad \text{implies} \quad U_{kj}^n \cap \varepsilon'_l = 0 ; \quad (5.25)$$

$$\nu(\Delta_{kj}^n) \leq \frac{1}{32(N_2 + 1)^2 A^4} ; \quad (5.26)$$

$$m_1 |g(z, \zeta_1) - g(z, \zeta_2)| < \log 2 \quad (5.27)$$

for all  $z \in \gamma_1$  and  $\zeta_1, \zeta_2 \in \Delta_{kj}^n$ , provided that  $\Delta_{kj}^n \cap \omega_1 \neq 0$ ;

$$|w(t_k) - w(\tau_k)| < \log 2 \quad (5.28)$$

and

$$|r_k(t_k, \tau_k) - r_k(t'_k, \tau'_k)| < \log 2 , \quad (5.29)$$

whenever  $t_k, \tau_k, t'_k$  and  $\tau'_k$  lie in the same  $V_{kj}^n$ .

There exists a positive number  $L$  such that, for all  $z \in \omega$

$$\nu(E[\zeta | g(z, \zeta) > L]) < \frac{1}{4} .$$

Let

$$W = \max_k \left[ \sup_{t_k \in G_k \cap C} w(t_k) \right] \quad \text{and} \quad R = \max_k \left[ \sup_{t_k, \tau_k \in G_k} r_k(t_k, \tau_k) \right] .$$

We postulate that, for  $0 < \lambda \leq 1/n$

$$\frac{2^9 m_1 A^{\frac{17}{4}} (N_2 + 1)^2}{(N_2 - N_1 - 1)^{\frac{1}{4}}} e^{W+R+m_1 L} \cdot \lambda^{\frac{1}{2}} |\log \lambda| < 1 . \quad (5.30)$$

On each  $\Delta_{kj}^n$  intersecting  $\omega_1$  there is a point  $\zeta_{\min}(\Delta_{kj}^n)$  such that

$$\int_{\gamma_1} \frac{\partial g(z, \zeta_{\min})}{\partial n_z} |dz| \leq \int_{\gamma_1} \frac{\partial g(z, \zeta)}{\partial n_z} |dz| \quad (5.31)$$

for all  $\zeta \in \Delta_{kj}^n$ ,  $n_z$  denoting the normal to  $\gamma_1$  which points into  $(\gamma_1, \delta_1)$ .

Now we concentrate in each  $\zeta_{\min}(\Delta_{kj}^n)$  the mass  $\nu$  which is associated with the interior of  $\Delta_{kj}^n$  and part of its boundary, defining the latter such that every point is covered exactly once. Let these be the masses  $p_{m+1}$  in  $\zeta_{m+1}, \dots, p_r$  in  $\zeta_r$ . We introduce the metric  $e^{w_1(z)} |dz|$ , where

$$w_1(z) = h(z) + \sum_{l=1}^r p_l g(z, \zeta_l) = w(z) + \sum_{l=m+1}^r p_l g(z, \zeta_l) . \quad (5.32)$$

Let  $\gamma$  be a JORDAN curve in  $\omega$  which is homotopic to  $\beta_1$ . We state that

$$I(e^{w_1} |dz|, \gamma; \beta_2) < -2\pi\eta . \quad (5.33)$$

In order to prove this inequality we first observe that, by the GAUSS-BONNET formula

$$I(e^{w_1} |dz|, \gamma; \beta_2) \leq I(e^{w_1} |dz|, \gamma_1; \beta_2) . \quad (5.34)$$

From (5.31) and the construction of  $w_1$  we infer that

$$I(e^{w_1} |dz|, \gamma_1; \beta_2) \leq I(e^{v_2} |dz|, \gamma_1; \beta_2) . \quad (5.35)$$

By (4.8), (5.22), (5.6) and (5.7)

$$\begin{aligned} I(e^{v_2} |dz|, \gamma_1; \beta_2) &= I(e^{v_2} |dz|, \gamma_0; \beta_2) \\ &\leq I(e^{v_1} |dz|, \gamma_0; \beta_2) \leq I(e^v |dz|, \gamma_0; \beta_2) + 2\pi m_0 \\ &\leq I(e^u |dz|, \gamma_0; \beta_2) + 2\pi[m_0 + \mu_1(\Omega)] . \end{aligned} \quad (5.36)$$

From (4.8) and (5.2) we infer that

$$I(e^u|dz|, \gamma_0; \beta_2) = I(e^u|dz|, \beta_1; \beta_2) + 2\pi\mu(\beta_1, \gamma_0) < -2\pi[\mu_1(\Omega) + 2\eta] . \quad (5.37)$$

(5.33) is implied by relations (5.4) and (5.34) to (5.37).

We state that there exists a rectifiable curve  $\beta$ , leading from  $\gamma_1$  to  $\delta_2$ , such that

$$\int_{\beta} e^{w_1}|dz| < \frac{2}{1 - \cos(\pi\eta)} \int_{\gamma_1} e^{w_1}|dz| . \quad (5.38)$$

The proof of this inequality is so similar to the one of (2.39) that we do not reproduce it here. We limit ourselves to the following remarks:

(a) In the definition

$$\Lambda(z_0) = \inf_{\gamma} \int_{\gamma} e^{w_1}|dz|$$

we admit to competition all rectifiable JORDAN curves  $\gamma$  which lie in the closure of  $(\gamma_1, \gamma_2)$ , pass through  $z_0$  and are homotopic to  $\beta_1$ . Then (2.44) holds again for every minimal curve  $\bar{\gamma}(z_0)$  which has neither double points nor points in common with  $\gamma_2$ . This follows from (5.33).

(b)  $\beta$  can be constructed as a "polygon", i. e. a contiguous chain of "straight line segments". Thereby a "straight line segment"  $\sigma$  in  $\omega$  is defined to be a smooth curve with the property that, for all  $k$ , the set  $\varphi_k(\sigma \cap \Delta_k^0)$  in the  $t_k$ -plane consists of (Euclidean) straight line segments.

(c)  $\beta$  does not intersect any  $\varepsilon_i$  for which  $p_i \geq 1$ . Indeed, by (5.32)

$$\int_{\beta} e^w|dz| \leq \int_{\beta} e^{w_1}|dz| . \quad (5.39)$$

From (5.27) and (5.22) we infer that

$$\int_{\gamma_1} e^{w_1}|dz| \leq 2 \int_{\gamma_1} e^{v_2}|dz| \leq 2 \int_{\gamma_1} e^{v_1}|dz| . \quad (5.40)$$

(5.38), (5.39) and (5.40) imply the inequality

$$\int_{\beta} e^w|dz| < \frac{4}{1 - \cos(\pi\eta)} \int_{\gamma_1} e^{v_1}|dz| , \quad (5.41)$$

which would contradict (5.20) if  $\beta$  would intersect any  $\varepsilon_i$  for which  $p_i \geq 1$ .

With the "polygon"  $\beta$  we now associate a curve  $\alpha$  which leads from  $\delta_1$  to  $\delta_2$ . This is done by the following construction: Let  $E(z)$  denote the set consisting of those  $\Delta_{k_i}^n$ 's which contain the point  $z$ . We join the endpoint  $z_0$  of  $\beta$  on  $\gamma_1$  with the last point of intersection  $z'_1$  of  $\beta$  with  $E(z_0)$  by a "straight line segment",  $z'_1$  with the last point of intersection  $z'_2$  of  $\beta$  with  $E(z'_1)$ , and so forth,

until we arrive at the endpoint of  $\beta$  on  $\delta_2$ . We obtain a "polygon"  $\beta'$ . Now, if  $\beta'$  should penetrate into  $\bar{\omega} - K_1$  or into the interiors of some curves  $\varepsilon_i$  for which  $p_i < 1$ , then we replace the "subpolygon" between the first entry and the last exit by a boundary arc. The resulting curve contains a portion  $\alpha$  which leads from  $\delta_1$  to  $\delta_2$  and is contained in the closure of  $(\delta_1, \delta_2)$ . We state that

$$\int_{\alpha} e^{v_2} |dz| \leq \frac{32}{3} \int_{\beta} e^{w_1} |dz| + 3. \quad (5.42)$$

We decompose  $\alpha = \alpha_P + \alpha_C$ ,  $\alpha_P$  consisting of a finite number of "polygons",  $\alpha_C$  of the detours introduced above. By (5.12), (5.21) and (5.22)

$$\int_{\alpha_C} e^{v_2} |dz| < 2.$$

We are left to show that

$$\int_{\alpha_P} e^{v_2} |dz| \leq \frac{32}{3} \int_{\beta} e^{w_1} |dz| + 1. \quad (5.43)$$

The verification of this inequality is quite analogous to the proof of (2.65). We leave it to the reader and limit ourselves to the following remarks:

(a) From (5.26) and property (II) of  $N$ -neighborhoods we infer

$$\nu(V_{kj}^n) \leq \frac{1}{4}. \quad (5.44)$$

Every point  $z$  in  $K_2$  belongs to at most 6  $\Delta_{kj}^n$ 's. From this and property (III) of  $N$ -neighborhoods we conclude that

$$\sum_{k,j} \nu(V_{kj}^n) \leq 48 m_1 (N_2 + 1)^2 A^4. \quad (5.45)$$

Relations (5.44) and (5.45) correspond to (2.57) and (2.58), respectively.

(b) In the estimations corresponding to (5.59), (5.60) and (5.61) it is convenient to integrate in the plane of the respective uniformizer  $t_k$ . One makes use of the decomposition (5.16) of GREEN's function. The logarithmic term is handled in the same way as in the proof of (2.65). The additional function  $r_k$  is taken care of by relation (5.29) and the fact that  $R$  occurs in (5.30).

Inequality (5.11) follows from (5.22), (5.42), (5.38) and (5.40). We have thus proved Lemma 6 and, with it, Theorem 13.

## 6. Further results

**Theorem 14.** *Let  $S$  be an open RIEMANN surface on which a complete conformal metric  $e^{u(z)} |dz|$  is defined. Suppose that the measure  $\mu^+$  has a compact support. Then the total area  $A = \iint_S e^{2u} dx dy$  is infinite.*

**Remark.** For manifolds which possess a continuous GAUSSIAN curvature  $K$  our hypothesis simply means that  $K \geq 0$  outside a compact subdomain. We

mention that Theorem 14 has already been demonstrated by F. FIALA (Theorem A, p. 300 in [12]) for (necessarily simply connected) analytic manifolds whose curvature is everywhere non-negative.

**Proof.** Since  $C^- = 2\pi\mu^+(S) < +\infty$ , we conclude from Theorem 13 that  $S$  is finitely connected. Furthermore, the curvatura integra  $C$  exists. Consequently<sup>16</sup>,  $I(\Gamma_r) \geq 0$  for  $r = 1, 2, \dots, N$ . Consider an arbitrary one of the regions  $\Omega_r$ . It is conformally equivalent to a schlicht circular ring  $R_1 < |z| < R_2$  ( $0 < R_1 < R_2 \leq +\infty$ ). Again we distinguish between two cases:

(I)  $R_2 < +\infty$ . It has been verified in the proof of Theorem 12 that the completeness of the metric yields indeed  $A = +\infty$ . (Actually this case does not occur at all. For,  $R_2 < +\infty$  implies that  $S$  is hyperbolic, and it will be shown in Theorem 15 that this is incompatible with  $C^- < +\infty$ .)

(II)  $R_2 = +\infty$ . We may assume that  $\Omega_r$  does not intersect the support of  $\mu^+$ . (If necessary we increase  $R_1$ .) Then  $u$  is superharmonic throughout  $\Omega_r$ . Since  $I(\Gamma_r) \geq 0$ , we have, by (4.7)

$$\lim_{\varrho \rightarrow \infty} \Phi(u, |z| = \varrho; \Gamma_r) = \Phi(\Gamma_r) \geq -1. \quad (6.1)$$

But,  $u$  being superharmonic,  $\Phi(u, |z| = \varrho; \Gamma_r)$  is a non-increasing function of  $\varrho$ . Hence (6.1) implies

$$\Phi(u, |z| = \varrho; \Gamma_r) \geq -1 \quad (R_1 < \varrho < +\infty). \quad (6.2)$$

Furthermore

$$\int_0^{2\pi} u(\varrho_2 e^{i\varphi}) d\varphi - \int_0^{2\pi} u(\varrho_1 e^{i\varphi}) d\varphi = 2\pi \int_{\varrho_1}^{\varrho_2} \Phi(u, |z| = \varrho; \Gamma_r) d \log \varrho, \quad (6.3)$$

where  $R_1 \leq \varrho_1 \leq \varrho_2 < +\infty$ . In the case of sufficiently regular  $u$  this relation can be verified immediately by a direct calculation. It is more generally true (and essentially known) for all functions  $u$  which admit the decomposition (1.9). (The reader looking for a proof will find section 5.14, p. 35 in [25], helpful.) From (6.2) and (6.3) we infer that

$$\frac{1}{2\pi} \int_0^{2\pi} u(\varrho e^{i\varphi}) d\varphi \geq B_1 - \log \varrho \quad (6.4)$$

for some real constant  $B_1$  and arbitrary  $\varrho$  ( $R_1 \leq \varrho < +\infty$ ). By making use of the theorem of the arithmetic and geometric means [15, p. 137] we obtain, for arbitrary  $\varrho$

$$\int_0^{2\pi} e^{2u(\varrho e^{i\varphi})} d\varphi \geq 2\pi e^{\frac{2}{2\pi} \int_0^{2\pi} u(\varrho e^{i\varphi}) d\varphi} \geq \frac{B_2}{\varrho^2}, \quad (6.5)$$

<sup>16</sup>) See proof of Theorem 10.

$B_2$  denoting a positive constant. (6.5) yields

$$A > \int_{R_1}^{+\infty} \int_0^{2\pi} e^{2u(\varrho e^{i\varphi})} \varrho d\varrho d\varphi \geq B_2 \int_{R_1}^{+\infty} \frac{d\varrho}{\varrho} = +\infty. \quad \text{Q. E. D.}$$

**Theorem 15.** *If an open RIEMANN surface  $S$  admits a complete conformal metric  $e^{u(z)}|dz|$  with finite  $C^-$ , then it is parabolic.*

**Remark.** This result is known in the simply connected case, where it has been proved by CH. BLANC and F. FIALA [5].

**Proof.** We assume that  $S$  is hyperbolic and show that this leads to a contradiction.

Under this hypothesis  $S$  would possess a GREEN's function (cf. P. J. MYRBERG [21], R. NEVANLINNA [24, chapter 10]). Consider the conformal metric  $e^{v(z)}|dz|$ , where

$$v(z) = u(z) + 2g(z, \zeta_0) + \int_S g(z, \zeta) d\mu^+(e_\zeta). \quad (6.6)$$

Here  $\zeta_0$  denotes an arbitrary, but fixed point on  $S$ . The integral on the right-hand side of (6.6) is not identically infinite, since  $\mu^+(S) = C^-/2\pi < +\infty$ . One verifies easily that  $C^+ \geq 4\pi$  and  $C^- = 0$  for this metric. But since always  $\chi \leq 1$  we conclude from Theorems 10 and 13 that  $e^{v(z)}|dz|$  cannot be complete. Hence  $e^{u(z)}|dz|$  would not be complete either, contrary to hypothesis.

We observe that the assumption  $C^- < +\infty$  has only been used for the purpose of showing that the integral in (6.6) is not identically infinite. Hence we have actually proved a statement which is slightly stronger than Theorem 15. *Let  $S$  be a hyperbolic RIEMANN surface carrying a complete conformal metric  $e^{u(z)}|dz|$  whose curvatura integra  $C$  may or may not exist. Then*

$$\int_S g(z, \zeta) d\mu^+(e_\zeta) \equiv +\infty,$$

where  $g$  denotes GREEN's function for  $S$ . In the case of infinitely connected  $S$  Theorem 15 is obviously superseded by Theorem 13, whereas the stronger result contains new information.

The author wishes to thank Mr. J. H. BRAMBLE, M. A., for helping with the English translation.

#### REFERENCES

- [1] B. ANDERSSON, *On an inequality concerning the integrals of moduli of regular analytic functions*, Arkiv Mat. 1 (1951) 367–373, No. 27.
- [2] M. G. ARSOVE, *Functions representable as differences of subharmonic functions*, Trans. Amer. Math. Soc. 75 (1953) 327–365.
- [3] E. F. BECKENBACH and T. RADÓ, *Subharmonic functions and surfaces of negative curvature*, Trans. Amer. Math. Soc. 35 (1933) 662–674.
- [4] A. BEURLING, *Sur la géométrie métrique des surfaces à courbure totale  $\leq 0$* , Meddelanden Lunds Univ. Mat. Sem., Supplementband 1952 (Marcel Riesz Anniv. Vol.) 7–11.

- [5] CH. BLANC et F. FIALA, *Le type d'une surface et sa courbure totale*, Comment. Math. Helv. 14 (1941–42) 230–233.
- [6] W. BLASCHKE, *Vorlesungen über Differentialgeometrie*, Bd. 1, Springer, Berlin 1924.
- [7] C. CARATHÉODORY, *Funktionentheorie*, Bd. 1, Birkhäuser, Basel 1950.
- [8] F. CARLSON, *Quelques inégalités concernant les fonctions analytiques*, Arkiv Mat., Astr. Fysik 29 B, No. 11 (1943).
- [9] S. S. CHERN, P. HARTMAN and A. WINTNER, *On isothermic coordinates*, Comment. Math. Helv. 28 (1954) 301–309.
- [10] S. COHN-VOSSEN, *Kürzeste Wege und Totalkrümmung auf Flächen*, Compositio Math. 2 (1935) 69–133.
- [11] L. FEJÉR und F. RIESZ, *Über einige funktionentheoretische Ungleichungen*, Math. Z. 11 (1921) 305–314.
- [12] F. FIALA, *Le problème des isopérimètres sur les surfaces ouvertes à courbure positive*, Comment. Math. Helv. 13 (1940–41) 293–346.
- [13] R. M. GABRIEL, *Some results concerning the integrals of moduli of regular functions along curves of certain types*, Proc. London Math. Soc. (II), 28 (1928) 121–127.
- [14] C. GATTEGNO et A. OSTROWSKI, *Représentation conforme à la frontière; domaines généraux*, Mém. sci. math., fasc. 109, Paris 1949.
- [15] G. H. HARDY, J. E. LITTLEWOOD and G. PÓLYA, *Inequalities*, second edition, Cambridge University Press 1952.
- [16] H. HOPF, *Sulla geometria globale delle superficie*, Rendiconti del Seminario Matematico e Fisico di Milano 23 (1952) 48–63.
- [17] H. HOPF und W. RINOW, *Über den Begriff der vollständigen differentialgeometrischen Fläche*, Comment. Math. Helv. 3 (1931) 209–225.
- [18] A. HUBER, *On the isoperimetric inequality on surfaces of variable GAUSSIAN curvature*, Ann. Math. 60 (1954) 237–247.
- [19] A. HUBER, *On an inequality of FEJÉR and RIESZ*, Ann. Math. 63 (1956) 572–587.
- [20] B. v. KERÉKJÁRTÓ, *Vorlesungen über Topologie I*, Springer, Berlin 1923.
- [21] P. J. MYRBERG, *Über die Existenz der GREENSchen Funktion auf einer gegebenen RIEMANNschen Fläche*, Acta Math. 61 (1933) 39–79.
- [22] R. NEVANLINNA, *Beitrag zur Theorie der ABELSchen Integrale*, Ann. Acad. Sci. Fenn., Ser. AI, No. 100 (1951).
- [23] R. NEVANLINNA, *Eindeutige analytische Funktionen*, 2. Aufl., Springer 1953.
- [24] R. NEVANLINNA, *Uniformisierung*, Springer, Berlin 1953.
- [25] T. RADÓ, *Subharmonic functions*, Ergebnisse der Mathematik und ihrer Grenzgebiete V1, Springer, Berlin 1937.
- [26] F. RIESZ, *Über die Randwerte einer analytischen Funktion*, Math. Z. 18 (1923) 87–95.
- [27] F. RIESZ, *Sur les fonctions subharmoniques et leur rapport à la théorie du potentiel*, part I and II, Acta Math. 48 (1926) 329–343 and 54 (1930) 321–360.
- [28] F. und M. RIESZ, *Über die Randwerte einer analytischen Funktion*, Quatrième congrès des mathématiciens scandinaves à Stockholm (1916) 27–44.
- [29] M. RIESZ, *Remarque sur les fonctions analytiques*, Acta Sci. Math. Szeged 12 (1950) 53–56.
- [30] S. SAKS, *Theory of the integral*, Second edition, Hafner, New York 1937.
- [31] M. SCHIFFER and D. C. SPENCER, *Functionals of finite RIEMANN surfaces*, Princeton University Press 1954.
- [32] H. WEYL, *Über die Bestimmung einer geschlossenen konvexen Fläche durch ihr Linienelement*, Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich 61 (1916) 40–72.
- [33] H. WEYL, *Die Idee der RIEMANNschen Fläche*, Teubner 1923.
- [34] A. WINTNER, *On the local rôle of the logarithmic potential in differential geometry*, Amer. J. Math. 75 (1953) 679–690.

(Eingegangen 25. September 1956.)