

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 35 (1961)

Artikel: Projectively Flat Spaces with Recurrent Curvature.
Autor: Wong, Yung-Chow / Yano, Kentaro
DOI: <https://doi.org/10.5169/seals-27343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Projectively Flat Spaces with Recurrent Curvature

To the University of Hong Kong on its Golden Jubilee in 1961

By YUNG-CHOW WONG and KENTARO YANO

Introduction

Let A_N be an affinely connected N -dimensional space with a symmetric connection (i.e. a connection without torsion). A_N is a *projectively flat space*, or simply, a P_N if there exists a coordinate system in terms of which the finite equations of the paths are linear. A_N is of *recurrent curvature*¹⁾ if the covariant derivative of its curvature tensor is the tensor product of a non-zero covariant vector and the curvature tensor itself.

The purpose of this paper is to determine all the projectively flat spaces with recurrent curvature. For convenience we shall denote such a space by P_N^* . It is found that the space P_N^* (or rather its connection) depends on 2 arbitrary functions of one variable or on 3 arbitrary functions of one variable according as its RICCI tensor is symmetric or non-symmetric. The actual construction of the connection of the P_N^* depends on the solution of a differential equation of the RICCATI type and on the solution of a completely integrable system of differential equations.

Projectively flat space with covariantly constant curvature tensor (i.e. projectively flat symmetric space) is also considered. We prove that it is a well-known type of projectively flat space characterized by its RICCI tensor being symmetric and covariantly constant.

1. Preliminaries²⁾

Throughout this paper, each of the indices a, h, i, j, \dots runs through the range $1, \dots, N$; ∂_k denotes partial differentiation with respect to the k th coordinate; and a repeated index implies summation.

Let A_N be a linearly connected N -dimensional space with a symmetric connection Γ_{ji}^h (i.e. $\Gamma_{ji}^h = \Gamma_{ij}^h$). The curvature tensor, the RICCI tensor, and the tensor P_{ji} are defined respectively by:

$$R_{kji}^h = \partial_k \Gamma_{ji}^h - \partial_j \Gamma_{ki}^h + \Gamma_{ka}^h \Gamma_{ji}^a - \Gamma_{ja}^h \Gamma_{ki}^a, \quad (1.1)$$

$$R_{ji} = R_{a ji}^a, \quad (1.2)$$

¹⁾ RIEMANNIAN spaces with recurrent curvature have been studied in great detail by H. S. RUSE [2] and A. G. WALKER [3]. Certain classes of non-RIEMANNIAN spaces with recurrent curvature have been studied by Y. C. WONG [4].

²⁾ See EISENHART [1], but our notation is slightly different from his.

$$P_{ji} = -\frac{N}{N^2-1} R_{ji} - \frac{1}{N^2-1} R_{ij}. \quad (1.3)$$

It follows from (1.3) that P_{ji} is symmetric iff the RICCI tensor R_{ji} is symmetric.

By definition, A_N is of *recurrent curvature* if its curvature tensor satisfies the condition

$$\nabla_i R_{kji}{}^h = r_i R_{kji}{}^h, \quad (1.4)$$

where ∇ denotes the covariant differentiation and r_i a non-zero covariant vector. If

$$\nabla_i R_{kji}{}^h = 0, \quad (1.5)$$

the space A_N is said to be *symmetric*.

A_N is *projectively flat* iff there exists a coordinate system in terms of which the components Γ_{ji}^h have the form

$$\Gamma_{ji}^h = \phi_j A_i^h + \phi_i A_j^h, \quad (1.6)$$

where A_i^h is the KRONECKER delta and ϕ_j a set of N functions. The special form (1.6) for Γ_{ji}^h is preserved by affine transformations of coordinates. Expressed in terms of P_{ji} , a well-known necessary and sufficient condition for A_N to be projectively flat is

$$\left. \begin{aligned} R_{kji}{}^h + A_k^h P_{ji} - A_j^h P_{ki} - (P_{kj} - P_{jk}) A_i^h &= 0, \\ \nabla_k P_{ji} - \nabla_j P_{ki} &= 0. \end{aligned} \right\} \quad (1.7)$$

For $N > 2$, (1.7)₂ is a consequence of (1.7)₁; for $N = 2$, (1.7)₁ is an identity. There exists, for (1.6), a function ϕ such that $\phi_j = \partial_j \phi$ iff $P_{ji} = P_{ij}$. Since on account of (1.7) a projectively flat space with $P_{ji} = 0$ is flat, we shall always assume that

$$P_{ji} \neq 0. \quad (1.8)$$

2. Some necessary conditions

By definition, a P_N^* is a non-flat projectively flat A_N with recurrent curvature and therefore it is characterized by the conditions (1.4), (1.7) and (1.8). To find all the P_N^* 's, we first derive a set of necessary and sufficient conditions and then determine the functions ϕ_j in (1.6) which satisfy these conditions.

From (1.4) we obtain $\nabla_i R_{ji} = r_i R_{ji}$, and consequently

$$\nabla_k P_{ji} = r_k P_{ji} \quad (r_k \neq 0). \quad (2.1)$$

It is easy to see that conversely (1.4) is a consequence of (2.1) and (1.7)₁.

On account of (2.1), equation (1.7)₂ becomes

$$r_k P_{ji} = r_j P_{ki} ,$$

which is equivalent to

$$P_{ji} = r_j p_i , \quad (2.2)$$

where p_i is some non-zero vector (since $P_{ji} \neq 0$).

Using (2.2) in (1.7)₁, we see that the curvature tensor of a P_N^* has the form

$$R_{kji}{}^h = - (A_k^h r_j - A_j^h r_k) p_i + (r_k p_j - r_j p_k) A_i^h . \quad (2.3)$$

On account of (2.2), equation (2.1) becomes

$$(\nabla_k r_j) p_i + r_j (\nabla_k p_i) = r_k r_j p_i , \quad (2.4)$$

from which it follows that

$$\begin{aligned} \nabla_j r_i &= s_j r_i , \\ \nabla_j p_i &= t_j p_i , \end{aligned} \quad (2.5)$$

for some vectors s_j, t_j . These equations show that r_i and p_i are of the form

$$\begin{aligned} r_i &= \lambda \xi_i \quad (\xi_i = \partial_i \xi \neq 0) , \\ p_i &= \mu \eta_i \quad (\eta_i = \partial_i \eta \neq 0) , \end{aligned} \quad (2.6)$$

for some scalars ξ, η, λ, μ .

Substitution of (2.6) in (2.3) and (2.5) gives

$$R_{kji}{}^h = \psi [(A_k^h \xi_j - A_j^h \xi_k) \eta_i - (\xi_k \eta_j - \xi_j \eta_k) A_i^h] \quad (2.7)$$

$$\begin{aligned} \nabla_j \xi_i &= \alpha_j \xi_i , \\ \nabla_j \eta_i &= \beta_j \eta_i , \end{aligned} \quad (2.8)$$

where ψ, α, β are some scalars.

We now proceed to consider the integrability conditions of the differential equations (2.8). For any covariant vector u_i , we have the Ricci identity

$$\nabla_k \nabla_j u_i - \nabla_j \nabla_k u_i = - R_{kji}{}^h u_h . \quad (2.9)$$

Using (2.7) in (2.9) and putting $u_i = \xi_i$, we get, since $\xi_i \neq 0$,

$$(\xi_k \alpha_j - \xi_j \alpha_k) = - \psi (\xi_k \eta_j - \xi_j \eta_k) \quad (\alpha_j = \partial_j \alpha) ,$$

i.e.

$$\xi_k (\alpha_j + \psi \eta_j) = \xi_j (\alpha_k + \psi \eta_k)$$

which is equivalent to

$$\alpha_j + \psi \eta_j = \varrho \xi_j \quad (2.10)_1$$

for some scalar ϱ . Similarly, using (2.7) in (2.9) and putting $u_i = \eta_i$, we get

$$\beta_j - 2\psi\xi_j = \sigma\eta_j \quad (\beta_j = \partial_i\beta) \quad (2.10)_2$$

for some scalar σ .

Equations (2.10) tell us that

The scalars $\alpha, \beta, \psi, \varrho, \sigma$ can be expressed as functions of ξ, η alone. (2.11)

On account of (2.11), equations (2.10) can be re-written as

$$\left. \begin{aligned} (\alpha_\xi - \varrho)\xi_j + (\alpha_\eta + \psi)\eta_j &= 0 \\ (\beta_\xi - 2\psi)\xi_j + (\beta_\eta - \sigma)\eta_j &= 0 \end{aligned} \right\} \quad (2.12)$$

where $\alpha_\xi = \frac{\partial\alpha}{\partial\xi}$, $\alpha_\eta = \frac{\partial\alpha}{\partial\eta}$, etc. Two cases arise according as whether or not η is functionally dependent on ξ , i.e. whether or not η_i and ξ_i are proportional. Now, by (2.2) and (2.6), η_i and ξ_i are proportional or not according as the tensor P_{ji} (or the RICCI tensor) is symmetric or not. We shall consider these two cases separately in § 3 and § 4 respectively.

3. P_N^* with symmetric RICCI tensor

In this case, $\eta = \eta(\xi)$ so that (2.12) reduce to

$$\begin{aligned} (\alpha_\xi - \varrho) + (\alpha_\eta + \psi)\eta' &= 0, \\ (\beta_\xi - 2\psi) + (\beta_\eta - \sigma)\eta' &= 0, \end{aligned}$$

where $\eta' = d\eta/d\xi$. Since these equations just determine the unspecified functions ϱ, σ of ξ which appear in (2.10), the *integrability conditions* of (2.8) reduce to the mere fact that α, β and ψ are functions of ξ .

On account of $\eta = \eta(\xi)$, equations (2.7) and (2.8) become

$$\begin{aligned} R_{kji}{}^h &= \psi\eta'(A_k^h\xi_j - A_j^h\xi_k)\xi_i, \\ \nabla_j\xi_i &= \alpha\xi_j\xi_i, \\ \nabla_j\xi_i &= (\beta\eta' - \eta''/\eta')\xi_j\xi_i. \end{aligned}$$

Hence, we have the first part of the following

Theorem 3.1. *For a P_N^* with symmetric RICCI tensor, there exists a scalar ξ and functions θ and α of ξ such that*

$$R_{kji}{}^h = \theta(A_k^h\xi_j - A_j^h\xi_k)\xi_i, \quad (3.1)$$

$$\nabla_j\xi_i = \alpha\xi_j\xi_i, \quad (3.2)$$

where $\xi_i = \partial_i\xi$.

[The integrability condition of (3.2) is identically satisfied.]

Conversely, an A_N satisfying this condition is a P_N^* (or a symmetric P_N) with symmetric Ricci tensor.

Proof. We need only prove the last part of the theorem. It follows from (3.1) that

$$P_{ji} = -\theta \xi_j \xi_i, \quad (3.3)$$

and so P_{ji} is symmetric. On account of (3.3) and (3.1), condition (1.7)₁ is satisfied. Furthermore, a simple calculation involving (3.3) and (3.2) will show that $\nabla_k P_{ji} = r_k P_{ji}$, with

$$r_j = -(\theta_\xi/\theta + 2\alpha)\xi_j \quad \left(\theta_\xi = \frac{d\theta}{d\xi}\right). \quad (3.4)$$

It follows from this that (1.7)₂ is satisfied and that $\nabla_i R_{kji}^h = r_i R_{kji}^h$. The A_N is therefore a P_N^* or a symmetric P_N according as $\theta_\xi/\theta + 2\alpha \neq 0$ or $= 0$. Hence our theorem is completely proved.

In order to construct the Γ_{ji}^h of a P_N^* with symmetric Ricci tensor, we choose a coordinate system in which Γ_{ji}^h have the form

$$\Gamma_{ji}^h = \vartheta_j A_i^h + \vartheta_i A_j^h, \quad (3.5)$$

where ϑ_j are N functions of the coordinates. With respect to the Γ_{ji}^h given by (3.5), the covariant derivation of a covariant vector u_i is

$$\nabla_j u_i = \partial_j u_i - \vartheta_j u_i - \vartheta_i u_j, \quad (3.6)$$

and the curvature tensor has the components

$$R_{kji}^h = -A_k^h \vartheta_{ji} + A_j^h \vartheta_{ki} + (\vartheta_{kj} - \vartheta_{jk}) A_i^h. \quad (3.7)$$

where

$$\vartheta_{ji} = \partial_j \vartheta_i - \vartheta_j \vartheta_i. \quad (3.8)$$

Now identifying (3.7) with (3.1) and rewriting (3.2) by means of (3.6), we get

$$\begin{aligned} \partial_j \vartheta_i &= \vartheta_j \vartheta_i - \theta(\xi) \xi_j \xi_i, \\ \partial_j \xi_i &= \vartheta_j \xi_i + \vartheta_i \xi_j + \alpha(\xi) \xi_j \xi_i, \end{aligned} \quad (3.9)$$

where $\xi_i = \partial_i \xi$.

It is easy to verify that for any functions $\theta(\xi)$ and $\alpha(\xi)$ of ξ , the integrability conditions of the differential equations (3.9) in the $N+1$ unknown functions ϑ_i, ξ are satisfied, on account of the equations (3.9) themselves. Now the solution of a completely integrable system of differential equations contains a finite number of arbitrary constants, while the special form (3.5) for the connection of a P_N is preserved by affine transformations of coordinates

which depend on a finite number of constants. But neither of these would alter the fact that Γ_{ji}^h as given by (3.5) depends on two arbitrary functions of one variable, namely, the functions θ, α of ξ . Hence (cf. (3.4)) we have

Theorem 3.2. *If we take any functions θ, α ($\alpha \neq -\theta_\xi/2\theta$) of one variable ξ , and any solution φ_i, ξ of the completely integrable system (3.9), then the connection $\Gamma_{ji}^h = \varphi_j A_i^h + \varphi_i A_j^h$ defines a P_N^* with symmetric Ricci tensor; and any P_N^* with symmetric Ricci tensor can be constructed in the way. Thus the most general P_N^* with symmetric Ricci tensor depends on 2 arbitrary functions of one variable, and the actual construction of the connection of such P_N^* depends on the solution of a completely integrable system of differential equations.*

4. P_N^* with non-symmetric Ricci tensors

Let us now return to (2.12). Since η_i, ξ_i are not proportional, (2.12) are equivalent to

$$\begin{aligned} \alpha_\xi &= \varrho, & \beta_\eta &= \sigma, \\ \alpha_\eta + \psi &= 0, & \beta_\xi - 2\psi &= 0. \end{aligned}$$

The first two equations merely determine the unspecified scalars ϱ, σ , which appear in (2.10). Therefore,

$$\alpha_\eta + \psi = 0, \quad \beta_\xi - 2\psi = 0$$

are the integrability conditions of equations (2.8).

From this and (2.7), (2.8) and (2.11), we have the first part of the following

Theorem 4.1. *For a P_N^* with non-symmetric Ricci tensor, there exist functionally independent scalars ξ and η and functions ψ, α, β of ξ and η such that*

$$R_{kji}^h = \psi[(A_k^h \xi_j - A_j^h \xi_k)\eta_i - (\xi_k \eta_j - \xi_j \eta_k)A_i^h] \quad (4.1)$$

$$\left. \begin{aligned} \nabla_j \xi_i &= \alpha \xi_j \xi_i, \\ \nabla_j \eta_i &= \beta \eta_j \eta_i, \end{aligned} \right\} \quad (4.2)$$

where $\xi_i = \partial_i \xi, \eta_i = \partial_i \eta$.

[The integrability conditions of (4.2) are

$$\alpha_\eta + \psi = 0, \quad \beta_\xi - 2\psi = 0.] \quad (4.3)$$

Conversely, any A_N which satisfies the above condition is a P_N^* with non-symmetric Ricci tensor.

Proof. We need only prove the last part of this theorem. It follows from (4.1) that

$$P_{ji} = -\psi \xi_j \eta_i, \quad (4.4)$$

and so P_{ji} is not symmetric. Now (4.4), together with (4.1) and (4.2) show that condition (1.7) for a P_N is satisfied. Furthermore, a simple calculation involving (4.4) and (4.2) will show that $\nabla_k P_{ji} = r_k P_{ji}$, and hence $\nabla_i R_{kji}^h = r_i R_{kji}^h$, with

$$r_k = \left(\frac{\psi_\xi}{\psi} + \alpha \right) \xi_k + \left(\frac{\psi_\eta}{\psi} + \beta \right) \eta_k .$$

This vector r_k is not zero. In fact, since η_k, ξ_k are not proportional, $r_k = 0$ would imply that

$$(\log \psi)_\xi + \alpha = 0, \quad (\log \psi)_\eta + \beta = 0 .$$

Differentiations of these give

$$\alpha_\eta = \beta_\xi ,$$

from which and (4.3) it follows that $\psi = 0$. But this gives a flat space. Hence our theorem is completely proved.

In order to construct the Γ_{ji}^h of a P_N^* with non-symmetric RICCI tensor, we proceed as in the case of symmetric RICCI tensor, and have (3.5), (3.6), (3.7) and (3.8). Identifying (3.7) with (4.1) and rewriting (4.2) by means of (3.6), we get

$$\left. \begin{aligned} \partial_j \varnothing_i &= \varnothing_j \varnothing_i + \psi(\xi, \eta) \xi_j \eta_i, \\ \partial_j \xi_i &= \varnothing_j \xi_i + \varnothing_i \xi_j + \alpha(\xi, \eta) \xi_j \eta_i, \\ \partial_j \eta_i &= \varnothing_j \eta_i + \varnothing_i \eta_j + \beta(\xi, \eta) \xi_j \eta_i, \end{aligned} \right\} \quad (4.4)$$

where $\xi_i = \partial_i \xi$, $\eta_i = \partial_i \eta$. From Theorem 4.1 it follows that an A_N with connection (3.5) is a P_N^* with non-symmetric RICCI tensor if the functions \varnothing_j together with two functionally independent scalars ξ, η and some functions ψ, α, β of ξ, η , satisfy the differential equations (4.4). We know already (from Theorem 4.1) that the integrability conditions of $(4.4)_{2,3}$ are (4.3). It is easy to verify that on account of (4.4), the integrability condition of $(4.4)_1$ reduces to

$$(\psi_\eta + \psi\beta)(\xi_k \eta_j - \xi_j \eta_k) = 0 .$$

Since $\eta \neq \eta(\xi)$, this is equivalent to

$$\psi_\eta + \psi\beta = 0 .$$

Hence combining this with (4.3), we obtain the following complete set of integrability conditions of (4.4):

$$\alpha_\eta = -\psi, \quad \beta_\xi = 2\psi, \quad \psi_\eta + \psi\beta = 0 .$$

Rewrite these as

$$\frac{\partial \alpha}{\partial \eta} = -\psi, \quad \frac{\partial \beta}{\partial \xi} = 2\psi, \quad \frac{\partial \psi}{\partial \eta} = -\psi\beta. \quad (4.5)$$

From the last two equations, we obtain

$$\frac{\partial^2 \beta}{\partial \xi \partial \eta} = 2 \frac{\partial \psi}{\partial \eta} = -2\psi\beta = -\beta \frac{\partial \beta}{\partial \xi},$$

i.e.
$$\frac{\partial^2 \beta}{\partial \xi \partial \eta} + \frac{1}{2} \frac{\partial \beta^2}{\partial \xi} = 0.$$

Integration of this with respect to ξ gives

$$\frac{\partial \beta}{\partial \eta} + \frac{1}{2} \beta^2 = f(\eta), \quad (4.6)_1$$

where $f(\eta)$ is an arbitrary function of η . If β is any function of ξ, η satisfying (4.6)₁, then in virtue of (4.5), ψ and α can be expressed in terms of β as follows:

$$\psi = \frac{1}{2} \frac{\partial \beta}{\partial \xi}, \quad (4.6)_2$$

$$\alpha = -\frac{1}{2} \int \frac{\partial \beta}{\partial \xi} d\eta + h(\xi). \quad (4.6)_3$$

where $h(\xi)$ is an arbitrary function of ξ . Equations (4.5) are equivalent to equations (4.6).

Since equation (4.6)₁ is of the RICCATI type, the most general function $\beta(\xi, \eta)$ satisfying it is of the form

$$\beta(\xi, \eta) = \frac{\beta_1(\eta) + g(\xi)\beta_2(\eta)}{\beta_3(\eta) + g(\xi)\beta_4(\eta)}, \quad (4.7)$$

where $\beta_1(\eta), \dots, \beta_4(\eta)$ are certain functions of η and $g(\xi)$ is an arbitrary function of ξ .

Hence, with an observation similar to that immediately before Theorem 3.2, we have

Theorem 4.2. *With any functions $f(\eta)$ and $h(\xi)$ of one variable η and ξ respectively, any solution $\beta(\xi, \eta)$ [cf. (4.7)] of equation (4.6)₁, and the functions $\psi(\xi, \eta)$, $\alpha(\xi, \eta)$ given by (4.6)₂, (4.6)₃, the system of differential equations (4.4) is completely integrable. If ϕ_i, ξ, η are any solution of the completely*

integrable system (4.4), then the connection $\Gamma_{ji}^h = \phi_j A_i^h + \phi_i A_j^h$ defines a P_N^* with non-symmetric Ricci tensor; and any such P_N^* can be constructed in this way. Thus the most general P_N^* with non-symmetric Ricci tensor depends on 3 arbitrary functions of one variable, and the actual construction of the connection of such P_N^* depends on the solution of a differential equation of the RICCATI type and on the solution of a completely integrable system of differential equations.

5. Projectively flat symmetric N -space

We now consider the A_N 's characterized by

$$R_{kji}^h = -A_k^h P_{ji} + A_j^h P_{ki} + (P_{kj} - P_{jk}) A_i^h \quad (5.1)$$

$$\nabla_i R_{kji}^h = 0. \quad (5.2)$$

On account of (5.1), the condition (5.2) is equivalent to

$$\nabla_k P_{ji} = 0. \quad (5.2')$$

Substituting (5.2') into the Ricci identity

$$\nabla_i \nabla_k P_{ji} - \nabla_k \nabla_i P_{ji} = -R_{ikj}^a P_{ai} - R_{ikl}^a P_{ja}$$

and taking account of (5.1), we obtain

$$\begin{aligned} 0 &= P_{li} P_{kj} - P_{ki} P_{lj} - (P_{lk} - P_{kl}) P_{ji} \\ &\quad + P_{jl} P_{ki} - P_{jk} P_{li} - (P_{lk} - P_{kl}) P_{ji}, \end{aligned}$$

$$\text{i.e.} \quad P_{li} (P_{kj} - P_{jk}) + P_{ki} (P_{jl} - P_{lj}) - 2(P_{lk} - P_{kl}) P_{ji} = 0.$$

Putting $l = j$, $k = i$, we get

$$P_{ji} (P_{ij} - P_{ji}) - 2(P_{ji} - P_{ij}) P_{ji} = 0,$$

$$\text{i.e.} \quad 3(P_{ji} - P_{ij}) P_{ji} = 0 \quad (i, j \text{ not summed}).$$

This gives

$$P_{ji} = P_{ij}, \quad (5.3)$$

and consequently, (5.1) becomes

$$R_{kji}^h = -A_k^h P_{ji} + A_j^h P_{ki}. \quad (5.4)$$

Since, conversely, (5.1) and (5.2) are easy consequences of (5.4), (5.3) and (5.2'), we have

Theorem 5.1. *An A_N is a symmetric P_N , iff*

$$R_{kji}{}^h = -A_k^h P_{ji} + A_j^h P_{ki},$$

$$P_{ji} = P_{ij}, \quad \nabla_k P_{ji} = 0.$$

The properties stated in Theorem 5.1 characterize a well-known type of projectively flat space; in particular, if P_{ji} is of rank n , it is RIEMANNIAN and of constant curvature (cf. EISENHART [1], p. 97 and p. 166). We can easily prove that in contrast, a P_N with recurrent curvature, i.e. a P_N^* , is never a RIEMANNIAN space.

Obviously, a P_N^* with a non-symmetric RICCI tensor cannot be RIEMANNIAN. If a P_N^* with a symmetric RICCI tensor is RIEMANNIAN, let g_{ji} be a fundamental tensor whose CHRISTOFFEL symbol $\left\{ \begin{smallmatrix} h \\ ji \end{smallmatrix} \right\}$ is equal to Γ_{ij}^h . Then the covariant components

$$R_{kji}{}^h = R_{kji}{}^a g_{ha}$$

of the curvature tensor satisfy the well-known equation

$$R_{kji}{}^h = R_{ihk}{}^j,$$

which on account of (3.1) reduces to

$$g_{kh} \xi_j \xi_i = g_{ij} \xi_h \xi_k.$$

But this contradicts the fact that g_{ji} is of rank $N > 1$. Therefore, no P_N^* is RIEMANNIAN.

*University of Hong Kong
Tokyo Institute of Technology*

REFERENCES

- [1] L. P. EISENHART, *Non-RIEMANNIAN Geometry*. New York (1927).
- [2] H. S. RUSE, *A classification of K^* -spaces*. Proc. London Math. Soc. (2), 53 (1951), 212–229.
- [3] H. S. WALKER, *On RUSE's space of recurrent curvature*. Proc. London Math. Soc. (2), 52 (1950), 36–64.
- [4] YUNG-CHOW WONG, *A class of Non-RIEMANNIAN K^* -space*. Proc. London Math. Soc. (3), 3 (1953).

(Received August 11, 1960)