

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 35 (1961)

Artikel: Fibrations and Cocategory.
Autor: Ganea, Tudor
DOI: <https://doi.org/10.5169/seals-27326>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Fibrations and Cocategory¹⁾

by TUDOR GANEA, Bucharest

1. Introduction and results

In a previous paper [3], I have defined a based homotopy type invariant, the cocategory, which appears as dual to LUSTERNIK-SCHNIRELMANN category within the framework of the ECKMANN-HILTON [1] duality in homotopy theory. The cocategory, $\text{cocat } X$, of an arbitrary topological space X with base-point is the (possibly infinite) strictly positive integer given by the following inductive

Definition 1.1. *$\text{cocat } X = 1$ if and only if X is contractible; $\text{cocat } X \leq n + 1$ whenever there exists a fibration $Q \rightarrow Y \rightarrow B$ such that the fibre Q dominates X and $\text{cocat } Y \leq n$. If the phrase $\text{cocat } X \leq n$ is false for all $n \geq 1$, we put $\text{cocat } X = \infty$.*

Evidently, $\text{cocat } X = n$ will mean that $\text{cocat } X \leq n$ is true but $\text{cocat } X \leq n - 1$ is false. It is assumed that all homotopies involved in 1.1 keep base-points fixed; the precise sense of the word fibration is explained in the next section.

In the ECKMANN-HILTON setting, fibrations are dual to cofibrations, i. e. sequences $Q \leftarrow Y \leftarrow B$ in which the first arrow results by pinching to a point the image of B in Y , while the second has the lowering homotopy property [5; p. 14]. It follows that cocategory is indeed dual to category since, provided X has the based homotopy type of a connected CW-complex, $\text{cat } X$ is equal to the invariant obtained by reversing the arrows and replacing in 1.1 the words *fibration* and *fibre* by *cofibration* and *cofibre* respectively [3; Th. 1.9]. The following relations between $\text{cocat } X$ and standard homotopy invariants of X have been established:

1.2. *If X is a 1-connected CW-complex with only n non-trivial Postnikov invariants k^{q+2} , then $\text{cocat } X \leq n + 2$; in particular, if the 1-connected CW-complex X has only n non-trivial homotopy groups, then $\text{cocat } X \leq n + 1$ [3; Th. 2.10 and Cor. 2.11].*

1.3. *If X has a non-trivial n -fold WHITEHEAD product, then $\text{cocat } X \geq n + 1$ [3; Cor. 2.13].*

¹⁾ Presented at the International Colloquium on Differential Geometry and Topology, Zürich, June 1960.

Also, for every $n \geq 1$ there exists a connected CW -complex X such that $\text{cocat } X = n$ [3; Remark 2.16].

The purpose of this paper is to present two further results.

Theorem 1.4. *If X is a $(p-1)$ -connected ($p \geq 2$) CW -complex such that $\pi_q(X) = 0$ for $q \geq r+1$, then $\text{cocat } X \leq [(r-1)/(p-1)] + 1$.*

Here $[a/b]$ stands for the largest integer $\leq a/b$. Theorem 1.4 dualizes a previous result by D. P. GROSSMAN [4] according to which $\text{cat } X \leq [r/p] + 1$ if X is a $(p-1)$ -connected complex of dimension $\leq r$. In fact, GROSSMAN's result may be restated for a 1-connected complex X such that $H^q(X; G) \neq 0$ only if $p \leq q \leq r$. The slight difference between the numerical estimation given in 1.4 and that of the GROSSMAN theorem agrees with the relations

$$\dim[\alpha, \beta] = \dim \alpha + \dim \beta - 1 \quad \text{and} \quad \dim u \cup v = \dim u + \dim v,$$

involving WHITEHEAD and cup products which are dual to each other. The proof of 1.4 is based on the extension, given in the next section, of a well known result concerning fibrations with a $K(\pi, n)$ as fibre.

Our next result refers to the cocategory of $(n-1)$ -connective spaces (X, n) over X , and to that of spaces (n, X) obtained by attaching cells to X so as to kill its homotopy groups in dimensions $\geq n$. When X is a CW -complex we shall assume, as we may, that both (X, n) and (n, X) have the based homotopy type of CW -complexes, and state

Theorem 1.5. *Let X be a connected CW -complex. Then, for all $n \geq 1$, $\text{cocat } (X, n) \leq \text{cocat } X$ and $\text{cocat } (n, X) \leq \text{cocat } X$.*

For $n = 2$ we have the

Corollary 1.6. *The simply connected covering space \tilde{X} of a connected CW -complex X satisfies $\text{cocat } \tilde{X} \leq \text{cocat } X$.*

2. A lemma on induced fibrations

All spaces, maps, and homotopies hereafter are assumed to possess, preserve, or keep fixed a base-point, generally denoted by $*$. A sequence

$\mathcal{F}: Q \xrightarrow{\eta} Y \xrightarrow{\beta} B$ of spaces and maps is a fibration with fibre $Q = \beta^{-1}(*)$ and inclusion map η if for any space E , any homotopy $h_t: E \rightarrow B$ and any map $k: E \rightarrow Y$ satisfying $\beta \circ k = h_0$, there is a homotopy $H_t: E \rightarrow Y$ such that $H_0 = k$ and $\beta \circ H_t = h_t$. We do not require that β be onto. The space of paths in B emanating from the base-point is denoted by EB , the loop-space by ΩB . Consider the fibration \mathcal{F} above, and let $\Phi: C \rightarrow B$

be a map; the sequence $Q \xrightarrow{\zeta} Z \xrightarrow{\gamma} C$ in which

$$Z = \{(c, y) \mid \Phi(c) = \beta(y)\} \subset C \times Y, \quad *_Z = (*_C, *_Y),$$

$$\zeta(q) = (*_C, \eta(q)) \quad \text{and} \quad \gamma(c, y) = c,$$

is the familiar fibration induced by \mathcal{F} via Φ . Suppose the rows in the diagram

$$\begin{array}{ccccc} Q_1 & \xrightarrow{\eta_1} & Y_1 & \xrightarrow{\beta_1} & B_1 \\ \downarrow h & & \downarrow g & & \downarrow f \\ Q_2 & \xrightarrow{\eta_2} & Y_2 & \xrightarrow{\beta_2} & B_2 \end{array}$$

are fibrations. The first is algebraically equivalent to the second if there are singular homotopy equivalences f, g, h such that each square commutes. A map $f: X \rightarrow Y$ is a singular homotopy equivalence if $f_q: \pi_q(X, x) \rightarrow \pi_q(Y, f(x))$ is isomorphic for all $q \geq 0$ and all $x \in X$; if X and Y are 0-connected, it suffices to take $x = *$. We shall often use the geometric realization $|S(X)|$ of the singular complex of an arbitrary space X and the canonical map $j_X: |S(X)| \rightarrow X$ which induces homotopy and homology isomorphisms in all dimensions [7].

Lemma 2.1. *Let $\mathcal{F}: Q \xrightarrow{\eta} Y \xrightarrow{\beta} B$ be a fibration with Y and B both having the based homotopy type of a CW-complex. Suppose that B is $(m-1)$ -connected and that $\pi_q(Q) \neq 0$ only if $n \leq q \leq n+m-2$, where $m \geq 2$ and $n \geq 1$. Suppose further that there exists a singular homotopy equivalence $\theta: Q \rightarrow \Omega W$, where W is a 1-connected space. Then, there exists a map $\Phi: B \rightarrow W$ such that \mathcal{F} is algebraically equivalent to the fibration induced by $\mathcal{G}: \Omega W \xrightarrow{i} EW \xrightarrow{p} W$ via Φ .*

Proof. Introduce the diagram

$$\begin{array}{ccccccc} Q & \xleftarrow{j_Q} & |S(Q)| & \xrightarrow{f} & \Omega W & & \\ \eta \downarrow & & \downarrow |\eta| & & \downarrow i & & \\ Y & \xleftarrow{j_Y} & |S(Y)| & \xrightarrow{g} & EW & & \\ \beta \downarrow & & \downarrow |\beta| & \searrow \varphi & & & \\ B & \xleftarrow{j_B} & |S(B)| & \xleftarrow{k} & C(\gamma) & \xrightarrow{h} & W \\ & & \downarrow \varrho & & \downarrow \varepsilon & \searrow \psi & \\ & & |S(Y)| / |S(Q)| & & & & \end{array} \tag{1}$$

The maps $|\eta|$ and $|\beta|$ are induced by η and β respectively, and the two squares on the left commute; $|S(Q)|$ is a subcomplex of $|S(Y)|$ and $|\eta|$ is the inclusion map. Since $|S(*)| = *$ and $\beta \circ \eta(Q) = *$, we have

$$|\beta| \circ |\eta| (|S(Q)|) = * . \quad (2)$$

Let $|S(Y)| / |S(Q)|$ and φ result by pinching the subset $|S(Q)|$ of $|S(Y)|$ to a point, which will serve as base-point in $|S(Y)| / |S(Q)|$. It follows from (2) that $\gamma = |\beta| \circ \varphi^{-1}$ is single-valued, and hence continuous. According to [11; § 8], the space $|S(Y)| / |S(Q)|$ may be given a CW-structure and γ is easily seen to be cellular; therefore, its reduced mapping cylinder $C(\gamma)$ is a CW-complex in which $|S(Y)| / |S(Q)|$ and $|S(B)|$ are embedded as subcomplexes by means of the maps ε and κ respectively. The standard retraction ϱ of $C(\gamma)$ onto $|S(B)|$ satisfies the relation $\varrho \circ \varepsilon = \gamma$. Let $f = \theta \circ j_Q$. Since the CW-pair $(|S(Y)|, |S(Q)|)$ has the homotopy extension property and since EW is contractible, there exists a map g such that

$$g \circ |\eta| = i \circ f . \quad (3)$$

Finally, since $p \circ i(\Omega W) = *$, (3) implies that $\psi = p \circ g \circ \varphi^{-1}$ is single-valued, and hence continuous.

We now prove that there is a map h such that

$$h \circ \varepsilon = \psi .$$

Since ε is an inclusion map, this amounts to extending ψ over the complex $C(\gamma)$. The diagram

$$\begin{array}{ccccc} H_q(Y, Q) & \xleftarrow{j_*} & H_q(|S(Y)|, |S(Q)|) & \xrightarrow{\varphi_*} & H_q(|S(Y)| / |S(Q)|, *) \\ \downarrow \beta_* & & \downarrow |\beta|_* & \searrow \gamma_* & \downarrow \varepsilon_* \\ H_q(B, *) & \xleftarrow{(j_B)_*} & H_q(|S(B)|, *) & \xleftarrow{\varrho_*} & H_q(C(\gamma), *) \end{array} \quad (4)$$

in which j_* is induced by the map of pairs defined by j_Y , is obviously commutative. Consideration of the upper left square in (1) and the five lemma show j_* isomorphic in all dimensions; excision in the CW-pair $(|S(Y)|, |S(Q)|)$ implies that so is also φ_* , while $(j_B)_*$ and ϱ_* are standard isomorphisms. Since $\pi_q(Q) = 0$ for $q < n$ and $\pi_q(B) = 0$ for $q < m$, a well known result by SERRE [9; p. 469] implies that β_* , whence $|\beta|_*$ and ε_* , are monomorphic for $q \leq n + m - 1$ and epimorphic for

$q \leq n + m$. Passing to cohomology, the universal coefficient theorem yields

$$H^{q+1}(C(\gamma), |S(Y)| / |S(Q)|; G) = 0 \quad \text{for all } q \leq n + m - 1 \quad (5)$$

and all coefficient groups G . Since $\pi_q(\Omega W) \approx \pi_q(Q)$, we also have

$$\pi_q(W) = 0 \quad \text{for all } q \geq n + m. \quad (6)$$

From (5) and (6) we finally obtain

$$H^{q+1}(C(\gamma), |S(Y)| / |S(Q)|; \pi_q(W)) = 0 \quad \text{for all } q \geq 0,$$

and a standard obstruction argument now yields the desired map h .

Since Y and B have the based homotopy type of a CW -complex, there exist homotopy inverses e_Y and e_B of j_Y and j_B respectively. Select a homotopy

$$b_t: |S(B)| \rightarrow |S(B)| \quad \text{with } b_0 = \text{id}, \quad b_1 = e_B \circ j_B, \quad b_t(*) = *.$$

Notice next that there is a homotopy

$$k_t: |S(Y)| / |S(Q)| \rightarrow C(\gamma) \quad \text{with } k_0 = \varepsilon, \quad k_1 = k \circ \gamma, \quad k_t(*) = *.$$

Define a homotopy $H_t: |S(Y)| \rightarrow W$ by

$$\begin{aligned} H_t(y) &= h \circ k_{2t} \circ \varphi(y) && \text{if } 0 \leq t \leq \frac{1}{2}, \\ &= h \circ k \circ b_{2t-1} \circ |\beta|(y) && \text{if } \frac{1}{2} \leq t \leq 1. \end{aligned}$$

Taking (2) into account, we obtain

$$H_t(|S(Q)|) = * = p \circ g(|S(Q)|) \quad \text{and} \quad H_0(y) = p \circ g(y).$$

Therefore, by [6], there is a map $g_1: |S(Y)| \rightarrow \Omega W$ such that

$$g_1 \circ |\eta| = g \circ |\eta| \quad \text{and} \quad h \circ k \circ e_B \circ j_B \circ |\beta| = p \circ g_1 \quad (7)$$

Let $\Phi = h \circ k \circ e_B$ and let $\mathcal{H}: \Omega W \xrightarrow{\zeta} Z \xrightarrow{\lambda} B$ be the fibration induced by \mathcal{G} via Φ . According to (7), a map $d: |S(Y)| \rightarrow Z$, satisfying

$$d \circ |\eta| = \zeta \circ f \quad \text{and} \quad j_B \circ |\beta| = \lambda \circ d, \quad (8)$$

is defined by setting $d(y) = (j_B \circ |\beta|(y), g_1(y))$. In the sequence

$$\pi_q(|S(Y)|, |S(Q)|) \xrightarrow{j_q} \pi_q(Y, Q) \xrightarrow{\beta_q} \pi_q(B, *) \xleftarrow{\lambda_q} \pi_q(Z, \Omega W),$$

where j_q is induced by the map of pairs defined by j_Y , the first arrow, as in (4), is isomorphic for all $q \geq 1$; since \mathcal{F} and \mathcal{H} are fibrations, so are also β_q and λ_q . Therefore, (8) and commutativity on the left in (1) imply that the map of pairs defined by d induces isomorphisms

$$\pi_q(|S(Y)|, |S(Q)|) \approx \pi_q(Z, \Omega W)$$

in all dimensions. Since $f = \theta \circ j_Q$ is a singular homotopy equivalence, the first of the relations (8) and the five lemma now imply that d also is a singular homotopy equivalence. As easily seen, the map $\lambda \circ d \circ e_Y: Y \rightarrow B$ is homotopic to β . Since \mathcal{H} is a fibration, the covering homotopy theorem yields a map $D: Y \rightarrow Z$, which is homotopic to $d \circ e_Y$, and satisfies $\lambda \circ D = \beta$; let $F: Q \rightarrow \Omega W$ be the map defined by D . Like $d \circ e_Y$, D is a singular homotopy equivalence; the five lemma implies that so is also F , and the required algebraic equivalence is now provided by the maps id_B , D , F .

Remark 2.2. Letting $m = 2$ in 2.1 we recover the well known result concerning fibrations with a $K(\pi, n)$ as fibre (see for instance [5; Th. 7.1, p. 43]).

Lemma 2.1 has a dual concerning induced cofibrations.

3. Proof of Theorem 1.4

It is well known that any $(n - 1)$ -connected CW -complex of dimension $< 2n$ has the homotopy type of a suspension. Dually, we have

Lemma 3.1. *Let X be an arbitrary space and let $n \geq 2$. If X is $(n - 1)$ -connected and $\pi_q(X) = 0$ for $q \geq 2n - 1$, then there exists a 1-connected space W and a singular homotopy equivalence $X \rightarrow \Omega W$.*

Proof. The space W is obtained by attaching cells to the reduced suspension ΣX so as to kill its homotopy groups in dimensions $\geq 2n$. Let $\sigma: \Sigma X \rightarrow W$ denote the inclusion map and consider the sequence

$$X \xrightarrow{e} \Omega \Sigma X \xrightarrow{\Omega \sigma} \Omega W,$$

in which e is the natural embedding. Evidently, $\Omega \sigma$ induces isomorphisms of homotopy groups in dimensions $\leq 2n - 2$; by the FREUDENTHAL theorem (see for instance [8; p. 05]), so does also e . Finally, for $q \geq 2n - 1$ we have $\pi_q(X) = \pi_q(\Omega W) = 0$.

Proof of 1.4. The result is obvious if $1 \leq r \leq p - 1$ since X then is contractible. Suppose $r \geq p$ and let X be an arbitrary $(p - 1)$ -connected

CW-complex such that $\pi_q(X) = 0$ for $q \geq r + 1$. Let the *CW*-complex B result by attaching cells to X so as to kill its homotopy groups in dimensions $\geq r - p + 2$. Replace the inclusion map $X \rightarrow B$ by a homotopy equivalent fibre map to obtain a fibration $\mathcal{F}: Q \rightarrow Y \rightarrow B$ such that

$$Y \text{ has the homotopy type of } X, \quad (9)$$

$$\pi_q(B) \neq 0 \quad \text{only if} \quad p \leq q \leq r - p + 1, \quad (10)$$

$$\pi_q(Q) \neq 0 \quad \text{only if} \quad \max(p, r - p + 2) \leq q \leq r. \quad (11)$$

Since $r - p + 1 < r$, we may assume as an induction hypothesis that (10) implies

$$\text{cocat } B \leq [(r - p)/(p - 1)] + 1. \quad (12)$$

It follows from (10), (11), 3.1, and Lemma 2.1 that there is a 1-connected space W and a map $\Phi: B \rightarrow W$ such that \mathcal{F} is algebraically equivalent to the fibration $\Omega W \rightarrow Z \rightarrow B$ induced by $\Omega W \rightarrow EW \rightarrow W$ via Φ . Therefore, Y has the homotopy type of the singular polytope of Z . By (9), [3; Prop. 2.8 and 2.9], and (12) we finally obtain

$$\text{cocat } X = \text{cocat } |S(Z)| \leq \text{cocat } Z \leq \text{cocat } B + 1 \leq [(r - 1)/(p - 1)] + 1.$$

4. Proof of Theorem 1.5

For any 0-connected space X and any $n \geq 1$ there is a space (X, n) and a map $p: (X, n) \rightarrow X$ such that $\pi_q(X, n) = 0$ if $q < n$ and $p_q: \pi_q(X, n) \approx \pi_q(X)$ if $q \geq n$. Similarly, there is a space (n, X) and a map $j: X \rightarrow (n, X)$ such that $\pi_q(n, X) = 0$ if $q \geq n$ and $j_q: \pi_q(n, X) \approx \pi_q(n, X)$ if $q < n$. When X has the homotopy type of a *CW*-complex, we shall assume, as we may, that the same holds for both (X, n) and (n, X) ; their homotopy type is then uniquely determined by that of X and n .

Proof of 1.5. If $\text{cocat } X = 1$, then X is contractible and so are both (X, n) and (n, X) . Suppose 1.5 is true for any connected *CW*-complex of cocategory $\leq m$ and suppose $\text{cocat } X = m + 1$. Let $Q \xrightarrow{\eta} Y \xrightarrow{\beta} B$ be a fibration such that Q dominates X and $\text{cocat } Y = m$.

Let $R \xrightarrow{\varphi} Z \xrightarrow{\psi} |S(B)|_0$ denote the fibration obtained by replacing the map $|\beta|_0: |S(Y)|_0 \rightarrow |S(B)|_0$ by a homotopy equivalent fibre map; the subscript 0 indicates restriction to the path-component of the base-point.

As easily seen, there is a map $r: R \rightarrow Q$ which, by the five lemma, induces isomorphisms

$$r_q: \pi_q(R) \approx \pi_q(Q) \quad \text{for all } q \geq 1. \quad (13)$$

Let C be a connected covering space of $|S(B)|_0$ such that $\pi_1(C)$ maps isomorphically onto the subgroup $\psi_1\pi_1(Z)$ of $\pi_1(|S(B)|_0)$ under the projection $f: C \rightarrow |S(B)|_0$. Since Z has the homotopy type of a connected CW -complex, the monodromy principle yields a map $g: Z \rightarrow C$ such that $f \circ g = \psi$. Let $T \xrightarrow{\varepsilon} W \xrightarrow{\gamma} C$ be the fibration obtained by replacing g by a homotopy equivalent fibre map. As above, there is a map $t: T \rightarrow R$ which, since f_1 is monomorphic, induces isomorphisms

$$t_q: \pi_q(T) \approx \pi_q(R) \quad \text{for all } q \geq 1. \quad (14)$$

For the same reason and since $f_1\pi_1(C) = \psi_1\pi_1(Z)$, the homomorphism $\gamma_1: \pi_1(W) \rightarrow \pi_1(C)$ is onto. Therefore, in $\pi_n(C)$ the subgroup

$$\gamma_n\pi_n(W) \text{ is closed under the operations of } \pi_1(C). \quad (15)$$

Introduce the diagram

$$\begin{array}{ccccc} U & \xrightarrow{\sigma} & (W, n) & \xrightarrow{\lambda} & D \\ \downarrow h & & \downarrow p & & \downarrow d \\ T & \xrightarrow{\varepsilon} & W & \xrightarrow{\gamma} & C \\ \downarrow k & & \downarrow j & & \downarrow e \\ V & \xrightarrow{\tau} & (n, W) & \xrightarrow{\mu} & E \end{array}$$

The space E and the inclusion map e are obtained by attaching cells to C in such a way that

$$e_q: \pi_q(C) \approx \pi_q(E) \quad \text{if } q < n,$$

the sequence

$$\pi_n(W) \xrightarrow{\gamma_n} \pi_n(C) \xrightarrow{e_n} \pi_n(E) \rightarrow 0$$

be exact, and $\pi_q(E) = 0$ if $q > n$; according to [10; Th. 2.10.1] this is possible in view of (15). The space D and the map d are selected so that $\pi_q(D) = 0$ if $q < n$,

$$d_n: \pi_n(D) \approx \gamma_n\pi_n(W) \quad \text{and} \quad d_q: \pi_q(D) \approx \pi_q(C)$$

if $q > n$. Since W has the homotopy type of a connected CW -complex, so have, by assumption, (W, n) and (n, W) , and standard arguments now

yield maps λ and μ for which the two squares on the right are homotopy commutative. Without altering the homotopy types of (W, n) and (n, W) , we may assume that λ and μ are fibre maps with U and V as fibres; the inclusion maps are denoted by σ and τ . Next, by means of the covering homotopy theorem, we may readjust the maps p and j within their own homotopy classes so as to obtain totally commutative squares on the right. Suppose this is so and let h and k be the maps defined by p and j respectively. Passing to homotopy groups, application of the five lemma, in the form given in [2; p. 16], to the resulting ladder yields

$$\pi_q(U) = 0 \quad \text{if} \quad q < n, \quad h_q : \pi_q(U) \approx \pi_q(T) \quad \text{if} \quad q \geq n, \quad (16)$$

$$\pi_q(V) = 0 \quad \text{if} \quad q \geq n, \quad k_q : \pi_q(T) \approx \pi_q(V) \quad \text{if} \quad q < n. \quad (17)$$

Since X is a connected CW -complex which is dominated by Q , X is also dominated by $|S(Q)|_0$. Since γ_1 is onto and W is 0-connected, T is 0-connected and, by (13) and (14), $|S(Q)|_0$ has the homotopy type of $|S(T)|$. It follows from (16) that $(|S(T)|, n)$ has the homotopy type of $|S(U)|$, while (17) implies that $(n, |S(T)|)$ has the homotopy type of $|S(V)|$. Since (X, n) and (n, X) have the homotopy type of CW -complexes, it follows easily that (X, n) is dominated by $|S(U)|$, and (n, X) by $|S(V)|$.

Since W , like Z , has the homotopy type of $|S(Y)|_0$, and since the component of the base-point in a CW -complex is a retract of the complex, by [3; Prop. 2.8] we have

$$\text{cocat } W = \text{cocat } |S(Y)|_0 \leq \text{cocat } |S(Y)| \leq \text{cocat } Y = m.$$

Since W has the homotopy type of a connected CW -complex, the induction hypothesis now implies that $\text{cocat } (W, n) \leq m$ and $\text{cocat } (n, W) \leq m$. By [3; Prop. 2.8] and 1.1 we obtain

$$\text{cocat } |S(U)| \leq \text{cocat } U \leq m + 1, \quad \text{cocat } |S(V)| \leq \text{cocat } V \leq m + 1,$$

and this clearly implies the desired result.

Appendix

(Added in proof)

The inductive arguments used in the proof of 1.4 are easily seen to yield the following more general result:

Let X be a $(p - 1)$ -connected CW -complex, $p \geq 2$. If the set of all integers q for which $\pi_q(X) \neq 0$ is contained in the union of k closed linear intervals, each of length $p - 2$, then $\text{cocat } X \leq k + 1$.

We allow the linear intervals to be degenerate, i.e. to have length 0. The second part of 1.2 now follows as the set $\{q_1, \dots, q_n\}$ is contained in the intervals $[q_j, q_j]$, $j = 1, \dots, n$; Theorem 1.4 follows upon noticing that the integers between p and r are all contained in the intervals

$$[j(p-1) + 1, j(p-1) + p-1], j = 1, \dots, \left[\frac{r-1}{p-1} \right].$$

Also, the author wishes to acknowledge that a result equivalent to Lemma 2.1 above has been obtained independently and with a different proof by P. J. HILTON as Theorem 3 in his paper "Excision and principal fibrations", Comment. Math. Helv. 35 (1961).

REFERENCES

- [1] B. ECKMANN et P. J. HILTON, *Groupes d'homotopie et dualité*. C. R. Acad. Sci. Paris, 246 (1958), 2444–2447, 2555–2558, 2991–2993.
- [2] S. EILENBERG and N. E. STEENROD, *Foundations of algebraic topology*, 1952.
- [3] T. GANEÀ, LUSTERNIK-SCHNIRELMANN *category and cocategory*. Proc. London Math. Soc. 10 (1960), 623–639.
- [4] D. P. GROSSMAN, *An estimation of the category of LUSTERNIK-SCHNIRELMANN*. C. R. (Doklady) Acad. Sci. URSS (N.S.) 54 (1946), 109–112.
- [5] P. J. HILTON, *Homotopy theory and duality*. Lecture notes, Cornell University (1959).
- [6] I. M. JAMES and J. H. C. WHITEHEAD, *Note on fibre spaces*. Proc. London Math. Soc. 4 (1954), 129–137.
- [7] J. MILNOR, *The geometric realization of a semi-simplicial complex*. Annals of Math. 65 (1957), 357–362.
- [8] J. C. MOORE, *Le théorème de FREUDENTHAL, la suite exacte de JAMES et l'invariant de HOPF généralisé*. Séminaire H. Cartan, E. N. S. (1955).
- [9] J. P. SERRE, *Homologie singulière des espaces fibrés*. Ann. of Math. 54 (1951), 425–505.
- [10] G. W. WHITEHEAD, *Homotopy theory*. Lecture notes, Massachusetts Institute of Technology (1953).
- [11] J. H. C. WHITEHEAD, *Combinatorial homotopy I*. Bull. Amer. Math. Soc. 55 (1949), 213–245.

Institute of Mathematics, R.P.R. Academy, Bucharest

(Received June 4, 1960)