

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 36 (1961-1962)

Artikel: On Combinatorial Submanifolds of Differentiable Manifolds.
Autor: Hirsch, Morris W.
DOI: <https://doi.org/10.5169/seals-515620>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On Combinatorial Submanifolds of Differentiable Manifolds¹⁾

by MORRIS W. HIRSCH, California (USA)

§ 1. Introduction

The purpose of this work is to prove the following results relating combinatorial and differentiable manifolds.

(A) *A combinatorial submanifold V of a differentiable manifold M of the same dimension possesses a compatible differentiable structure.*

(B) *Every compact and contractible combinatorial manifold V possesses a compatible differentiable structure.²⁾*

(A differentiable structure on a combinatorial manifold M is called *compatible* if M has a rectilinear subdivision, each simplex of which is differentiably imbedded.)

A. M. GLEASON has announced (unpublished) that a contractible *unbounded* combinatorial manifold has a compatible differentiable structure. Theorem (B) follows easily from this and Theorem (A). The proof of (B) given here is derived from JOHN STALLINGS' proof [11] of the generalized POINCARÉ conjecture.

(C) *The sequence*

$$\dots \xrightarrow{d} \Gamma^n \xrightarrow{j} \Theta^n \xrightarrow{k} \Lambda^n \xrightarrow{d} \Gamma^{n-1} \xrightarrow{\dots} \dots \quad (1)$$

is well defined and exact.

This result was announced in [3]. Here Γ^n is the group of differentiable structures on S^n compatible with the usual combinatorial structure; Θ^n is the group of differentiable homotopy n -spheres modulo J -equivalence, and Λ^n the combinatorial analogue of Ω^n . Using powerful intrinsic methods, STEPHEN SMALE has shown [9, 10] that for $n \geq 8$ or $n = 5$, the map $j: \Gamma^n \rightarrow \Theta^n$ is an isomorphism, and using (B) (but not (C)) that $\Lambda^n = 0$, for $n \geq 8$ or $n = 6$.

In [8] SMALE proves that $\Gamma^3 = 0$, (this was proved independently by J. MUNKRES, and J. H. C. WHITEHEAD.) The fact that every combinatorial 3-manifold possess a unique (up to a diffeomorphism) compatible differentiable structure [6] implies that $k: \Theta^3 \rightarrow \Lambda^3$ is an isomorphism. Thus only the

¹⁾ Presented at the International Colloquium on Differential Geometry and Topology, Zürich, June 1960.

²⁾ *Added in proof;* The hypothesis of compactness is unnecessary.

subsequences $0 \rightarrow \Gamma^7 \rightarrow \Theta^7 \rightarrow \Lambda^7 \rightarrow \Gamma^6 \rightarrow 0$ and $0 \rightarrow \Lambda^5 \rightarrow \Gamma^4 \rightarrow \Theta^4 \rightarrow \Lambda^4 \rightarrow 0$ remain. In proving (C), SMALE's results are not used.

§ 2. Proof of (A)

In order to prove (A) it suffices to establish the stronger result 2.5 below.

If K is a subcomplex of complex N , the n th *simplicial neighborhood* of K is the union of the closed simplexes of the n 'th barycentric subdivision of N that meet K .

Lemma 2.1. *Let K be the boundary of a combinatorial manifold M . The second simplicial neighborhood A of K is combinatorially equivalent to $K \times I$, where I is the unit interval.*

Proof. This is a well known result. It follows e. g. from theorems 22 and 23 of [14], which state that any two "regular neighborhoods" of M (in the sense of [14]) in the same manifold are combinatorially equivalent and that A is a regular neighborhood of K in M . If K is identified with $K \times 0$, then $M' = M \cup K \times I$ is a manifold, and in M' both A and $K \times I$ are regular neighborhoods of K .

Now let V be a bounded combinatorial n -manifold imbedded as a subcomplex of an unbounded combinatorial n -manifold M . Assume M has a metric $d(x, y)$.

Lemma 2.2. *Let U be a neighborhood of V in M , and ϵ a positive continuous function on M . There is a semi-linear homeomorphism $h: M \rightarrow M$ with the following properties:*

- a) $h(V)$ is the second simplicial neighborhood of V in a subdivision of M ;
- b) $h(V) \subset U$;
- c) $h(x) = x$ if $x \in M - U$;
- d) $d(x, h(x)) < \epsilon(x)$ for all $x \in M$.

Proof. By 2.1, the boundary of K of V has a neighborhood combinatorially equivalent to $K \times I$ in V , and another in $cl(M - V)$. The union B_0 of these two neighborhoods is again equivalent to $K \times I$. Moreover, we can take B_0 to be the second simplicial neighborhood of K in a subdivision of M ; if this subdivision is sufficiently fine, the second simplicial neighborhood B of B_0 will be in U . It will be clear that if the subdivision is sufficiently fine, d) will be satisfied. There is a combinatorial equivalence $u: B \rightarrow K \times I$ such that $u(B_0) = K \times [\frac{1}{4}, \frac{3}{4}]$ and $u(K) = K \times \frac{1}{2}$. We may assume that $h(B_0 \cap V) = K \times [0, \frac{1}{2}]$. Let $f: I \rightarrow I$ be a semi-linear homeomorphism

such that $f(x) = x$ for x in a neighborhood of 0 and 1, and $f(\frac{1}{2}) = 3/4$. Define $g: K \times I \rightarrow K \times I$ by $g(x, t) = (x, f(t))$. Now define $h: M \rightarrow M$ by $h(x) = \begin{cases} x & \text{if } x \in M - B \\ u^{-1}gu(x) & \text{if } x \in B. \end{cases}$

Then h is the desired homeomorphism.

Now let M be a differentiable manifold. A combinatorial manifold A which is a subcomplex of a smooth triangulation of M is called a *combinatorial submanifold of M* . A vector field Φ on A in M is *transverse* if it is transverse to A in every coordinate system, in the sense of [13]. The following lemma is well known.

Lemma 2.3. *Let A be the boundary of the second simplicial neighborhood B of a subcomplex K of M . Then there is a transverse field on A .*

Proof. Each simplex ϱ of the first simplicial neighborhood B' of K is the join $\sigma^* \tau$ of unique simplices $\sigma \subset K$ and $\tau \subset M - K$. Each closed simplex α of A lies in such a join $\sigma^* \tau$, disjoint from σ and τ , and each $x \in \alpha$ lies on a unique line p^*q with $p \in \sigma$, $q \in \tau$. It is easily seen that the unit tangent $\Phi(x)$ to p^*q , directed from p to q , is transverse to α at x , and that Φ is continuous. Thus Φ is a transverse field on A .

Lemma 2.4. *Let M be an unbounded differentiable n -manifold, and $V \subset M$ a combinatorial submanifold, also of dimension n . Let U be a neighborhood of the boundary A of V , d a metric on M , and ϵ a positive continuous function on M . There is a homeomorphism $h: M \rightarrow M$ such that*

- a) h is a diffeomorphism on each closed simplex of a subdivision of M ;
- b) $h(A)$ has a transverse field;
- c) $h(x) = x$ if $x \in M - U$;
- d) $d(x, h(x)) < \epsilon(x)$ for all $x \in M$.

Proof. Apply 2.2 and 2.3.

Let V be a combinatorial submanifold of a differentiable unbounded n -manifold M . Assume either

1. V has dimension n ; or
2. V has dimension $n - 1$, is unbounded, and admits a transverse field.

Let U be a neighborhood of bdV and ϵ a positive continuous function on M .

Theorem 2.5. *There is a homeomorphism $h: M \rightarrow M$ such that:*

- a) $h(V)$ is a differentiable submanifold of M , combinatorially equivalent to V ;

b) M has a smooth triangulation in which V is a subcomplex, every closed simplex of which is mapped diffeomorphically by h ;
 c) and d) as in 2.4.

Proof. Case 1) follows from 2.1 and case 2); Thus we assume 2).

By standard approximation methods, it may be assume that there is a differentiable non-zero vector field Φ on a neighborhood W of V contained in U , such that $\Phi|V$ is transverse field. A generalization of the CAIRNS-WHITEHEAD theory of transverse fields [1, 13] shows that there is a submanifold C of dimension $n - 1$ differentiably imbedded in an arbitrary neighborhood of V , such that $\Phi|C$ is transverse. (The CAIRNS-WHITEHEAD theory applies to a q -dimensional submanifold of EUCLIDEAN $(q + p)$ -space endowed with a transverse p -plane field. The present case follows, e. g., by imbedding M in R^{n+k} , and assigning to each point $x \in V$ the $k + 1$ plane generated by $\Phi(x)$ and the k -plane normal to M at x . Alternatively, the methods of [13] simplify considerably in the special case where the submanifold has codimension 1, if transverse lines are replaced by the integral curves of a transverse vector field.³⁾ We can assume that each integral curve of Φ meets V in a unique point and C in a unique point. This establishes a map $f: V \rightarrow C$ which is a diffeomorphism on each closed simplex of V . Thus C is combinatorially equivalent to V . Let A be the region bounded by V and C which is fibered by the integral curves of Φ . Define $G: V \times I \rightarrow A$ by $G(x, 0) = x$, $G(x, 1) = f(x)$ and $G(x, t)$ is the point dividing the length of the integral curve joining x and $f(x)$ in the ratio $t/(1-t)$, for $0 < t < 1$. Then if Δ is a closed simplex of V , $G| \Delta \times I$ is a diffeomorphism. Hence $G: K \times I \rightarrow M$ is a non-degenerate C^∞ subcomplex of M in the sense of [15]. By [15, p. 822, *addendum*] this triangulation of A can be extended to a smooth triangulation of M , after possible subdivision. An easy application of 2.1 (cf. proof of 2.2) establishes the desired extension $h: M \rightarrow M$ of $f: V \rightarrow C$. This completes the proof.

Remark. It can be shown that if a neighborhood in V of a closed subset $X \subset V$ is a differentiable submanifold of M , h can be chosen so that for some neighborhood Y of X in M , $h(x) = x$ if $x \in Y$.

The following theorem was announced by S. S. Cairns [16].

Theorem 2.6 (Cairns). *If M is a combinatorial manifold and if for some p , $M \times R^p$ has a compatible differentiable structure, then so has M .*

Proof. By induction on p . The case $p = 0$ is trivial. Let $F^p \subset R^p$ be a closed half-space. First assume M is unbounded. If $p > 0$ and if $M \times R^p$

³⁾ Cf. [18].

has a compatible differentiable structure, then so has $M \times F^p$, by (A). So therefore does its boundary $M \times R^{p-1}$, completing the induction. The case where M is bounded follows easily now from (A).

§ 3. Proof of (B)

Let V be an n -dimensional compact combinatorial manifold which is contractible. Let \tilde{V} be the double of V , obtained by identifying two disjoint copies of V along their boundary. Then \tilde{V} is a closed combinatorial n -manifold of the same homotopy type as S^n , and V is a submanifold. J. STALLINGS proves in [11] that if x is any point of \tilde{V} , then $\tilde{V} - x$ is combinatorially equivalent to EUCLIDEAN n -space, provided $n \geq 7$, and states that E. C. ZEEMAN has extended the result to the case $n \geq 5$. (If $n \leq 4$, theorem (B) is a consequence of well known results of CAIRNS [1, 2].) Thus we can assume that V is a submanifold of the differentiable manifold R^n . (Alternatively, $\tilde{V} - x$ is an unbounded contractible manifold, and one can apply GLEASON's theorem that $\tilde{V} - x$ has a compatible differentiable structure). Theorem (B) now follows from (A). Actually, this proves the following stronger result.

Theorem 3.1. *A compact, contractible, combinatorial n -manifold is combinatorially equivalent to a differentiable submanifold of R^n .*

GLEASON's theorem is proved by showing that an *unbounded* combinatorial contractible n -manifold can be *immersed* in R^n . This follows from the mere existence of a compatible differentiable structure by observing that such a structure is necessarily parallelizable, and applying a theorem of [4].⁴⁾

A plausible conjecture along these lines is that *any combinatorial manifold all of whose cohomology groups vanish has a compatible differentiable structure*. J. MUNKRES [7] has proved this except for compatibility.

§ 4. Proof of (C)

We must show that the sequence

$$\xrightarrow{d} \Gamma^n \xrightarrow{j} \Theta^n \xrightarrow{k} A^n \xrightarrow{d} \Gamma^{n-1} \longrightarrow \quad (1)$$

is well defined and exact.

The group Θ^n is defined as follows. An element of Θ^n is an equivalence class $[M]$ of oriented, closed differentiable manifolds M which have the homo-

⁴⁾ GLEASON's theorem follows from 2.6 and [17], in which it is proved that if M^n is a contractible combinatorial unbounded manifold, then $M^n \times R^p$ is combinatorially equivalent to R^{n+p} for some p .

topy type of the n -sphere S^n , under the relation of J -equivalence. Two oriented differentiable manifolds M_0, M_1 are J -equivalent if there is an oriented differentiable manifold N whose boundary is (diffeomorphic to) the disjoint union of M_1 and $-M_0$ (where $-M_0$ means M_0 with the opposite orientation), and such that both M_0 and M_1 are deformation retracts of N . Addition in Θ^n is defined by $[A] + [B] = [A \# B]$ where $A \# B$ is the *connected sum* of A and B . This is defined by removing the interior of an n -ball from each of A and B and joining the two boundary $(n-1)$ -spheres by an orientation reversing diffeomorphism which is extendable to the whole n -ball, and then smoothing the resulting corner. It can be shown that the diffeomorphism class of $A \# B$ is independent of the choices made, and the J -equivalence class $[A \# B]$ is independent of the representatives of $[A]$ and $[B]$ that are chosen. See [5] for details. Define $-[A] = [-A]$, and Θ^n becomes an abelian group, with $[S^n]$ as identity element.

Using combinatorial instead of differentiable manifolds, J -equivalence and connected sum are analogously defined, and Λ^n is the group of J -equivalence classes $\langle M \rangle$ of oriented combinatorial closed n -manifolds M which are homotopy spheres.

The elements of Γ^n are (diffeomorphism classes of) oriented differentiable manifolds which are combinatorially equivalent to the boundary of an $(n+1)$ -simplex. Addition is defined using $\#$, and the inverse of $M \in \Gamma^n$ is $-M$. Using these definitions, Γ^n is an abelian group, with S^n for 0, although this is not obvious. It is a consequence of the following result.

Theorem 4.1. (MUNKRES-THOM). *There is at most one compatible differentiable structure on a contractible combinatorial n -manifold, up to a diffeomorphism.*

Proof. See [6, 12]

The only difficulty about proving that Γ^n is a group is showing

$$M \# (-M) = S^n.$$

We can obtain $M \# (-M)$ by removing the interior E of an n -ball from M , and taking the boundary V of $(M - E) \times I$ and smoothing the corner. But $(M - E) \times I$ is then a differentiable manifold combinatorially equivalent to $\Delta^n \times I$, where Δ^n is an n -simplex, and by 4.1 its boundary $V = M \# (-M)$ is diffeomorphic to S^n , which is the zero element of Γ^n .

The map $k: \Theta^n \rightarrow \Lambda^n$ is defined as follows. If M is a differentiable manifold, let kM be the corresponding combinatorial manifold, i. e., kM is a simplicial complex L such that there is a homeomorphism $t: L \rightarrow M$ which is a smooth triangulation of M . Such an L exists and is unique up to combinatorial equivalence [15].

Now define $k: \Theta^n \rightarrow \Lambda^n$ by $k[M] = \langle kM \rangle$. It is obvious that k preserves sums and J -equivalence, so k is a well defined homomorphism.

The map $j: \Gamma^n \rightarrow \Theta^n$ is defined by $jM = [M]$.

To define $d: \Lambda^n \rightarrow \Gamma^{n-1}$, let M represent an element of Λ^n , and let E be the interior of an n -simplex of M . Then $M - E$ is contractible, and by (B) possesses a compatible differentiable structure, which is unique by 4.1, up to diffeomorphism. The combinatorial structure of $M - E$ is independent of E , and dM is defined to be the boundary of $M - E$. We shall see shortly that if A and B are J -equivalent, then $dA = dB$.

Lemma 4.2. a) $d(M \# N) = dM + dN$

b) $d(-M) = -dM$.

Proof. Let C and D be closed n -simplices in M and N respectively. In forming $M \# N$, remove the interiors of n -simplices disjoint from C and D . Now M -int C and N -int D have unique compatible differentiable structures by (B) and 4.1, and then $(M \# N) - (\text{int } C \cup \text{int } D) = (M\text{-int } C) \# (N\text{-int } D)$ has a compatible differentiable structure. Now join C to D in $M \# N$ by a simple differentiable arc, meeting C and D only at its end points. A tubular neighborhood Q of this arc can be chosen so that $C \cup D \cup Q$ is a combinatorial n -cell in $M \# N$. Then M -int $(C \cup D \cup Q)$ has $d(M \# N)$ for its boundary (after smoothing). On the other hand, this boundary is diffeomorphic to $\partial(M\text{-int } C) \# \partial(N\text{-int } D) = d(M) \# d(N)$, which proves a). The proof of b) is obvious.

Lemma 4.3. *Let M be a closed oriented combinatorial homotopy n -sphere. If M has a compatible differentiable structure, $dM = 0$.*

Proof. If E is the interior of an n -simplex Δ of M , and B the interior of an n -ball differentiably imbedded in E , then $M - E$ and $M - B$ are combinatorially equivalent. Assuming M has a compatible differentiable structure, take Δ to be a simplex of a smooth triangulation of M . Then $M - B$ is a differentiable submanifold of M and hence $\partial(M - B) = \partial B = S^{n-1} = 0 \in \Gamma^{n-1}$. By 4.1, compatible differentiable structures on $M - E$ and $M - B$ are diffeomorphic; hence $\partial(M - E) = d(M) = 0$.

Corollary 4.4. *If M bounds a contractible manifold, $d(M) = 0$.*

Proof. By (A), M has a compatible differentiable structure and 4.3 applies.

Theorem 4.5. *$d: \Lambda^n \rightarrow \Gamma^{n-1}$ defined by $d \langle M \rangle = dM$ is a well defined homomorphism.*

Proof. We must show first that if $\langle M \rangle = \langle N \rangle$, then $dM = dN$. If M is J -equivalent to N , then $M \# (-N)$ is J -equivalent to $\partial \Delta^{n+1}$. Since

$\partial\Delta^{n+1}$ bounds Δ^{n+1} , $M \# (-N)$ bounds a contractible manifold. By 4.4 $d(M \# (-N)) = O$, and by 4.2, $d(M \# (-N)) = d(M) - d(N)$. Thus $d(M) - d(N) = O$, so $d: \Delta^n \rightarrow \Gamma^{n-1}$ is well defined, and 4.4 proves d to be a homomorphism.

Now we prove that the sequence (1) is exact. We leave the proof that $jd = kj = dk = O$ to the reader as an exercise; the last equality, for example, follows from 4.3.

Let M be an element of Γ^n such that $j(M) = O$. This means M bounds a contractible differentiable manifold V . Since $\partial V = M$ is combinatorially equivalent to $\partial\Delta^{n+1}$, $V \cup \Delta^{n+1}$ is a combinatorial homotopy sphere and hence represents an element λ of Δ^{n+1} . It is obvious that $d(\lambda) = \partial(V \text{-int } \Delta^{n+1}) = M$. This establishes the exactness of the sequence jd .

Let $\langle M \rangle \in \Delta^n$ be such that $d\langle M \rangle = O$. This means for some n -simplex Δ in M , M -int Δ has a compatible differentiable structure making $\partial(M \text{-int } \Delta)$ diffeomorphic to S^{n-1} . Choosing such a diffeomorphism, attach the n -ball D^n to M -int Δ to obtain a differentiable manifold N which is combinatorially equivalent to M . Thus $k[M] = \langle N \rangle$ and dk is exact.

Finally let M represent an element of Θ^n annihilated by k . This means M is J -equivalent to $\partial\Delta^{n+1}$ in the combinatorial sense. Let V be a combinatorial $(n+1)$ -manifold realizing this J -equivalence. Let T be a «tube» joining M to $\partial\Delta^{n+1}$ in V , i.e., T is equivalent to $I \times \Delta^n$ with $0 \times \Delta^n \subset M$ and $1 \times \Delta^n \subset \partial\Delta^{n+1}$, and no other points of T in ∂V . (Such a T can easily be constructed by first putting a compatible differentiable structure on the contractible manifold $V \cup \Delta^{n+1}$.) Then V -int T is contractible if T is “unknotted”, which is always true if $n+1 > 3$ and which is the case for $n+1 = 3$ provided T is chosen properly. (In fact, $\Theta^2 = O$, so this case is unnecessary.) Thus V -int T has a compatible differentiable structure, by (B). By 2.5, we can assume that M -int $(0 \times \Delta^n)$ is a differentiable submanifold A of the boundary of V -int T . The closure of the complement of A is combinatorially equivalent to $\partial\Delta^{n+1}$, and hence is diffeomorphic to D^n , while A is combinatorially equivalent, and hence diffeomorphic to $M - E$, where E is the interior of an n -ball, \overline{E} differentiably imbedded in M . Thus there is a diffeomorphism $f: \partial(M - E) \rightarrow S^{n-1}$ such that $\partial(V \text{-int } T)$ is diffeomorphic to $(M - E) \cup, D^n$. Let P be the manifold $\overline{E} \cup, D^n$. Then P is an element of Γ^n , and $\partial(V \text{-int } T)$ is the same as $M \# (-P)$. Since V -int T is contractible, $[M - P] = O$, and so $[M] = [P] = jP$. This establishes the exactness of the sequence (1).

REFERENCES

- [1] S. CAIRNS, *Homeomorphisms between topological manifolds and analytic manifolds*. Ann. of Math., **41** (1940), 796–808.
- [2] S. CAIRNS, *Introduction of a RIEMANNian geometry on a triangulable 4-manifold*. Ann. of Math. **45** (1944), 218–219.
- [3] M. HIRSCH, *An exact sequence in differential topology*. Bull. Amer. Math. Soc., **66** (1960), 322–323.
- [4] M. HIRSCH, *On imbedding differentiable manifolds in EUCLIDEAN space*. Ann. of Math., **73** (1960).
- [5] J. MILNOR, *Differentiable manifolds which are homotopy spheres*. Princeton University 1959 (mimeographed).
- [6] J. MUNKRES, *Obstructions to the smoothing of piecewise-differentiable homeomorphisms*. Ann. of Math., **72** (1960), 521–554.
- [7] J. MUNKRES, *Obstructions to imposing differentiable structures*. To appear.
- [8] S. SMALE, *Diffeomorphisms of the 2-sphere*. Proc. Amer. Math. Soc., **10** (1959), 621–626.
- [9] S. SMALE, *Generalized POINCARÉ's conjecture in dimensions greater than 4*. Ann. of Math., to appear.
- [10] S. SMALE, *Differentiable and combinatorial structures on manifolds*. Ann. of Math., to appear.
- [11] J. STALLINGS, *Polyhedral homotopy spheres*. Bull. Amer. Math. Soc., **66** (1960), 485–488.
- [12] R. THOM, *Des variétés triangulées aux variétés différentiables*. Proc. Int. Cong. Math., 1958, Cambridge Univ. Press, (1960), 248–255.
- [13] J. H. C. WHITEHEAD, *Manifolds with transverse fields in EUCLIDEAN space*. Ann. of Math., **73** (1960), 154–212.
- [14] J. H. C. WHITEHEAD, *Simplicial spaces, nuclei and m-groups*. Proc. Lond. Math. Soc., **45** (1939), 243–327.
- [15] J. H. C. WHITEHEAD, *On C^1 complexes*. Ann. of Math. **41** (1940), 809–824.
- [16] S. CAIRNS, *The manifold smoothing problem*, Bull. Amer. Math. Soc. **67** (1961), 237–238.
- [17] D. McMILLAN, *Cartesian products of contractible manifolds*, Bull. Amer. Math. Soc. **67** (1961), 510–514.
- [18] D. MONTGOMERY and H. SAMELSON, *Examples for differentiable group actions on spheres* Proc. Nat. Acad. Sci. U.S.A. **47** (1961), 1202–1205.

(Received March 27, 1961)