

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 37 (1962-1963)

Artikel: On a characterization of quasiconformal mappings.
Autor: Reich, Edgar
DOI: <https://doi.org/10.5169/seals-28605>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On a characterization of quasiconformal mappings¹

by EDGAR REICH

1. Introduction

Let $M(Q)$ denote the module of the quadrilateral Q (a JORDAN domain with four distinguished boundary points and two non-adjacent distinguished sides), A sense-preserving homeomorphism f of a region Ω onto the image region $f(\Omega)$ is said to be a quasiconformal mapping with maximal dilatation $K(\Omega) = K_r(\Omega)$ if

$$\sup_{\bar{Q} \subset \Omega} \frac{M(f(Q))}{M(Q)} = K(\Omega) < \infty. \quad (1.1)$$

($f(Q)$ denotes the quadrilateral with preimage Q .)

Instead of considering the effect of f on quadrilaterals it is natural to study the effect of f on ring domains (doubly connected regions). The modulus $\mu(R)$ of a ring R is defined as $(2\pi)^{-1} \log(r_2/r_1)$, where $\{r_1 < |w| < r_2\}$ is an annulus conformally equivalent to R . It is well known [1] that if (1.1) holds then

$$\frac{\mu(f(R))}{\mu(R)} \leq K(\Omega), \quad \bar{R} \subset \Omega. \quad (1.2)$$

In fact, the following has been proved by GEHRING and VÄISÄLÄ ([2], Theorem 3):

$$\sup_{\bar{R} \subset \Omega} \frac{\mu(f(R))}{\mu(R)} = K(\Omega). \quad (1.3)$$

Thus quasiconformal mappings f with maximal dilatation $K(\Omega)$ may be characterized by (1.3) instead of (1.1).

In the proof of (1.3) in [2] essential use is made of the deep and rather difficult "analytic" characterization of quasiconformal maps (See, for instance, [4], Chapter 4). In view of the significance of (1.3) it appears desirable to obtain a more direct and more elementary proof of (1.3), starting with the definition (1.1). It is the object of the present note to provide such a proof.

2. A class of ring domains

Let Q be a quadrilateral with distinguished sides α_1, α_2 . A ring R will be said to *link* Q if every closed curve γ , in R , separating the boundary com-

¹) This work was carried out with partial support from the Office of Naval Research under Contract Nonr 710 (16) with the University of Minnesota.

ponents of R , contains an arc lying in Q which joins α_1 and α_2 . (See Fig.1 where the heavy lines are the boundary components of R , and the dashed line represents Q .)




Fig. 1

We will first show that

$$\mu(R) \leq M(Q), \text{ if } R \text{ links } Q. \tag{2.1}$$

Let $\varrho^*(z), z \in Q$, be the extremal metric for Q ; that is,

$$M(Q) = \frac{A_{\varrho^*}(Q)}{L_{\varrho^*}^2(Q)}, \quad A_{\varrho^*}(Q) = \iint_Q \varrho^{*2} |dz|^2, \quad L_{\varrho^*}(Q) = \inf_{\gamma} \int_{\gamma} \varrho^* |dz|,$$

where $\{\gamma\}$ is the set of locally rectifiable curves in Q joining α_1, α_2 . On the other hand, (See, for instance, [3], Chapter 2),

$$\mu(R) = \inf_e \frac{A_e(R)}{L_e^2(R)} \tag{2.2}$$

where, this time, $\{\gamma\}$ is the set of closed curves in R separating the boundary components. For $z \in R$, let

$$\varrho(z) = \begin{cases} \varrho^*(z), & z \in R \cap Q \\ 0, & z \in R - Q \end{cases}.$$

Clearly,

$$A_{\varrho}(R) \leq A_{\varrho^*}(Q).$$

Since R links Q , $L_{\varrho}(R) \geq L_{\varrho^*}(Q)$. Hence, by (2.2),

$$\mu(R) \leq \frac{A_{\varrho^*}(Q)}{L_{\varrho^*}^2(Q)} = M(Q),$$

as was to be shown.

In general, to obtain a bound on the modulus of a ring domain R from below, one attempts to make use of the fact ([3], Chapter 2) that

$$\frac{1}{\mu(R)} = \inf_e \frac{A_e(R)}{L_e^2(R)} \tag{2.3}$$

where, now, $\{\gamma\}$ is the set of locally rectifiable curves in R joining the boun-

dary components. Fortunately, we shall require a bound on $\mu(R)$ from below, for rings linking Q , only in the very special case considered next.

Suppose the set of positive numbers, a, p, h, s , is given, and

$$2p < \min(a, 1), h < a - 2p, p < s < a - p - h, h < 1.$$

We consider the ring R' (Fig. 2) obtained as follows.




Fig. 2

The outer boundary component of R' consists of the union of (a) the boundary of the rectangle with vertices $0, 1, 1 + ia, ia$ (b) the vertical segment joining the points $\frac{1}{2} + ia, \frac{1}{2} + i(a - s)$ (c) the horizontal segment $I_1 = \{z \mid \Im z = a - s, p \leq \Re z \leq 1 - p\}$.

The inner boundary component of R' is the horizontal segment

$$I_2 = \{z \mid \Im z = a - s - h, p \leq \Re z \leq 1 - p\}.$$

Let T be the rectangle with horizontal sides I_1, I_2 , with the vertical sides (of length h) distinguished. The ring R' links T , and, therefore, by (2.1), $\mu(R') \leq h/(1 - 2p)$. We shall now show that

$$\mu(R') \geq \frac{h}{1 + ah^{\frac{1}{2}}}, \text{ if } h \leq p^4. \tag{2.4}$$

Let

$$\varrho(z) = \begin{cases} 1, & 0 < \Re z < 1, a - s - h < \Im z < a - s \\ h^{\frac{3}{4}}, & \text{elsewhere in } R'. \end{cases}$$

Let $\{\gamma\}$ be the family of curves in R' joining I_2 to the outer boundary component of R' . Clearly,

$$L_\varrho(R') = \min(h, h^{\frac{3}{4}}p), A_\varrho = h + h^{\frac{3}{2}}(a - h) \leq h + ah^{\frac{3}{2}}.$$

Hence, by (2.3),

$$\frac{1}{\mu(R')} \leq \frac{A_\varrho(R')}{L_\varrho^2(R')} \leq \frac{h + ah^{\frac{3}{2}}}{h^2} = \frac{1 + ah^{\frac{1}{2}}}{h}, \text{ if } h^{\frac{3}{4}}p \geq h.$$

This is equivalent to (2.4).

3. Proof of (1.3)

Let ϵ , $0 < \epsilon < \frac{1}{2}$, be given. In the notation of Section 1, there exists a quadrilateral $Q, \bar{Q} \subset \Omega$, with

$$\frac{M(f(Q))}{M(Q)} \geq (1 - \epsilon) K(\Omega).$$

Let the transformation Φ map $f(Q)$ conformally onto a rectangle V with vertices $0, 1, 1 + ia, ia$, with the vertical sides distinguished. Thus

$$a = M(V) = M(f(Q)).$$

Let V_p be the rectangle, interior to V and oriented as V , such that the distance between corresponding sides of V_p and V is p , $0 < p < \frac{1}{2} \min(a, 1)$. Let $Q_p = f^{-1} \Phi^{-1}(V_p)$. There exists [1] a number $\delta(\epsilon) > 0$ such that

$$\frac{M(V_p)}{M(Q_p)} = \frac{a - 2p}{(1 - 2p) M(Q_p)} \geq (1 - 2\epsilon) K(\Omega), \text{ if } p \leq \delta(\epsilon);$$

that is, Q_p has nearly the same module as Q if p is small.

We now divide V_p into n equal horizontal strips, σ_k , $k = 1, 2, \dots, n$. σ_k is an $h_n \times 1 - 2p$ rectangle, $h_n = (a - 2p) n^{-1}$, whose vertical sides are distinguished. The quadrilaterals

$$q_k = f^{-1} \Phi^{-1}(\sigma_k), \quad k = 1, 2, \dots$$

which lie in Ω , subdivide Q_p . According to the subadditivity property for modules [1],

$$\sum_{k=1}^n M(q_k) \leq M(Q_p) \leq \frac{a - 2p/1 - 2p}{(1 - 2\epsilon) K(\Omega)}.$$

Hence for some k , say $k = k_n$,

$$M(q_{k_n}) \leq \frac{a - 2p/1 - 2p}{(1 - 2\epsilon) n K(\Omega)} = \frac{h_n/1 - 2p}{(1 - 2\epsilon) K(\Omega)}.$$

If we define $N(p)$ as $N(p) = (a - 2p) p^{-4}$, then $n \geq N(p)$ implies $h_n \leq p^4$.

Let R'_{pn} be the ring linking σ_{k_n} formed as in Fig. 2, with the present h_n serving as the h of Fig. 2. Then, by (2.4),

$$\mu(R'_{pn}) \geq \frac{h_n}{1 + ah_n^{\frac{1}{2}}}, \text{ if } n \geq N(p).$$

Consider the ring $R_{pn} = f^{-1} \Phi^{-1}(R'_{pn})$, in Ω . Since R_{pn} links q_{k_n} , we have, by (2.1),

$$\mu(R_{pn}) \leq M(q_{kn}) \leq \frac{h_n/1 - 2p}{(1 - 2\varepsilon)K(\Omega)}, \text{ if } p \leq \delta(\varepsilon).$$

Now $\mu(f(R_{pn})) = \mu(R'_{pn})$. Therefore,

$$\frac{\mu(f(R_{pn}))}{\mu(R_{pn})} \geq \frac{(1 - 2\varepsilon)(1 - 2p)K(\Omega)}{1 + ah_n^{\frac{1}{2}}}, \text{ if } p \leq \delta(\varepsilon), n \geq N(p). \quad (3.1)$$

Once p is chosen, h_n can be made arbitrarily small by taking n sufficiently large. Thus (3.1) shows that there exist rings $R, \bar{R} \subset \Omega$, such that

$$\frac{\mu(f(R))}{\mu(R)} \geq (1 - 3\varepsilon)K(\Omega).$$

Together with (1.2) the above establishes (1.3).

University of Minnesota

December 1961

REFERENCES

- [1] L. V. AHLFORS, "On quasiconformal mappings" J. Analyse Math. 3 (1954), 1-58.
- [2] F. W. GEHRING and J. VÄISÄLÄ, "On the geometric definition for quasiconformal mappings", Comment Math. Helv. 36 (1961), 19-32.
- [3] J. A. JENKINS, *Univalent Functions and Conformal Mapping*, Springer Verlag, Berlin, 1958.
- [4] H. P. KÜNZI, *Quasikonforme Abbildungen*, Springer Verlag, Berlin, 1960.