

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 38 (1963-1964)

Artikel: On the Local Triviality of the Restriction Map for Embeddings.
Autor: Lima, Elon L.
DOI: <https://doi.org/10.5169/seals-29440>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the Local Triviality of the Restriction Map for Embeddings

by ELON L. LIMA¹⁾

Let V, M be C^∞ manifolds, V compact. A map $f: M \rightarrow M$ is said to have *compact support* if it agrees with the identity outside of a compact set. For $1 \leq r \leq \infty$, we consider the following spaces endowed with the C^r -topology: $\mathcal{E}^r(V, M) =$ all C^∞ embeddings of V in M ; $\mathcal{C}^r(M) =$ all C^∞ maps, with compact support, of M into M ; $\mathcal{D}^r(M) =$ all C^∞ diffeomorphisms, with compact support, of M onto M . We remark that $\mathcal{D}^r(M)$ is an open subset of $\mathcal{C}^r(M)$.

R. PALAIS proved [1] that if V is a submanifold of W then the restriction map $j: \mathcal{E}^r(W, M) \rightarrow \mathcal{E}^r(V, M)$ is a locally trivial fibration. Previously, R. THOM had observed [2] that j has the covering homotopy property for polyhedra. The local triviality of j follows easily from the theorem below, (see [1], or Remark 2), at the end of this note), which was also proved by J. CERF [3]. We present here a very simple proof of this theorem. For implications and applications, see the bibliography.

Theorem: *Given $f \in \mathcal{E}^r(V, M)$, there is a neighborhood U of f and a continuous map $\xi: U \rightarrow \mathcal{D}^r(M)$ such that $g = \xi(g) \circ f$ for every $g \in U$.*

Proof: We may assume that V is a submanifold of M , $f =$ inclusion, and M is embedded in some euclidean space R^k . Let $\pi': T' \rightarrow M$ be a tubular neighborhood of M in R^k and $\pi: T \rightarrow V$ a tubular neighborhood, of radius $\varepsilon > 0$, of V in R^k , with $T \subset T'$. Denote by $\frac{1}{2}T$ the tubular neighborhood of V with radius $\varepsilon/2$. Since the shortest line from a point in R^k to V is a normal segment, any line segment of length $< \varepsilon/2$ which intersects $\frac{1}{2}T$ lies entirely within T . Choose a neighborhood U' of f in $\mathcal{E}^r(V, M)$ so small that $|g(y) - y| < \varepsilon/2$ for all $g \in U'$ and all $y \in V$. Let $\lambda: R \rightarrow [0, 1]$ be a C^∞ function with $\lambda(t) = 1$ for $|t| \leq \varepsilon/4$ and $\lambda(t) = 0$ for $|t| \geq \varepsilon/2$. Define a map $\xi': U' \rightarrow \mathcal{C}^r(M)$ as follows. Given $g \in U'$, put $\xi'(g)(x) = x$, if $x \in M - T$, and $\xi'(g)(x) = \pi'\{x + \lambda(|x - \pi x|) \cdot [g(\pi x) - \pi x]\}$ if $x \in T$. One sees that ξ' is continuous and $\xi'(f)$ is the identity map of M , so $\xi'(f) \in \mathcal{D}^r(M)$. Since $\mathcal{D}^r(M)$ is open in $\mathcal{C}^r(M)$, a smaller neighborhood U of f can be chosen so that $\xi'(U) \subset \mathcal{D}^r(M)$. Put $\xi = \xi'|_U$.

Remarks: 1) Let $\mathcal{D}_0^r(M) \subset \mathcal{D}^r(M)$ be the subset of C^∞ diffeomorphisms, with compact support, that are diffeotopic to the identity. It is known that

¹⁾ The author holds a Guggenheim Fellowship. Work partially supported by NSF-G 21514.

$\mathcal{D}'_0(M)$ is open in $\mathcal{D}'(M)$. (This can be seen by a construction similar to, and simpler than, the above one.) So, if needed, U may be taken such that $\xi(U) \subset \mathcal{D}'_0(M)$.

2) Given $f \in \mathcal{D}'(V, M)$, take ξ and U as in the theorem, let $F = j^{-1}(f)$ and define a homeomorphism $\psi: F \times U \rightarrow j^{-1}(U)$ by $\psi(\bar{f}, g) = \xi(g) \circ \bar{f}$, for $\bar{f} \in F$, $g \in U$. This shows that j is a locally trivial fibration.

3) When $r = \infty$, $\mathcal{E}^\infty(V, M)$ and $\mathcal{D}^\infty(J)$ are C^∞ (infinite dimensional) manifolds, locally homeomorphic with FRÉCHET spaces. The reason why ξ is continuous is that, in the last analysis, it is obtained as a series of compositions of the variable map g with fixed C^∞ maps. Now, composition is a differentiable map in the C^∞ topology. (See [4], pages 182, 183.) So, by the same token, ξ is a C^∞ map when $r = \infty$. It follows from this and Remark 2) above that $j: \mathcal{E}^\infty(W, M) \rightarrow \mathcal{E}^\infty(V, M)$ would be a C^∞ fibration, in the sense that the local trivializing maps $\psi: F \times U \rightarrow j^{-1}(U)$ are C^∞ , provided one could show that F is a differentiable manifold.

Institute for Advanced Study, Princeton, N. J. and
I.M.P.A., Rio de Janeiro, Brazil

BIBLIOGRAPHY

- [1] R. PALAIS, *Local triviality of the restriction map for embeddings*. Comment. Math. Helv. 34 (1960) pp. 305–312.
- [2] R. THOM, *La classification des immersions (d'après Smale)*. Séminaire Bourbaki, 10e année: 1957/1958, exposé 157.
- [3] J. CERF, *Topologie de certains espaces de plongement*. Bull. Soc. Math. France 89 (1961) pp. 227–380.
- [4] J. DIEUDONNÉ, *Foundations of Modern Analysis*. Academic Press, New York, 1960.

(Received April 29, 1963)