

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 38 (1963-1964)

Artikel: On a Certain Property of Closed Hypersurfaces in an EINSTEIN Space.
Autor: Katsurada, Yoshie
DOI: <https://doi.org/10.5169/seals-29441>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On a Certain Property of Closed Hypersurfaces in an EINSTEIN Space

by YOSHIE KATSURADA, Sapporo

Introduction.

The following theorem, due to H. LIEBMANN (1900) [1], has been, and still is, the starting point of various interesting investigations within the Differential Geometry in the Large:

The only ovaloids with constant mean curvature H in EUCLIDEAN space E^3 are the spheres.

The analogous theorem for convex m -dimensional hypersurfaces in E^{m+1} has been proved by W. SÜSS (1929), [2], (cf. also [3], p. 118, and [4]). Recently (1958), A. D. ALEXANDROV has achieved the striking result that the convexity is not necessary for the validity of the LIEBMANN-SÜSS theorem, [5]: the theorem holds for arbitrary closed m -dimensional surfaces (hypersurfaces) without double points in E^{m+1} (i.e., for 1-1-embeddings of closed m -manifolds). Already previously (1951), H. HOPF had shown that, for $n = 2$ and for surfaces of genus 0, the theorem holds even without the hypothesis that there are no double points (i.e., it holds for immersions, not necessarily one-one, of 2-spheres into E^3), [6]. It remains an open question whether there exists an immersion, not one-one, of a closed surface of higher genus into E^3 such that $H = \text{constant}$.

There are also interesting investigations about generalizing the condition $H = \text{constant}$ in LIEBMANN's theorem. But we shall not discuss these problems here.

It is the aim of the present author to investigate the question whether the mentioned theorems, especially the LIEBMANN-SÜSS theorem, are special cases of theorems which hold in more general RIEMANN spaces. One step in this direction has already been made in a previous paper dealing with RIEMANN spaces with constant RIEMANN curvature [7]. The present paper deals with EINSTEIN spaces and generalizes the paper [7], without making use of it. Our result is Theorem 3.1 which, as is easily seen, contains the LIEBMANN-SÜSS theorem as special case (so does, by the way, also the main theorem of [7]).

It is well known that the LIEBMANN-SÜSS theorem is closely related to classical integral formulas of MINKOWSKI (cf. the paper of Süss). The base of our proof of Theorem 3.1 is a formula of MINKOWSKI type which holds in arbitrary RIEMANN spaces (formula (I) in § 1). This formula had already been established in [7]; a new proof is given in § 1 below. In § 2, some integral formulas for

hypersurfaces with $H = \text{constant}$ in EINSTEIN spaces are derived, and in § 3, the main theorem is proved.

The author wishes to express to Professor HEINZ HOPF her very sincere thanks for his valuable advice and suggestions.

§ 1. Another proof of the generalized MINKOWSKI formula (I).

In this section, we shall give a different proof of the generalized MINKOWSKI formula (I) derived in the previous paper ([7], p.288).

We consider a RIEMANN space R^{m+1} ($m + 1 \geq 3$) of class C^ν ($\nu \geq 3$) which admits an one-parameter continuous group G of transformations generated by an infinitesimal transformation

$$\bar{x}^i = x^i + \xi^i(x) \delta\tau \quad (1.1)$$

(where x^i are local coordinates in R^{m+1} and ξ^i are the components of a contra-variant vector ξ). We suppose that the paths of these transformations cover R^{m+1} simply and that ξ is everywhere continuous and $\neq 0$. If ξ is a KILLING vector, a homothetic KILLING, a conformal KILLING vector etc. ([8], p.32), then the group G is called isometric, homothetic, conformal etc., respectively.

We now consider a closed orientable hypersurface V^m of class C^3 imbedded in R^{m+1} , locally given by

$$x^i = x^i(u^\alpha); \quad (1.2)$$

here and henceforth, Latin indices run from 1 to $m + 1$ and Greek indices from 1 to m .

To the vector ξ introduced above, there belongs a covariant vector $\bar{\xi}$ of V^m with the components

$$\bar{\xi}_\alpha = \frac{\partial x^i}{\partial u^\alpha} \xi_i$$

where ξ_i are the covariant components of ξ ; we shall compute its covariant derivatives along V^m : by virtue of the fact that the covariant derivatives of $\frac{\partial x^i}{\partial u^\alpha}$ are

$$\frac{\delta}{\partial u^\beta} \left(\frac{\partial x^i}{\partial u^\alpha} \right) = b_{\alpha\beta} n^i$$

where $b_{\alpha\beta}$ is the second fundamental tensor and n^i is the unit normal vector of V^m , we find

$$\bar{\xi}_{\alpha;\beta} = b_{\alpha\beta} n^i \xi_i + \frac{\partial x^i}{\partial u^\alpha} \frac{\partial x^j}{\partial u^\beta} \xi_{i;j} \quad (1.3)$$

(the symbol " $;$ " always means the covariant derivative).

Multiplying (1.3) by the contravariant metric tensor $g^{\alpha\beta}$ of V^m and contracting, we get

$$g^{\alpha\beta} \bar{\xi}_{\alpha;\beta} = m H_1 n^i \xi_i + \frac{1}{2} g^{\alpha\beta} \frac{\partial x^i}{\partial u^\alpha} \frac{\partial x^j}{\partial u^\beta} \cdot \mathcal{L}_\xi g_{ij}, \quad (1.4)$$

where H_1 is the first mean curvature $\frac{1}{m} g^{\alpha\beta} b_{\alpha\beta}$ of V^m and $\mathcal{L}_\xi g_{ij}$ is the LIE derivative of the fundamental tensor g_{ij} of R^{m+1} with respect to the infinitesimal transformation (1.1) (cf. [8], p.5). If we put

$$\frac{\partial x^i}{\partial u^\alpha} \frac{\partial x^j}{\partial u^\beta} \mathcal{L}_\xi g_{ij} = \mathcal{L}_\xi g_{\alpha\beta}$$

then (1.4) rewritten is as follows:

$$\frac{1}{m} \bar{\xi}^\alpha_{;\alpha} = H_1 n^i \xi_i + \frac{1}{2m} g^{\alpha\beta} \mathcal{L}_\xi g_{\alpha\beta}.$$

dA being the area element of V^m , there holds

$$\int \dots \int_{V^m} \bar{\xi}^\alpha_{;\alpha} dA = 0$$

because V^m is closed and orientable ([9], p.31). Thus we obtain the integral formula

$$\int \dots \int_{V^m} H_1 n^i \xi_i dA + \frac{1}{2m} \int \dots \int_{V^m} g^{\alpha\beta} \mathcal{L}_\xi g_{\alpha\beta} dA = 0 \quad (I)$$

which is nothing but the formula (I) of the previous paper [7], p.288.

Let the group G be conformal, that is, ξ^i satisfy the equation

$$\mathcal{L}_\xi g_{ij} \equiv \xi_{i;j} + \xi_{j;i} = 2\Phi g_{ij}$$

(cf. [8], p.32), then (I) becomes

$$\int \dots \int_{V^m} H_1 n^i \xi_i dA + \int \dots \int_{V^m} \Phi dA = 0; \quad (I)_c$$

let G be homothetic, that is, $\Phi \equiv C = \text{constant}$, then

$$\int \dots \int_{V^m} H_1 n^i \xi_i dA + C \int \dots \int_{V^m} dA = 0; \quad (I)_h$$

and let G be isometric, that is, $C = 0$, then

$$\int \dots \int_{V^m} H_1 n^i \xi_i dA = 0. \quad (I)_i$$

§ 2. Some integral formulas for a closed hypersurface with $H_1 = \text{constant}$ in an EINSTEIN space.

Hereafter we shall assume that the RIEMANN space R^{m+1} is an EINSTEIN space and V^m is a closed orientable hypersurface with $H_1 = \text{constant}$.

If we take the covariant vector of the hypersurface V^m , defined by

$$\eta_\alpha = n^i{}_{;\alpha} \xi_i$$

and calculate its covariant derivatives along V^m , we have

$$\eta_{\alpha;\beta} = n^i{}_{;\alpha;\beta} \xi_i + n^i{}_{;\alpha} \xi_{i;j} \frac{\partial x^j}{\partial u^\beta}.$$

Remembering the following formulas for hypersurfaces

$$n^i{}_{;\alpha} = -b_\alpha^\gamma \frac{\partial x^i}{\partial u^\gamma}, \quad (2.1)$$

$$\frac{\delta}{\partial u^\beta} \left(\frac{\partial x^i}{\partial u^\gamma} \right) = b_{\beta\gamma} n^i, \quad (2.2)$$

where b_α^γ means $g^{\gamma\delta} b_{\alpha\delta}$ ([10] p. 136, p. 127), we find that

$$\eta_{\alpha;\beta} = - \left(\xi_i b_{\alpha;\beta}^\gamma \frac{\partial x^i}{\partial u^\gamma} + b_\alpha^\gamma b_{\gamma\beta} \xi_i n^i + \xi_{i;j} b_\alpha^\gamma \frac{\partial x^i}{\partial u^\gamma} \frac{\partial x^j}{\partial u^\beta} \right).$$

Multiplying by $g^{\alpha\beta}$ and contracting, we obtain

$$g^{\alpha\beta} \eta_{\alpha;\beta} = - g^{\alpha\beta} \left(\xi_i b_{\alpha;\beta}^\gamma \frac{\partial x^i}{\partial u^\gamma} + b_\alpha^\gamma b_{\gamma\beta} \xi_i n^i + \xi_{i;j} b_\alpha^\gamma \frac{\partial x^i}{\partial u^\gamma} \frac{\partial x^j}{\partial u^\beta} \right). \quad (2.3)$$

We shall first calculate the first term of the right-hand side of (2.3):

$$g^{\alpha\beta} \xi_i b_{\alpha;\beta}^\gamma \frac{\partial x^i}{\partial u^\gamma} = g^{\alpha\beta} g^{\gamma\delta} b_{\alpha\delta;\beta} \xi_i \frac{\partial x^i}{\partial u^\gamma}. \quad (2.4)$$

As well-known, an hypersurface in a RIEMANN space has the following property

$$b_{\alpha\delta;\beta} - b_{\alpha\beta;\delta} = - R_{ijkl} \frac{\partial x^i}{\partial u^\alpha} n^j \frac{\partial x^k}{\partial u^\delta} \frac{\partial x^l}{\partial u^\beta}, \quad ([10] \text{ p. 138})$$

where R_{ijkl} is the curvature tensor of R^{m+1} . Multiplying both sides of this equation by $g^{\alpha\beta}$ and contracting, we get

$$g^{\alpha\beta} (b_{\alpha\delta;\beta} - b_{\alpha\beta;\delta}) = - R_{ijkl} \frac{\partial x^i}{\partial u^\alpha} n^j \frac{\partial x^k}{\partial u^\delta} \frac{\partial x^l}{\partial u^\beta} g^{\alpha\beta}; \quad (2.5)$$

substituting

$$\frac{\partial x^i}{\partial u^\alpha} \frac{\partial x^l}{\partial u^\beta} g^{\alpha\beta} = g^{il} - n^i n^l$$

into the right-hand side of (2.5), we obtain

$$g^{\alpha\beta}(b_{\alpha\delta;\beta} - b_{\alpha\beta;\delta}) = -R_{jk}n^j \frac{\partial x^k}{\partial u^\delta} \quad (2.6)$$

R_{jk} being the Ricci tensor of R^{m+1} ($R_{jk} = g^{il}R_{ijkl}$). Because R^{m+1} is an EINSTEIN space and V^m has the property $H_1 = \text{constant}$, the right-hand side of (2.6) and the second term of the left-hand side vanish; it follows that

$$g^{\alpha\beta}b_{\alpha\delta;\beta} = 0. \quad (2.7)$$

Therefore, and with respect to (2.4), the first term of the right-hand side of (2.3) is equal to zero.

Next, we discuss the second term of the right-hand side of (2.3):

$$g^{\alpha\beta}b_\alpha^\gamma b_{\gamma\beta}n^i\xi_i = g^{\alpha\beta}g^{\gamma\delta}b_{\alpha\delta}b_{\gamma\beta}n^i\xi_i. \quad (2.8)$$

Let k_1, k_2, \dots, k_m be the principal curvatures at a point P of V^m , and let H_2 be the second mean curvature of V^m at the point P which is defined to be the second elementary symmetric function of k_1, k_2, \dots, k_m divided by the number of terms, that is,

$$\binom{m}{2} H_2 = \sum_{(\alpha, \beta)} k_\alpha k_\beta \quad (\alpha < \beta);$$

since, furthermore the following relation holds

$$\frac{1}{2}(g^{\alpha\delta}g^{\gamma\beta}b_{\alpha\delta}b_{\gamma\beta} - g^{\alpha\beta}g^{\gamma\delta}b_{\alpha\delta}b_{\gamma\beta}) = \binom{m}{2} H_2,$$

(2.8) can be written as follows

$$g^{\alpha\beta}b_\alpha^\gamma b_{\gamma\beta}n^i\xi_i = \{m^2 H_1^2 - 2 \binom{m}{2} H_2\} n^i\xi_i. \quad (2.9)$$

At last, for the third term of the right-hand side of (2.3), we calculate as follows

$$\begin{aligned} g^{\alpha\beta}b_\alpha^\gamma \frac{\partial x^i}{\partial u^\gamma} \frac{\partial x^j}{\partial u^\beta} \xi_{i;j} &= g^{\alpha\beta}g^{\gamma\delta}b_{\alpha\delta} \frac{\partial x^i}{\partial u^\gamma} \frac{\partial x^j}{\partial u^\beta} \xi_{i;j} \\ &= \frac{1}{2}g^{\alpha\beta}g^{\gamma\delta}b_{\alpha\delta} \frac{\partial x^i}{\partial u^\gamma} \frac{\partial x^j}{\partial u^\beta} (\xi_{i;j} + \xi_{j;i}) \\ &= \frac{1}{2}H^{\beta\gamma} \mathcal{L}_\xi g_{\beta\gamma} \end{aligned} \quad (2.10)$$

where $H^{\beta\gamma}$ denotes $b_{\alpha\delta}g^{\alpha\beta}g^{\gamma\delta}$.

Accordingly, from (2.7), (2.9), and (2.10), (2.3) becomes

$$\frac{1}{m}\eta^\alpha_{;\alpha} = -\{(mH_1^2 - (m-1)H_2)n^i\xi_i + \frac{1}{2m}H^{\beta\gamma} \mathcal{L}_\xi g_{\beta\gamma}\}.$$

And also on making use of

$$\int \dots \int_{V^m} \eta^\alpha_{;\alpha} dA = 0$$

by virtue of V^m being closed orientable, we finally reach the integral formula

$$\int \dots \int_{V^m} \{mH_1^2 - (m-1)H_2\} n^i \xi_i dA + \frac{1}{2m} \int \dots \int_{V^m} H^{\alpha\beta} \mathcal{L}_{\xi} g_{\alpha\beta} dA = 0. \quad (\text{II})$$

If the group G of transformation is conformal, (II) becomes

$$\int \dots \int_{V^m} \{mH_1^2 - (m-1)H_2\} n^i \xi_i dA + \int \dots \int_{V^m} \Phi H_1 dA = 0; \quad (\text{II})_c$$

if G is homothetic (i.e. $\Phi \equiv \text{constant} = C$),

$$\int \dots \int_{V^m} \{mH_1^2 - (m-1)H_2\} n^i \xi_i dA + C \int \dots \int_{V^m} H_1 dA = 0; \quad (\text{II})_h$$

and if G is isometric (i.e. $\Phi \equiv 0$),

$$\int \dots \int_{V^m} \{mH_1^2 - (m-1)H_2\} n^i \xi_i dA = 0. \quad (\text{II})_i$$

§ 3. Closed orientable hypersurfaces with $H_1 = \text{constant}$ in an EINSTEIN space.

In this section, we shall prove the following theorem:

Theorem 3.1. *Let R^{m+1} be an EINSTEIN space, V^m a closed orientable hypersurface with $H_1 = \text{constant}$ in R^{m+1} ; we suppose that there exists a continuous one-parameter group G of conformal transformations of R^{m+1} such that the scalar product $\tilde{p} = n^i \xi_i$ of the normal vector n of V^m and the vector ξ belonging to G does not change the sign (and is not $\equiv 0$) on V^m . Then every point of V^m is umbilic.*

Proof. Multiplying the formula (I)_c in § 1 by $H_1 (= \text{const.})$, we obtain

$$\int \dots \int_{V^m} H_1^2 \tilde{p} dA + \int \dots \int_{V^m} \Phi H_1 dA = 0,$$

and subtracting this formula from the formula (II)_c in § 2, we find

$$\int \dots \int_{V^m} (m-1)(H_1^2 - H_2) \tilde{p} dA = 0. \quad (3.1)$$

From

$$H_1^2 - H_2 = \frac{1}{m^2} \cdot (\sum k_\alpha)^2 - \frac{2}{m(m-1)} \sum_{\alpha, \beta} k_\alpha k_\beta = \frac{1}{m^2(m-1)} \sum (k_\alpha - k_\beta)^2 \quad (3.2)$$

(with $\alpha \neq \beta$) we see that

$$H_1^2 - H_2 \geq 0. \quad (3.3)$$

From (3.1), (3.3) and the fact that \tilde{p} has a fixed sign we conclude that

$$H_1^2 - H_2 = 0$$

and therefore, because of (3.2), that

$$k_1 = k_2 = \dots = k_m$$

at each point of V^m . This means, that each point of V^m is umbilic.

We wish now to show that the LIEBMANN-SÜSS Theorem is a special case of our Theorem 3.1. Because in euclidean E^{m+1} an hypersurface is a sphere if all its points are umbilical we have only to verify that, for a convex V^m in E^{m+1} , there exists a vector field ξ having the properties formulated in Theorem 3.1. We take a point in the interior of V^m as origin of the euclidean coordinates x^i and attach to each point x the vector $\xi(x)$ with the components $\xi^i = x^i$ (i.e. the position vector of x). Then, the transformations (1.1) are homothetic, thus conformal; furthermore, for $x \in V^m$, $\tilde{p}(x)$ is the support function and, because V^m is convex, $\tilde{p}(x) \neq 0$.

REFERENCES

- [1] H. LIEBMANN: *Über die Verbiegung der geschlossenen Flächen positiver Krümmung*, Math. Ann. 53 (1900), 91–112.
- [2] W. SÜSS: *Zur relativen Differentialgeometrie V.*, Tôhoku Math. J. 30 (1929), 202–209.
- [3] T. BONNESEN und W. FENCHEL: *Theorie der konvexen Körper* (Springer, Berlin 1934).
- [4] C. C. HSIUNG: *Some integral formulas for closed hypersurfaces*, Math. Scand. 2 (1954), 286–294.
- [5] A. D. ALEXANDROV: *Uniqueness theorems for surfaces in the large*, V. Vestnik Leningrad University 13 (1958), 5–8 (Russian, with English summary).
- [6] H. HOPF: *Über Flächen mit einer Relation zwischen den Hauptkrümmungen*, Math. Nachr. 4 (1951), 232–249.
- [7] Y. KATSURADA: *Generalized Minkowski formulas for closed hypersurfaces in a RIEMANN space*, Annali di Matematica, Serie IV, 57 (1962), 283–294.
- [8] K. YANO: *The theory of LIE derivatives and its applications* (Amsterdam 1957).
- [9] K. YANO and S. BOCHNER: *Curvature and Betti numbers* (Princeton, Annals of Math. Studies, 1953).
- [10] C. E. WEATHERBURN: *An introduction to RIEMANN geometry and the tensor calculus* (Cambridge University Press 1938).

(Received May 12, 1963)